

(19) World Intellectual Property  
Organization  
International Bureau



(43) International Publication Date  
17 February 2005 (17.02.2005)

PCT

(10) International Publication Number  
WO 2005/014650 A2

(51) International Patent Classification<sup>7</sup>: C07K 16/22,  
A61K 39/395, C07K 14/51, G01N 33/53, A61P 19/10

WA 98115 (US). LATHAM, John [US/US]; 2409 10th Avenue West, Seattle, WA 98119 (US).

(21) International Application Number:  
PCT/US2004/018912

(74) Agents: POTTER, Jane, E., R. et al.; 2600 Century Square, 1501 Fourth Avenue, Seattle, WA 98101-1688 (US).

(22) International Filing Date: 15 June 2004 (15.06.2004)

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
60/478,977 16 June 2003 (16.06.2003) US

(71) Applicant (for all designated States except US): CELL-TECH R & D, INC. [US/US]; 1631 220th Street SE, Bothell, WA 98021 (US).


(72) Inventors; and

(75) Inventors/Applicants (for US only): WINKLER, David, G. [US/US]; 7037 20th Avenue NE, Seattle, WA 98115 (US). SHI, Jiye [CN/US]; 9115 32nd Avenue NE, Seattle,

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI,

[Continued on next page]

(54) Title: ANTIBODIES SPECIFIC FOR SCLEROSTIN AND METHODS FOR INCREASING BONE MINERALIZATION



(57) Abstract: Compositions and methods relating to antibodies that specifically bind to TGF-beta binding proteins are provided. These methods and compositions relate to altering bone mineral density by interfering with the interaction between a TGF-beta binding protein sclerostin and a TGF-beta superfamily member, particularly a bone morphogenic protein. Increasing bone mineral density has uses in diseases and conditions in which low bone mineral density typifies the condition, such as osteopenia, osteoporosis, and bone fractures.

WO 2005/014650 A2



SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

*For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*

**Published:**

- *without international search report and to be republished upon receipt of that report*

## ANTIBODIES SPECIFIC FOR SCLEROSTIN AND METHODS FOR INCREASING BONE MINERALIZATION

### TECHNICAL FIELD

The present invention relates generally to pharmaceutical products and methods  
5 and, more specifically, to methods and sclerostin-specific antibodies suitable for increasing the mineral content of bone. Such compositions and methods may be used to treat a wide variety of conditions including, for example, osteopenia, osteoporosis, fractures, and other disorders in which low bone mineral density is a hallmark of the disease.

### BACKGROUND OF THE INVENTION

10 Two or three distinct phases of changes to bone mass occur over the life of an individual (*see* Riggs, *West J. Med.* 154:63-77 (1991)). The first phase occurs in both men and women and proceeds to attainment of a peak bone mass. This first phase is achieved through linear growth of the endochondral growth plates and radial growth due to a rate of periosteal apposition. The second phase begins around age 30 for trabecular bone (flat bones such as the  
15 vertebrae and pelvis) and about age 40 for cortical bone (*e.g.*, long bones found in the limbs) and continues to old age. This phase is characterized by slow bone loss and occurs in both men and women. In women, a third phase of bone loss also occurs, most likely due to postmenopausal estrogen deficiencies. During this phase alone, women may lose an additional 10% of bone mass from the cortical bone and 25% from the trabecular compartment (*see* Riggs, *supra*).

20 Loss of bone mineral content can be caused by a wide variety of conditions and may result in significant medical problems. For example, osteoporosis is a debilitating disease in humans and is characterized by marked decreases in skeletal bone mass and mineral density, structural deterioration of bone, including degradation of bone microarchitecture and corresponding increases in bone fragility, and susceptibility to fracture in afflicted individuals.  
25 Osteoporosis in humans is preceded by clinical osteopenia (bone mineral density that is greater than one standard deviation but less than 2.5 standard deviations below the mean value for young adult bone), a condition found in approximately 25 million people in the United States. Another 7-8 million patients in the United States have been diagnosed with clinical osteoporosis (defined as bone mineral content greater than 2.5 standard deviations below that of mature  
30 young adult bone). Osteoporosis is one of the most expensive diseases for the health care

system, costing tens of billions of dollars annually in the United States. In addition to health care-related costs, long-term residential care, and lost working days add to the financial and social costs of this disease. Worldwide, approximately 75 million people are at risk for osteoporosis.

5           The frequency of osteoporosis in the human population increases with age. Among Caucasians, osteoporosis is predominant in women who, in the United States, comprise 80% of the osteoporosis patient pool. The increased fragility and susceptibility to fracture of skeletal bone in the aged is aggravated by the greater risk of accidental falls in this population. More than 1.5 million osteoporosis-related bone fractures are reported in the United States each  
10   year. Fractured hips, wrists, and vertebrae are among the most common injuries associated with osteoporosis. Hip fractures in particular are extremely uncomfortable and expensive for the patient, and for women, correlate with high rates of mortality and morbidity.

          Although osteoporosis has been regarded as an increase in the risk of fracture due to decreased bone mass, none of the presently available treatments for skeletal disorders can  
15   substantially increase the bone density of adults. A strong perception among many physicians is that drugs are needed that could increase bone density in adults, particularly in the bones of the wrist, spinal column, and hip that are at risk in osteopenia and osteoporosis.

          Current strategies for the prevention of osteoporosis may offer some benefit to individuals but cannot ensure resolution of the disease. These strategies include moderating  
20   physical activity (particularly in weight-bearing activities) with the onset of advanced age, providing adequate calcium in the diet, and avoiding consumption of products containing alcohol or tobacco. For patients presenting clinical osteopenia or osteoporosis, the prevalent current therapeutic drugs and strategies are directed to reducing further loss of bone mass by inhibiting the process of bone absorption, a natural aspect of the bone remodeling process that  
25   occurs constitutively.

          For example, estrogen is now being prescribed to retard bone loss. However, some controversy exists over whether patients gain any long-term benefit and whether estrogen has any effect at all on patients over 75 years old. Moreover, use of estrogen is believed to increase the risk of breast and endometrial cancer. Calcitonin, osteocalcin with vitamin K, or  
30   high doses of dietary calcium, with or without vitamin D, have also been suggested for postmenopausal women. High doses of calcium, however, can often have unpleasant gastrointestinal side effects, and serum and urinary calcium levels must be continuously monitored (e.g., Khosla and Riggs, *Mayo Clin. Proc.* 70:978-982, 1995).

Other therapeutic approaches to osteoporosis include bisphosphonates (e.g., Fosamax™, Actonel™, Bonviva™, Zometa™, olpadronate, neridronate, skelid, bonefos), parathyroid hormone, calcilytics, calcimimetics (e.g., cinacalcet), statins, anabolic steroids, lanthanum and strontium salts, and sodium fluoride. Such therapeutics, however, are often  
5 associated with undesirable side effects (for example, calcitonin and steroids may cause nausea and provoke an immune reaction, bisphosphonates and sodium fluoride may inhibit repair of fractures, even though bone density increases modestly) that may preclude their efficacious use (see Khosla and Riggs, *supra*).

No currently practiced therapeutic strategy for treating a condition associated  
10 with excessive or insufficient bone mineralization, such as osteoporosis or other disorders characterized by loss of bone mineralization, involves a drug that alters (*i.e.*, increases or decreases in a statistically significant manner) bone mass. In particular, no current strategy therapeutically stimulates or enhances the growth of new bone mass. The present invention provides compositions and methods which can be used to increase bone mineralization, and  
15 which therefore may be used to treat a wide variety of conditions in which an increase in bone mass is desirable. The present invention also offers other related advantages.

#### BRIEF SUMMARY OF THE INVENTION

Briefly stated, the present invention provides antibodies that specifically bind to a TGF-beta binding protein, sclerostin (SOST), and provides immunogens comprising SOST  
20 peptides derived from regions of SOST that interact with a member of the TGF-beta superfamily such as a bone morphogenic protein. In one embodiment, the invention provides an isolated antibody, or an antigen-binding fragment thereof, that binds specifically to a SOST polypeptide, which comprises an amino acid sequence set forth in SEQ ID NO:1, 20, 58, 60, 62, or 68, wherein the antibody competitively inhibits binding of the SOST polypeptide to at least one of  
25 (i) a bone morphogenic protein (BMP) Type I Receptor binding site and (ii) a BMP Type II Receptor binding site, wherein the BMP Type I Receptor binding site is capable of binding to a BMP Type I Receptor polypeptide comprising an amino acid sequence set forth in GenBank Acc. Nos. NM\_004329 (SEQ ID NO:71); D89675 (SEQ ID NO:72); NM\_001203 (SEQ ID NO:73); S75359 (SEQ ID NO:74); NM\_030849 (SEQ ID NO:75); D38082 (SEQ ID NO:76);  
30 NP\_001194 (SEQ ID NO:77); BAA19765 (SEQ ID NO:78); or AAB33865 (SEQ ID NO:79), and wherein the BMP Type II Receptor binding site is capable of binding to a BMP Type II Receptor polypeptide comprising the amino acid sequence set forth in GenBank Acc. NOs.

U25110 (SEQ ID NO:80); NM\_033346 (SEQ ID NO:81); Z48923 (SEQ ID NO:83); CAA88759 (SEQ ID NO:84); or NM\_001204 (SEQ ID NO:82). In a particular embodiment, the antibody, or the antigen-binding fragment thereof, binds specifically to a polypeptide that comprises an amino acid sequence set forth in SEQ ID NO:2-6, 12-15, 21, 22, 25, 26, 47, 48, 49, 50, and in other particular embodiments, the antibody binds specifically to a polypeptide that comprises an amino acid sequence set forth in SEQ ID NO: 5, 6, or 14.

In one embodiment, the invention provides an isolated antibody, or an antigen-binding fragment thereof, that is produced by immunizing a non-human animal with a peptide of at least 20 amino acids and no more than 75 amino acids that comprises an amino acid sequence of SEQ ID NOs:2-19, 21-28, or 47-50, wherein the antibody binds specifically to a SOST polypeptide, which comprises an amino acid sequence set forth in SEQ ID NOs:1, 20, 58, 60, 62, or 68, and wherein the antibody competitively inhibits binding of the SOST polypeptide to at least one of (i) a bone morphogenic protein (BMP) Type I Receptor binding site and (ii) a BMP Type II Receptor binding site, wherein the BMP Type I Receptor binding site is capable of binding to a BMP Type I Receptor polypeptide (set forth in any one of GenBank sequences provided herein), and wherein the BMP Type II Receptor binding sites is capable of binding to a BMP Type II Receptor polypeptide (set forth in any one of GenBank sequences provided herein). In certain preferred embodiments, an antibody is produced by immunizing a non-human animal with a peptide of at least 20 amino acids and no more than 75 amino acids that comprises an amino acid sequence of SEQ ID NO:5, 6, 10, 11, 14, or 18.

The present invention further provides an antibody, or an antigen-binding fragment thereof, that binds specifically to a SOST polypeptide and that impairs formation of a SOST homodimer, wherein the SOST polypeptide comprises an amino acid sequence set forth in SEQ ID NOs:1, 20, 58, 60, 62, or 68, and wherein the antibody binds to a polypeptide that comprises an amino acid sequence of SEQ ID NOs:29-31, 33-36, 41-43, or 51-54.

In another embodiment, the invention provides an isolated antibody, or an antigen-binding fragment thereof, that is produced by immunizing a non-human animal with a peptide of at least 20 amino acids and no more than 75 amino acids comprising an amino acid sequence of SEQ ID NOs: 29-46, 51, 52, 53, or 54, wherein the antibody, or the antigen-binding fragment thereof, binds specifically to a SOST polypeptide that comprises an amino acid sequence set forth in SEQ ID NO:1, 20, 58, 60, 62, or 68, and wherein the antibody impairs formation of a SOST homodimer.

In certain particular embodiments of the invention, the antibody is a polyclonal

antibody. In other embodiments, the antibody is a monoclonal antibody, which is a mouse, human, rat, or hamster monoclonal antibody. The invention also provides a hybridoma cell or a host cell that is capable of producing the monoclonal antibody. In other embodiments of the invention, the antibody is a humanized antibody or a chimeric antibody. The invention further provides a host cell that produces the humanized or chimeric antibody. In certain embodiments the antigen-binding fragment of the antibody is a F(ab')<sub>2</sub>, Fab', Fab, Fd, or Fv fragment. The invention also provides an antibody that is a single chain antibody and provides a host cell that is capable of expressing the single chain antibody. In another embodiment, the invention provides a composition comprising such antibodies and a physiologically acceptable carrier.

The invention provides an immunogen comprising a peptide comprising 6, 7, 8, 9, 10, 11, or 12 consecutive amino acids of a SOST polypeptide, which SOST polypeptide comprises an amino acid sequence set forth in SEQ ID NOs:1, 20, 58, 60, 62, or 68, and wherein the peptide is capable of eliciting in a non-human animal an antibody that binds specifically to the SOST polypeptide and that competitively inhibits binding of the SOST polypeptide to at least one of (i) a bone morphogenic protein (BMP) Type I Receptor binding site and (ii) a BMP Type II Receptor binding site, wherein the BMP Type I Receptor binding site is capable of binding to a BMP Type I Receptor polypeptide, wherein wherein the BMP Type I Receptor binding site is capable of binding to a BMP Type I Receptor polypeptide comprising an amino acid sequence set forth in GenBank Acc. Nos. NM\_004329 (SEQ ID NO:71); D89675 (SEQ ID NO:72); NM\_001203 (SEQ ID NO:73); S75359 (SEQ ID NO:74); NM\_030849 (SEQ ID NO:75); D38082 (SEQ ID NO:76); NP\_001194 (SEQ ID NO:77); BAA19765 (SEQ ID NO:78); or AAB33865 (SEQ ID NO:79), and wherein the BMP Type II Receptor binding site is capable of binding to a BMP Type II Receptor polypeptide comprising the amino acid sequence set forth in GenBank Acc. NOs. U25110 (SEQ ID NO:80); NM\_033346 (SEQ ID NO:81); Z48923 (SEQ ID NO:83); CAA88759 (SEQ ID NO:84); or NM\_001204 (SEQ ID NO:82).

In another embodiment, an immunogen comprises a peptide comprising at least 21 consecutive amino acids and no more than 50 consecutive amino acids of a SOST polypeptide, said SOST polypeptide comprising an amino acid sequence set forth in SEQ ID NOs:1, 20, 58, 60, 62, or 68, wherein the peptide is capable of eliciting in a non-human animal an antibody that binds specifically to the SOST polypeptide and that competitively inhibits binding of the SOST polypeptide to at least one of (i) a bone morphogenic protein (BMP) Type I Receptor binding site and (ii) a BMP Type II Receptor binding site, wherein the BMP Type I Receptor binding site is capable of binding to a BMP Type I Receptor polypeptide (set forth in

any one of GenBank sequences provided herein), and wherein the BMP Type II Receptor binding site is capable of binding to a BMP Type II Receptor polypeptide (set forth in any one of GenBank sequences provided herein).

In certain embodiments the invention provides an immunogen comprising a peptide of at least 20 amino acids and no more than 75 amino acids comprising an amino acid sequence of SEQ ID NOs: 2-19, 21-28, 47, 48, 49, or 50, wherein the peptide is capable of eliciting in a non-human animal an antibody that binds specifically to a SOST polypeptide, which comprises an amino acid sequence set forth in SEQ ID NO:1, 20, 58, 60, 62, or 68, wherein the antibody competitively inhibits binding of the SOST polypeptide to at least one of (i) a bone morphogenic protein (BMP) Type I Receptor binding site and (ii) a BMP Type II Receptor binding site, wherein the BMP Type I Receptor binding site is capable of binding to a BMP Type I Receptor polypeptide comprising an amino acid sequence set forth in GenBank Acc. Nos. NM\_004329 (SEQ ID NO:71); D89675 (SEQ ID NO:72); NM\_001203 (SEQ ID NO:73); S75359 (SEQ ID NO:74); NM\_030849 (SEQ ID NO:75); D38082 (SEQ ID NO:76); NP\_001194 (SEQ ID NO:77); BAA19765 (SEQ ID NO:78); or AAB33865 (SEQ ID NO:79), and wherein the BMP Type II Receptor binding site is capable of binding to a BMP Type II Receptor polypeptide comprising the amino acid sequence set forth in GenBank Acc. NOs. U25110 (SEQ ID NO:80); NM\_033346 (SEQ ID NO:81); Z48923 (SEQ ID NO:83); CAA88759 (SEQ ID NO:84); or NM\_001204 (SEQ ID NO:82). In a particular embodiment, the invention provides an immunogen comprising a peptide of at least 20 amino acids and no more than 75 amino acids comprising an amino acid sequence of SEQ ID NO:5, 6, 10, 11, 14, or 18.

The invention also provides an immunogen comprising a peptide of at least 20 amino acids and no more than 75 amino acids comprising an amino acid sequence of SEQ ID NOs: 29-46, 51, 52, 53, or 54, wherein the peptide is capable of eliciting in a non-human animal an antibody that binds specifically to a SOST polypeptide, which comprises an amino acid sequence set forth in SEQ ID NO:1, 20, 58, 60, 62, or 68, and wherein the antibody impairs formation of a SOST homodimer. In another embodiment, an immunogen comprises a peptide comprising 6, 7, 8, 9, 10, 11, or 12 consecutive amino acids of a SOST polypeptide, wherein the SOST polypeptide comprises an amino acid sequence set forth in SEQ ID NO:1, 20, 58, 60, 62, or 68, and wherein the peptide is capable of eliciting in a non-human animal an antibody that binds specifically to the SOST polypeptide and that impairs formation of a SOST homodimer. In certain embodiments, the invention provides an immunogen that comprises a peptide



comprising at least 21 consecutive amino acids and no more than 50 consecutive amino acids of a SOST polypeptide, which polypeptide comprises SEQ ID NO:1, 20, 58, 60, 62, or 68, and wherein the peptide is capable of eliciting in a non-human animal an antibody that binds specifically to the SOST polypeptide and that impairs formation of a SOST homodimer.

5               In certain particular embodiments, the subject invention immunogens are associated with a carrier molecule. In certain embodiments, the carrier molecule is a carrier polypeptide, and in particular embodiments, the carrier polypeptide is keyhole limpet hemocyanin.

              The present invention also provides a method for producing an antibody that  
10       specifically binds to a SOST polypeptide, comprising immunizing a non-human animal with an immunogen comprising a peptide of 6, 7, 8, 9, 10, 11, or 12 consecutive amino acids or at least 21 and no more than 50 consecutive amino acids of a SOST polypeptide having the sequence set forth in SEQ ID NO:1, 20, 58, 60, 62, or 68, that is capable of eliciting in a non-human animal an antibody that binds specifically to the SOST polypeptide and that competitively inhibits  
15       binding of the SOST polypeptide to at least one of (i) a bone morphogenic protein (BMP) Type I Receptor binding site and (ii) a BMP Type II Receptor binding site, wherein the BMP Type I Receptor binding site is capable of binding to a BMP Type I Receptor polypeptide, wherein the BMP Type I Receptor binding site is capable of binding to a BMP Type I Receptor polypeptide comprising an amino acid sequence set forth in GenBank Acc. Nos. NM\_004329 (SEQ ID  
20       NO:71); D89675 (SEQ ID NO:72); NM\_001203 (SEQ ID NO:73); S75359 (SEQ ID NO:74); NM\_030849 (SEQ ID NO:75); D38082 (SEQ ID NO:76); NP\_001194 (SEQ ID NO:77); BAA19765 (SEQ ID NO:78); or AAB33865 (SEQ ID NO:79), and wherein the BMP Type II Receptor binding site is capable of binding to a BMP Type II Receptor polypeptide comprising the amino acid sequence set forth in GenBank Acc. NOs. U25110 (SEQ ID NO:80);  
25       NM\_033346 (SEQ ID NO:81); Z48923 (SEQ ID NO:83); CAA88759 (SEQ ID NO:84); or NM\_001204 (SEQ ID NO:82). In another embodiment, the present invention provides a method for producing an antibody that specifically binds to a SOST polypeptide, comprising immunizing a non-human animal with an immunogen comprising a peptide at least 20 amino acids and no more than 75 amino acids comprising an amino acid sequence of SEQ ID NO: 2-  
30       19, 21-28, 47, 48, 49, or 50, wherein the peptide is capable of eliciting in a non-human animal an antibody that binds specifically to a SOST polypeptide, which comprises an amino acid sequence set forth in SEQ ID NO:1, 20, 58, 60, 62, or 68, wherein the antibody competitively inhibits binding of the SOST polypeptide to at least one of (i) a bone morphogenic protein

(BMP) Type I Receptor binding site and (ii) a BMP Type II Receptor binding site, wherein the BMP Type I Receptor binding site is capable of binding to a BMP Type I Receptor polypeptide. In a particular embodiment, the immunogen comprises a peptide at least 20 amino acids and no more than 75 amino acids comprising an amino acid sequence of SEQ ID NO:5, 6, 10, 11, 14, or 18.

In other preferred embodiments, the invention provides a method for producing an antibody that specifically binds to a SOST polypeptide, comprising immunizing a non-human animal with an immunogen comprising a peptide of 6, 7, 8, 9, 10, 11, or 12 consecutive amino acids or at least 21 and no more than 50 consecutive amino acids of a SOST polypeptide having the sequence set forth in SEQ ID NO:1, 20, 58, 60, 62, or 68, that is capable of eliciting in a non-human animal an antibody that binds specifically to the SOST polypeptide and that is capable of impairing formation of a SOST homodimer. In another embodiment, the invention provides a method for producing an antibody that specifically binds to a SOST polypeptide, comprising immunizing a non-human animal with an immunogen comprising a peptide at least 20 amino acids and no more than 75 amino acids comprising an amino acid sequence of SEQ ID NO: 29-46, 51, 52, 53, or 54, wherein the peptide is capable of eliciting in a non-human animal an antibody that binds specifically to a SOST polypeptide and impairs formation of a SOST homodimer.

The present invention also provides a method for identifying an antibody that modulates a TGF-beta signaling pathway, comprising contacting an antibody that specifically binds to a SOST polypeptide which comprises an amino acid sequence set forth in any one of SEQ ID NOS: 1, 20, 58, 60, 62, or 68 with at least one SOST peptide comprising an amino acid sequence of SEQ ID NO: 2-19, or 21-54, under conditions and for a time sufficient to permit formation of an antibody/SOST peptide complex; and detecting a level of antibody/SOST peptide complex, and thereby detecting the presence of an antibody that modulates a TGF-beta signaling pathway. In another embodiment, the invention provides a method for identifying an antibody that impairs binding of a BMP to a SOST polypeptide, comprising contacting (i) an antibody that specifically binds to a SOST polypeptide which comprises an amino acid sequence set forth in any one of SEQ ID NOS: 1, 20, 58, 60, 62, or 68 with (ii) at least one SOST peptide comprising an amino acid sequence of SEQ ID NO: 2-19, 21-28, or 47-50 under conditions and for a time sufficient to permit formation of an antibody/SOST peptide complex; and detecting a level of antibody/SOST peptide complex, and thereby detecting the presence of an antibody that impairs binding of a BMP to a SOST polypeptide. In particular embodiments of the method for

identifying an antibody that modulates a TGF-beta signaling pathway and for identifying an antibody that impairs binding of a BMP to a SOST polypeptide, the SOST peptide comprises an amino acid sequence of SEQ ID NO:5, 6, 10, 11, 14, or 18.

5 In another embodiment, the invention provides a method for identifying an antibody that impairs SOST homodimer formation, comprising contacting (i) an antibody that specifically binds to a SOST polypeptide which comprises an amino acid sequence set forth in any one of SEQ ID NOS: 1, 20, 58, 60, 62, or 68, with (ii) at least one SOST peptide comprising an amino acid sequence of SEQ ID NO: 29-46 or 51-54 under conditions and for a time sufficient to permit formation of an antibody/SOST peptide complex; and detecting a level of  
10 antibody/SOST peptide complex, and thereby detecting the presence of an antibody that that impairs SOST homodimer formation.

In another embodiment, the invention provides a method for identifying an antibody that increases bone mineral content, comprising contacting (i) an antibody that specifically binds to a SOST polypeptide which comprises an amino acid sequence set forth in  
15 any one of SEQ ID NOS: 1, 20, 58, 60, 62, or 68 with (ii) at least one SOST peptide comprising an amino acid sequence of SEQ ID NO: 2-19, or 21-54 under conditions and for a time sufficient to permit formation of an antibody/SOST peptide complex; and detecting a level of antibody/SOST peptide complex, and thereby detecting the presence of an antibody that increases bone mineral content. In a particular embodiment, the peptide comprises an amino  
20 acid sequence of SEQ ID NO:5, 6, 10, 11, 14, or 18.

In certain particular embodiments for identifying an antibody that modulates a TGF-beta signaling pathway, for identifying an antibody that impairs binding of a BMP to a SOST polypeptide, for identifying an antibody that impairs SOST homodimer formation, or for identifying an antibody that increases bone mineral content, the antibody is present in a  
25 biological sample or the antibody is a purified antibody. In certain embodiments the purified antibody is a polyclonal antibody, a monoclonal antibody, a chimeric antibody, a humanized antibody, or an antigen-binding fragment of any of these antibodies.

These and other embodiments of the present invention will become apparent upon reference to the following detailed description and attached drawings. All references  
30 disclosed herein are hereby incorporated by reference in their entireties as if each was incorporated individually.

## BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 presents an alignment of the region containing the characteristic cystine-knot of a SOST polypeptide and its closest homologues. Three disulphide bonds that form the cystine-knot are illustrated as solid lines. An extra disulphide bond, shown by a dotted line, is unique to this family, which connects two  $\beta$ -hairpin tips in the 3D structure. The polypeptides depicted are SOST: sclerostin (SEQ ID NO:95); CGHB: Human Chorionic Gonadotropin  $\beta$  (SEQ ID NO:96); FSHB: follicle-stimulating hormone beta subunit (SEQ ID NO:97); TSHB: thyrotropin beta chain precursor (SEQ ID NO:98); VWF: Von Willebrand factor (SEQ ID NO:99); MUC2: human mucin 2 precursor (SEQ ID NO:100); CER1: Cerberus 1 (*Xenopus laevis* homolog) (SEQ ID NO:101); DRM: gremlin (SEQ ID NO:102); DAN: (SEQ ID NO:103); CTGF: connective tissue growth factor precursor (SEQ ID NO:104); NOV: NovH (nephroblastoma overexpressed gene protein homolog) (SEQ ID NO:105); CYR6: (SEQ ID NO:106).

Figure 2 illustrates a 3D model of the core region of SOST (SOST\_Core).

Figure 3 presents a 3D model of the core region of SOST homodimer.

Figure 4A and B provides an amino acid sequence alignment of Noggin from five different animals: human (NOGG\_HUMAN (SEQ ID NO:107); chicken (NOGG\_CHICK, SEQ ID NO:108); African clawed frog (NOGG\_XENLA, SEQ ID NO:109); NOGG\_FUGRU, SEQ ID NO:110); and zebrafish (NOGG\_ZEBRA, SEQ ID NO:111); and SOST from human (SOST\_HUMAN, SEQ ID NO:1), rat (SOST\_RAT, SEQ ID NO:20), and mouse (SOST\_Mouse, SEQ ID NO:112).

Figure 5 illustrates the Noggin/BMP-7 complex structure. The BMP homodimer is shown on the bottom portion of the figure in surface mode. The Noggin homodimer is shown on top of the BMP dimer in cartoon mode. The circles outline the N-terminal binding region, the core region, and the linker between the N-terminal and core regions.

Figure 6 depicts a 3D model of the potential BMP-binding fragment located at the SOST N-terminal region. A BMP dimer is shown in surface mode, and the potential BMP-binding fragment is shown in stick mode. A phenylalanine residue fitting into a hydrophobic pocket on the BMP surface is noted.

## DETAILED DESCRIPTION OF THE INVENTION

The present invention provides antibodies that specifically bind to a SOST polypeptide methods for using such antibodies. The present invention also provides SOST polypeptide immunogens that may be used for generation and analysis of these antibodies. The

antibodies may be useful to block or impair binding of a SOST polypeptide, which is a TGF-beta binding protein, to a ligand, particularly a bone morphogenic protein, and may also block or impair binding of the SOST polypeptide to one or more other ligands.

The invention relates in part to the surprising discovery, within the sclerostin  
5 polypeptide sequence, of specific short peptide sequences that are specifically recognized by anti-sclerostin antibodies capable of competitively inhibiting binding of sclerostin polypeptides to BMP, where such sclerostin-BMP binding otherwise would occur via a BMP Type I Receptor binding site and/or a BMP Type II Receptor binding site. A molecule such as an antibody that inhibits the binding of the TGF-beta binding protein to one or more members of the TGF-beta  
10 family of proteins, including one or more bone morphogenic proteins (BMPs), should be understood to include, for example, a molecule that allows the activation of a TGF-beta family member or BMP, or allows binding of TGF-beta family members including one or more BMPs to their respective receptors by removing or preventing the TGF-beta member from binding to the TGF-binding-protein.

15 The present invention also provides peptide and polypeptide immunogens that, unexpectedly, may be used to generate and/or identify antibodies or fragments thereof that are capable of inhibiting, preventing, or impairing (*e.g.*, decreasing in a statistically significant manner) binding of the TGF-beta binding protein SOST to one or more BMPs. Exemplary peptide immunogens may comprise 6, 7, 8, 9, 10, 11, 12, 20-25, 21-50, 26-30, 31-40, 41-50, 51-  
20 60, 61-70, or 71-75 consecutive amino acids of a sclerostin polypeptide as provided herein (or of a variant thereof), such peptides comprising, for instance, amino acid sequences as set forth in SEQ ID NOS:2-19, 21-53, and 54. The present invention also provides peptide and polypeptide immunogens that may be used to generate and/or identify antibodies or fragments thereof that are capable of inhibiting, preventing, or impairing the formation of SOST homodimers. The  
25 antibodies of the present invention are useful for increasing the mineral content and mineral density of bone, thereby ameliorating numerous conditions that result in the loss of bone mineral content, including for example, disease, genetic predisposition, accidents that result in the lack of use of bone (*e.g.*, due to fracture), therapeutics that effect bone resorption or that kill bone forming cells, and normal aging.

30

### Sclerosteosis

Sclerosteosis is a disease related to abnormal bone mineral density in humans. Sclerosteosis is a term that was applied by Hansen (Hansen, H. G., *Sklerosteose in*: Opitz et al.,

eds. Handbuch der Kinderheilkunde, (Berlin: Springer 1967) 351-355) to a disorder similar to van Buchem hyperostosis corticalis generalisata but possibly differing in radiologic appearance of the bone changes and in the presence of asymmetric cutaneous syndactyly of the index and middle fingers in many cases. Sclerosteosis is now understood to be an autosomal semi-dominant disorder that is characterized by widely disseminated sclerotic lesions of the bone in the adult. The condition is progressive. Sclerosteosis also has a developmental aspect that is associated with syndactyly (two or more fingers fused together). Sclerosteosis Syndrome is associated with large stature, and many affected individuals attain a height of six feet or more. In addition, the jaw of persons with this condition has an unusually square appearance. The bone mineral content of homozygotes can be 1 to 6 fold greater than normal individuals, and bone mineral density can be 1 to 4 fold above normal values (*e.g.*, in comparison to unaffected siblings).

Sclerosteosis Syndrome occurs primarily in Afrikaaners of Dutch descent in South Africa. Approximately 1 of every 140 individuals in the Afrikaaner population is a carrier of the mutated gene (heterozygotes). The mutation shows 100% penetrance. Anecdotal reports indicate that increased bone mineral density has been observed in heterozygotes but they have no associated pathologies (*e.g.*, syndactyly or skull overgrowth).

No abnormality of the pituitary-hypothalamus axis has been observed in patients with sclerosteosis. In particular, no over-production of growth hormone and cortisone occurs, and sex hormone levels are normal in affected individuals. Bone turnover markers (such as osteoblast specific alkaline phosphatase, osteocalcin, type 1 procollagen C' propeptide (PICP), and total alkaline phosphatase (*see Comier, Curr. Opin. in Rheu.* 7:243 (1995)) indicate that hyperosteoblastic activity is associated with the disease but normal to slightly decreased osteoclast activity is observed as measured by markers of bone resorption (pyridinoline, deoxypyridinoline, N-telopeptide, urinary hydroxyproline, plasma tartrate-resistant acid phosphatases, and galactosyl hydroxylysine (*see Comier, supra*)).

Sclerosteosis is characterized by the continual deposition of bone throughout the skeleton during the lifetime of the affected individuals. In homozygotes the continual deposition of bone mineral leads to an overgrowth of bone in areas of the skeleton where mechanoreceptors are absent (*e.g.*, skull, jaw, cranium). In homozygotes with Sclerosteosis, the overgrowth of the bones of the skull leads to cranial compression and eventually to death due to excessive hydrostatic pressure on the brain stem. A generalized and diffuse sclerosis is observed in all other parts of the skeleton. Cortical areas of the long bones are greatly thickened resulting in a

substantial increase in bone strength. Trabecular connections are increased in thickness, which in turn increases the strength of the trabecular bone. Sclerotic bones appear unusually opaque to x-rays.

The rare genetic mutation that is responsible for Sclerosteosis Syndrome has been localized to the region of human chromosome 17. A gene within this region encodes a novel member of the TGF-beta binding-protein family (see, e.g., U.S. Patent Nos. 6,395,511, 6,489,445, and 6,495,736; Brunkow et al., *Am. J. Hum. Genet.* 68:577-89 (2001)). Therapeutics related to altering or modulating the level of sclerostin may therefore be useful in treating conditions and diseases related to abnormal bone development or deterioration. As described in more detail below, antibodies that specifically bind to this TGF-beta binding-protein, sclerostin (also referred to herein as Beer or as SOST), may be used to increase bone mineral content thus treating, preventing, retarding progression, or ameliorating the symptoms of a number of diseases.

#### TGF-Beta Superfamily

Among significant molecules to which reference is made herein include any known or novel members of the Transforming Growth Factor-beta (TGF-beta) superfamily, which includes bone morphogenic proteins (BMPs). Also included in the TGF-beta superfamily are TGF-beta receptors, which should be understood to refer to one or more receptors that are specific for a particular member of the TGF-beta superfamily (including BMPs), and TGF-beta binding proteins, which should be understood to refer to one or more polypeptides with specific binding affinity for a particular member or subset of members of the TGF-beta superfamily (including BMPs). A specific example of a TGF-beta binding protein includes sclerostin or SOST. Polynucleotide sequences encoding SOST and SOST variants in various animals, including humans, are provided herein in SEQ ID NOs: 55, 56, 57, 59, 61, 63, 65, and 67 and 69 (the encoded polypeptide sequences are provided in SEQ ID NOs: 1, 20, 58, 60, 62, 64, 66, 68, and 70, respectively). (See, e.g., U.S. Patent Nos. 6,395,511; 6,489,445; and 6,495,736. See also e.g., Balemans et al., 2002 *Dev. Biol.* 250:231; Schmitt et al., 1999 *J. Orthopaed. Res.* 17:269; Khalil, 1999 *Microbes Infect.* 1:1255; Miyazono et al., 1993 *Growth Factors* 8:11; von Bubnoff et al., 2001 *Dev. Biol.* 239:1; Koli et al., 2001 *Microsc. Res. Tech.* 52:354; Ebara et al., 2002 *Spine* 27(16 Suppl. 1):S10; Bondestam, 2002, *Ligands & Signaling Components of the Transforming Growth Factor  $\beta$  Family*, Helsinki University Biomedical Dissertations No. 17).

The TGF-beta superfamily contains a variety of growth factors that share

common sequence elements and structural motifs at both secondary and tertiary levels. This protein family exerts a wide spectrum of biological responses that affect a large variety of cell types. Many of the members of the TGF-beta family have important functions in pattern formation and tissue specification during embryonal development. In adults, TGF-beta family members are involved, for example, in wound healing, bone repair and bone remodeling, and in modulation of the immune system. In addition to the TGF-beta polypeptides, this superfamily includes BMPs, activins, inhibins, Growth and Differentiation Factors (GDFs), and Glial-Derived Neurotrophic Factors (GDNFs). Primary classification is established through general sequence features that bin a specific protein into a general subfamily. Additional stratification within the sub-family is possible due to stricter sequence conservation between members of the smaller group. In certain instances, such as with BMP-5, BMP-6, and BMP-7, the amino acid percent identity may be as high as 75% among members of the smaller group. This level of identity enables a single representative sequence to illustrate the key biochemical elements of the sub-group that separates it from other members of the larger family.

The crystal structure of TGF-beta2 has been determined. The general fold of the TGF-beta2 monomer contains a stable, compact, cysteine knotlike structure formed by three disulphide bridges. Dimerization, which is stabilized by one disulphide bridge, is antiparallel.

TGF-beta family members signal by inducing the formation of hetero-oligomeric receptor complexes. Transduction of TGF-beta signals involves two distinct subfamilies of transmembrane serine/threonine kinase receptors, type I and type II. At least seven type I receptors and five type II receptors have been identified (see Kawabata et al., *Cytokine Growth Factor Rev.* 9:49-61 (1998); Miyazono et al., *Adv. Immunol.* 75:115-57 (2000)). Each member of the TGF-beta family binds to a characteristic combination of type I and type II receptors, both of which are needed for signaling. In the current model for TGF-beta receptor activation, a TGF-beta ligand first binds to the type II receptor (TbR-II), which then recruits a type I receptor (TbR-I) to form a ligand/type II/type I ternary complex. The type I receptor, cannot bind ligand in the absence of TbR-II. TbR-II then phosphorylates TbR-I predominantly in a domain rich in glycine and serine residues (GS domain) in the juxtamembrane region, thereby activating TbR-I. The activated type I receptor kinase then phosphorylates particular members of the Smad family of proteins that translocate to the nucleus where they modulate transcription of specific genes.

#### Bone Morphogenic Proteins (BMPs): Key Regulatory Proteins of Bone Mineral Density

A major advance in the understanding of bone formation was the identification of



the BMPs, also known as osteogenic proteins (OPs), which regulate cartilage and bone differentiation in vivo. BMPs/OPs induce endochondral bone differentiation through a cascade of events by which mesenchymal stem cells differentiate into chondrocytes that lay down a cartilage structure that is reabsorbed and replaced by bone tissue (*see* Balemans et al., *Dev. Biol.* 250:231-50 (2002)). Thus, this process involves formation of cartilage, hypertrophy and calcification of the cartilage, vascular invasion, differentiation of osteoblasts, and formation of bone. As described above, the BMPs/OPs (BMP 2-14, and osteogenic protein 1 and -2, OP-1 and OP-2) (*see, e.g.*, GenBank P12643 (BMP-2); GenBank P12645 (BMP3); GenBank P55107 (BMP-3b, Growth/differentiation factor 10) (GDF-10)); GenBank P12644 (BMP4); GenBank P22003 (BMP5); GenBank P22004 (BMP6); GenBank P18075 (BMP7); GenBank P34820 (BMP8); GenBank Q9UK05 (BMP9); GenBank O95393 (BM10); GenBank O95390 (BMP11, Growth/differentiation factor 11 precursor (GDF-11)); GenBank O95972 (BM15)) are members of the TGF-beta superfamily. The striking evolutionary conservation among members of the BMP/OP sub-family suggests that they are important in the normal development and function of animals. In addition to postfetal chondrogenesis and osteogenesis, the BMPs/OPs play multiple roles in skeletogenesis, including the development of craniofacial and dental tissues. Different BMP family members also have biological activities in various other cell types, including monocytes, epithelial cells, mesenchymal cells, and neuronal cells. BMPs regulate cell proliferation and differentiation, chemotaxis and apoptosis, and also control fundamental roles, for example, left-right asymmetry, neurogenesis, mesoderm patterning, and embryonic development and organogenesis of a number of organs including the kidney, gut, lung, teeth, limb, amnion, and testis (*see* Balemans, *supra*).

BMPs are synthesized as large precursor proteins. Upon dimerization, the BMPs are proteolytically cleaved within the cell to yield carboxy-terminal mature proteins that are then secreted from the cell. BMPs, like other TGF-beta family members, initiate signal transduction by binding cooperatively to both type I and type II serine/threonine kinase receptors. Type I receptors for which BMPs may act as ligands include BMPR-IA (also known as ALK-3), BMPR-IB (also known as ALK-6), ALK-1, and ALK-2 (also known as ActR-I). Of the type II receptors, BMPs bind to BMP type II receptor (BMPR-II), Activin type II (ActR-II), and Activin type IIB (ActR-IIB). (*See* Balemans et al., *supra*, and references cited therein). Polynucleotide sequences and the encoded amino acid sequence of BMP type I receptor polypeptides are provided in the GenBank database, for example, GenBank NM\_004329 (SEQ ID NO:71 encoded by SEQ ID NO:85); D89675 (SEQ ID NO:72 encoded by SEQ ID NO:86);

NM\_001203 (SEQ ID NO:73 encoded by SEQ ID NO:87); S75359 (SEQ ID NO:74 encoded by SEQ ID NO:88); NM\_030849 (SEQ ID NO:75 encoded by SEQ ID NO:89); and D38082 (SEQ ID NO:76 encoded by SEQ ID NO:90). Other polypeptide sequences of type I receptors are provided in the GenBank database, for example, NP\_001194 (SEQ ID NO:77); BAA19765 (SEQ ID NO:78); and AAB33865 (SEQ ID NO:79). Polynucleotide sequences and the encoded amino acid sequence of BMP type II receptor polypeptides are provided in the GenBank database and include, for example, U25110 (SEQ ID NO:80 encoded by SEQ ID NO:91); NM\_033346 (SEQ ID NO:81 encoded by SEQ ID NO:92); NM\_001204 (SEQ ID NO:82 encoded by SEQ ID NO:93); and Z48923 (SEQ ID NO:83 encoded by SEQ ID NO:94). Additional polypeptide sequences of type II receptors are also provided in the GenBank database, for example, CAA88759 (SEQ ID NO:84).

BMPs, similar to other cystine-knot proteins, form a homodimer structure (Scheufler et al., *J. Mol. Biol.* 287:103-15 (1999)). According to evolutionary trace analysis performed on the BMP/TGF- $\beta$  family, the BMP type I receptor binding site and type II receptor binding site were mapped to the surface of the BMP structure (Innis et al., *Protein Eng.* 13:839-47 (2000)). The location of the type I receptor binding site on BMP was later confirmed by the x-ray structure of BMP-2/BMP Receptor IA complex (Nickel et al., *J. Joint Surg. Am.* 83A(Suppl 1(Pt 1)):S7-S14 (2001)). The predicted type II receptor binding site is in good agreement with the x-ray structure of TGF- $\beta$ 3/TGF- $\beta$  Type II receptor complex (Hart et al., *Nat. Struct. Biol.* 9:203-208 (2002)), which is highly similar to the BMP/BMP Receptor IIA system.

The BMP and Activin sub-families are subject to significant post-translational regulation by TGF-beta binding proteins. An intricate extracellular control system exists, whereby a high affinity antagonist is synthesized and exported, and subsequently forms a complex selectively with BMPs or activins to disrupt their biological activity (Smith, *Trends Genet.* 15:3-6 (1999)). A number of such TGF-beta binding proteins have been identified, and on the basis of sequence divergence, the antagonists appear to have evolved independently because of the lack of primary sequence conservation. In vertebrates, antagonists include noggin, chordin, chordin-like, follistatin, FSRP, the DAN/Cerberus protein family, and sclerostin (SOST) (see Balemans et al., *supra*, and references cited therein). The mechanism of antagonism seems to differ for the different antagonists (Iemura et al. (1998) *Proc. Natl. Acad. Sci. USA* 95:9337-9342).

The type I and type II receptor binding sites on the BMP antagonist noggin have also been mapped. Noggin binds to BMPs with high affinity (Zimmerman et al., 1996). A

study of the noggin/BMP-7 complex structure revealed the binding interactions between the two proteins (Groppe et al., *Nature* 420:636-42 (2002)). Superposition of the noggin-BMP-7 structure onto a model of the BMP signaling complex showed that noggin binding effectively masks both pairs of binding epitopes (*i.e.*, BMP Type I and Type II receptor binding sites) on BMP-7. The cysteine-rich scaffold sequence of noggin is preceded by an N-terminal segment of about 20 amino acid residues that are referred to as the "clip" (residues 28-48). The type I receptor-binding site is occluded by the N-terminal portion of the clip domain of Noggin, and the type II receptor binding site is occluded by the carboxy terminal portion of the clip domain. Two  $\beta$ -strands in the core region near the C-terminus of noggin also contact BMP-7 at the type II receptor binding site. This binding mode enables a Noggin dimer to efficiently block all the receptor binding sites (two type I and two type II receptor binding sites) on a BMP dimer.

#### Sclerostin Polypeptides and Encoding Polynucleotides

The BMP antagonist sclerostin (U.S. Patent Nos. 67,395,511, 6,489,445, and 6,495,736; see also SEQ ID NOs: 1, 20, 58, 60, 62, 64, 66, 68, and 70), possesses a nearly identical cysteine (disulfide) scaffold when compared with Human DAN, Human Gremlin, and Human Cerberus, and SCGF (U.S. Patent No. 5,780,263) but almost no homology at the nucleotide level (for background information, *see generally* Hsu et al., *Mol. Cell* 1:673-683 (1998)). Sclerostin should also be understood to include variants of this TGF-beta binding-protein (*e.g.*, SEQ ID NOs. 60 and 62). As used herein, a TGF-beta binding-protein variant polynucleotide refers to nucleic acid molecule that encodes a polypeptide having an amino acid sequence that is a modification (insertion, deletion, or substitution of one or more nucleotides) of SEQ ID NOs: 55-57, 59, 61, 63, 65, 67, or 69. Such variants include naturally-occurring polymorphisms or allelic variants of TGF-beta binding-protein encoding polynucleotides, as well as synthetic polynucleotides that encode conservative amino acid substitutions of these amino acid sequences. A variety of criteria known to those skilled in the art indicate whether amino acids at a particular position in a peptide or polypeptide are similar. For example, a similar amino acid or a conservative amino acid substitution is one in which an amino acid residue is replaced with an amino acid residue having a similar side chain, which include amino acids with basic side chains (*e.g.*, lysine, arginine, histidine); acidic side chains (*e.g.*, aspartic acid, glutamic acid); uncharged polar side chains (*e.g.*, glycine, asparagine, glutamine, serine, threonine, tyrosine, cysteine, histidine); nonpolar side chains (*e.g.*, alanine, valine, leucine, isoleucine, proline, phenylalanine, methionine, tryptophan); beta-branched side chains (*e.g.*,

threonine, valine, isoleucine), and aromatic side chains (*e.g.*, tyrosine, phenylalanine, tryptophan). Proline, which is considered more difficult to classify, shares properties with amino acids that have aliphatic side chains (*e.g.*, Leu, Val, Ile, and Ala). In certain circumstances, substitution of glutamine for glutamic acid or asparagine for aspartic acid may be considered a similar substitution in that glutamine and asparagine are amide derivatives of glutamic acid and aspartic acid, respectively.

Additional variant forms of a TGF-beta binding protein encoding polynucleotide are nucleic acid molecules that contain substitutions, insertions or deletions of one or more nucleotides found within the nucleotide sequences described herein. A variant or mutant of a TGF-beta binding protein may be constructed or identified so that the altered version of the TGF-beta binding protein competes with the wildtype TGF-beta binding protein. Such competition would preferably block the activity of the wildtype TGF-beta binding protein and thus lead to increased bone density.

An isolated polynucleotide is a nucleic acid molecule (polynucleotide or oligonucleotide) that is not integrated in the genomic DNA of an organism. A polynucleotide is considered to be isolated if, for example, it is cloned into a vector that is not a part of the natural environment. For example, a DNA molecule that encodes a TGF-binding protein that has been separated from the genomic DNA of a eukaryotic cell is an isolated DNA molecule. Another example of an isolated nucleic acid molecule is a chemically-synthesized nucleic acid molecule that is not integrated into the genome of an organism. The isolated nucleic acid molecule may be DNA, cDNA, RNA, or composed at least in part of nucleic acid analogs.

TGF-beta binding-protein variant polynucleotides can be identified by determining whether the polynucleotides hybridize with a nucleic acid molecule having the nucleotide sequence of SEQ ID NOs: 55-57, 59, 61, 63, 65, 67, or 69 under stringent conditions. In addition, TGF-beta binding-protein variant polynucleotides encode a protein having a cysteine backbone. TGF-beta binding-protein variant polynucleotides may also be identified by sequence comparison. As used herein, two amino acid sequences have "100% amino acid sequence identity" if the amino acid residues of the two amino acid sequences are the same when aligned for maximal correspondence. Similarly, two nucleotide sequences have "100% nucleotide sequence identity" if the nucleotides of the two nucleotide sequences are the same when aligned for maximal correspondence. Sequence comparisons can be performed using standard software programs such as those included in the LASERGENE bioinformatics computing suite, which is produced by DNASTAR (Madison, Wisconsin), or the BLAST

algorithm available at the NCBI web site ([Internet]<:http:www.ncbi.nlm.nih.gov>). Other methods for comparing two or more nucleotide or amino acid sequences by determining optimal alignment are well-known to those of skill in the art (*see*, for example, Peruski and Peruski, *The Internet and the New Biology: Tools for Genomic and Molecular Research* (ASM Press, Inc. 1997), Wu et al. (eds.), "Information Superhighway and Computer Databases of Nucleic Acids and Proteins," in *Methods in Gene Biotechnology*, pages 123-151 (CRC Press, Inc. 1997), and Bishop (ed.), *Guide to Human Genome Computing*, 2nd Edition (Academic Press, Inc. 1998)).

A variant TGF-beta binding protein should have at least a 50% amino acid sequence identity to SEQ ID NOs: 1, 20, 58, 60, 62, 64, 66, and 68, and preferably greater than 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, or 98% identity. Alternatively, TGF-beta binding-protein variants can be identified by having at least a 70% nucleotide sequence identity to SEQ ID NOs: 55-57, 59, 61, 63, 65, 67, or 69. Moreover, the present invention contemplates TGF-beta binding-protein polynucleotide variants having greater than 75%, 80%, 85%, 90%, or 95% identity to SEQ ID NO:55 or 57. Regardless of the particular method used to identify a TGF-beta binding-protein variant polynucleotide or variant TGF-beta binding protein, such a variant can be functionally characterized by, for example, its ability to bind to and/or inhibit the signaling of a selected member of the TGF-beta family of proteins, or by its ability to bind specifically to an anti-TGF-beta binding-protein antibody.

Within the context of this invention, a "functional fragment" of a TGF-beta binding-protein polynucleotide refers to a nucleic acid molecule that encodes a portion of a TGF-beta binding-protein polypeptide that either (1) possesses a functional activity as described herein or (2) specifically binds to an anti-TGF-beta binding protein antibody. For example, a functional fragment of a TGF-beta binding protein encoding polynucleotide described herein comprises a portion of the nucleotide sequence of SEQ ID Nos: 55-57, 59, 61, 63, 65, 67, or 69.

An "isolated polypeptide" referred to herein is a polypeptide that is removed from its natural environment. For example, a naturally-occurring protein is isolated if it is separated from some or all of the coexisting materials in the natural system, such as carbohydrate, lipid, nucleic acids, or proteinaceous impurities associated with the polypeptide in nature. Preferably, such isolated polypeptides are at least about 90% pure, more preferably at least about 95% pure, and most preferably at least about 99% pure.

#### Antibodies Specific for TGF-Beta Binding Proteins

The present invention provides antibodies that specifically bind to SOST and also

provides SOST polypeptide immunogens that may be used for generation and analysis of these antibodies and also provides methods for using such antibodies. The antibodies of the present invention as described herein specifically bind to a SOST polypeptide and thereby block or inhibit the binding of a BMP to the SOST polypeptide, that is, prevent or impair the interaction between the BMP and SOST. The effect of blocking this interaction is to alter (increase or decrease in a statistically significant manner) bone mineral density, preferably to increase bone mineral density.

Polypeptides or peptides useful for immunization and/or analysis of SOST-specific antibodies may also be selected by analyzing the primary, secondary, and tertiary structure of a TGF-beta binding protein according to methods known to those skilled in the art and described herein, in order to determine amino acid sequences more likely to generate an antigenic response in a host animal. See, e.g., Novotny, *Mol. Immunol.* 28:201-207 (1991); Berzofsky, *Science* 229:932-40 (1985)). Modeling and x-ray crystallography data may also be used to predict and/or identify which portions or regions of a TGF-beta binding protein interact with which portions of a TGF-beta binding protein ligand, such as a BMP. TGF-beta binding protein peptide immunogens may be designed and prepared that include amino acid sequences within or surrounding the portions or regions of interaction. These antibodies may be useful to block or impair binding of the TGF-beta binding protein to the same ligand and may also block or impair binding of the TGF-beta binding protein to one or more other ligands.

Antibodies or antigen binding fragments thereof contemplated by the present invention include antibodies that are capable of specifically binding to SOST and competitively inhibiting binding of a TGF-beta polypeptide, such as a BMP, to SOST. For example, the antibodies contemplated by the present invention competitively inhibit binding of the SOST polypeptide to the BMP Type I receptor site on a BMP, or to the BMP Type II receptor binding site, or may competitively inhibit binding of SOST to both the Type I and Type II receptor binding sites on a BMP. Without wishing to be bound by theory, when an anti-SOST antibody competitively inhibits binding of the Type I and/or Type II binding sites of the BMP polypeptide to SOST, thus blocking the antagonistic activity of SOST, the receptor binding sites on BMP are available to bind to the Type I and Type II receptors, thereby increasing bone mineralization. The binding interaction between a TGF-beta binding protein such as SOST and a TGF-beta polypeptide such as a BMP generally occurs when each of the ligand pairs forms a homodimer. Therefore instead of or in addition to using an antibody specific for SOST to block, impair, or prevent binding of SOST to a BMP by competitively inhibiting binding of SOST to BMP, a

SOST specific antibody may be used to block or impair SOST homodimer formation.

By way of example, one dimer of human Noggin, which is a BMP antagonist that has the ability to bind a BMP with high affinity (Zimmerman et al., *supra*), was isolated in complex with one dimer of human BMP-7 and analyzed by multiwavelength anomalous diffraction (MAD) (Groppe et al., *Nature* 420:636-42 (2002)). As discussed herein, this study revealed that Noggin dimer may efficiently block all the receptor binding sites (two type I and two type II receptor binding sites) on a BMP dimer. The location of the amino acids of Noggin that contact BMP-7 may be useful in modeling the interaction between other TGF-beta binding proteins, such as sclerostin (SOST), and BMPs, and thus aiding the design of peptides that may be used as immunogens to generate antibodies that block or impair such an interaction.

In one embodiment of the present invention, an antibody, or an antigen-binding fragment thereof, that binds specifically to a SOST polypeptide competitively inhibits binding of the SOST polypeptide to at least one or both of a bone morphogenic protein (BMP) Type I Receptor binding site and a BMP Type II Receptor binding site that are located on a BMP. The epitopes on SOST to which these antibodies bind may include or be included within contiguous amino acid sequences that are located at the N-terminus of the SOST polypeptide (amino acids at about positions 1-56 of SEQ ID NO:1). The polypeptides may also include a short linker peptide sequence that connects the N-terminal region to the core region, for example, polypeptides as provided in SEQ ID NO:47 (human) and SEQ ID NO:48 (rat). Shorter representative N-terminus peptide sequences of human SOST (e.g., SEQ ID NO:1) include SEQ ID NOS:2-6, and representative rat SOST (e.g., SEQ ID NO:20) peptide sequences include SEQ ID NOS:12-15.

Antibodies that specifically bind to a SOST polypeptide and block or competitively inhibit binding of the SOST polypeptide to a BMP, for example, by blocking or inhibiting binding to amino acids of a BMP corresponding to one or more of the Type I and Type II receptor binding sites may also specifically bind to peptides that comprise an amino acid sequence corresponding to the core region of SOST (amino acids at about positions 57-146 of SEQ ID NO:1). Polypeptides that include the core region may also include additional amino acids extending at either or both the N-terminus and C-terminus, for example, to include cysteine residues that may be useful for conjugating the polypeptide to a carrier molecule. Representative core polypeptides of human and rat SOST, for example, comprise the amino acid sequences set forth in SEQ ID NO:49 and SEQ ID NO:50, respectively. Such antibodies may also bind shorter polypeptide sequences. Representative human SOST core peptide sequences

are provided in SEQ ID NOs:21-24 and representative rat SOST core sequences are provided in SEQ ID NOs:25-28.

In another embodiment, antibodies that specifically bind to a SOST polypeptide impair (inhibit, prevent, or block, *e.g.*, decrease in a statistically significant manner) formation of a SOST homodimer. Because the interaction between SOST and a BMP may involve a homodimer of SOST and a homodimer of the BMP, an antibody that prevents or impairs homodimer formation of SOST may thereby alter bone mineral density, preferably increasing bone mineral density. In one embodiment, antibodies that bind to the core region of SOST prevent homodimer formation. Such antibodies may also bind to peptides that comprise contiguous amino acid sequences corresponding the core region, for example, SEQ ID NOs: 29, 30 and 53 (human SOST) and SEQ ID NOs:31 and 54 (rat SOST). Antibodies that bind to an epitope located on the C-terminal region of a SOST polypeptide (at about amino acid positions 147-190 of either SEQ ID NO:1 or 20) may also impair homodimer formation. Representative C-terminal polypeptides of human and rat SOST, for example, comprise the amino acid sequences set forth in SEQ ID NO:51 and SEQ ID NO:52, respectively. Such antibodies may also bind shorter polypeptide sequences. Representative human SOST C-terminal peptide sequences are provided in SEQ ID NOs:33-36 and representative rat SOST C-terminal sequences are provided in SEQ ID NOs:41-43.

The SOST polypeptides and peptides disclosed herein to which antibodies may specifically bind are useful as immunogens. These immunogens of the present invention may be used for immunizing an animal to generate a humoral immune response that results in production of antibodies that specifically bind to a Type I or Type II receptor binding site or both located on a BMP include peptides derived from the N-terminal region of SOST or that may prevent SOST homodimer formation.

Such SOST polypeptides and peptides that are useful as immunogens may also be used in methods for screening samples containing antibodies, for example, samples of purified antibodies, antisera, or cell culture supernatants or any other biological sample that may contain one or more antibodies specific for SOST. These peptides may also be used in methods for identifying and selecting from a biological sample one or more B cells that are producing an antibody that specifically binds to SOST (*e.g.*, plaque forming assays and the like). The B cells may then be used as source of a SOST specific antibody-encoding polynucleotide that can be cloned and/or modified by recombinant molecular biology techniques known in the art and described herein.



A "biological sample" as used herein refers in certain embodiments to a sample containing at least one antibody specific for a SOST polypeptide, and a biological sample may be provided by obtaining a blood sample (from which serum or plasma may be prepared), biopsy specimen, tissue explant, organ culture, or any other tissue or cell preparation from a subject or a biological source. A sample may further refer to a tissue or cell preparation in which the morphological integrity or physical state has been disrupted, for example, by dissection, dissociation, solubilization, fractionation, homogenization, biochemical or chemical extraction, pulverization, lyophilization, sonication, or any other means for processing a sample derived from a subject or biological source. The subject or biological source may be a human or non-human animal, a primary cell culture (*e.g.*, B cells immunized *in vitro*), or culture adapted cell line including but not limited to genetically engineered cell lines that may contain chromosomally integrated or episomal recombinant nucleic acid sequences, immortalized or immortalizable cell lines, somatic cell hybrid cell lines, differentiated or differentiable cell lines, transformed cell lines, and the like.

SOST peptide immunogens may also be prepared by synthesizing a series of peptides that, in total, represent the entire polypeptide sequence of a SOST polypeptide and that each have a portion of the SOST amino acid sequence in common with another peptide in the series. This overlapping portion would preferably be at least four amino acids, and more preferably 5, 6, 7, 8, 9, or 10 amino acids. Each peptide may be used to immunize an animal, the sera collected from the animal, and tested in an assay to identify which animal is producing antibodies that impair or block binding of SOST to a TGF-beta protein. Antibodies are then prepared from such identified immunized animals according to methods known in the art and described herein.

Peptides, polypeptides, and other non-peptide molecules that specifically bind to a TGF-beta binding protein such as SOST are contemplated by this invention. As used herein, a molecule is said to "specifically bind" to a TGF-beta binding protein if it reacts at a detectable level with the TGF-beta binding protein, but does not react detectably with peptides and polypeptides containing an unrelated sequence, or a sequence of a different TGF-beta binding protein. Preferred binding molecules include antibodies, which may be, for example, polyclonal, monoclonal, single chain, chimeric, anti-idiotypic, or CDR-grafted immunoglobulins, or fragments thereof, such as proteolytically generated or recombinantly produced immunoglobulin F(ab')<sub>2</sub>, Fab, Fab' Fv, and Fd fragments.

Particularly useful are anti-TGF-beta binding-protein antibodies that "specifically

bind" TGF-beta binding-protein of SEQ ID NOs: 1, 20, 58, 60, 62, 64, 66, or 68, but not to other TGF-beta binding-proteins such as Dan, Cerberus, SCGF, or Gremlin. Antibodies are understood to specifically bind a TGF-beta binding-protein, or a specific TGF-beta family member, if they bind with a  $K_a$  of greater than or equal to  $10^4 \text{ M}^{-1}$ , more preferably greater than or equal to about  $10^5 \text{ M}^{-1}$ , more preferably greater than or equal to about  $10^6 \text{ M}^{-1}$ , still more preferably greater than or equal to  $10^7 \text{ M}^{-1}$ , and still more preferably greater than or equal to  $10^8 \text{ M}^{-1}$ , and do not bind to other TGF-beta binding-proteins. Affinity of an antibody for its cognate antigen is also commonly expressed as a dissociation constant  $K_D$ , and an anti-SOST antibody specifically binds to a TGF-beta family member if it binds with a  $K_D$  of less than or equal to  $10^{-4} \text{ M}$ , more preferably less than or equal to about  $10^{-5} \text{ M}$ , more preferably less than or equal to about  $10^{-6} \text{ M}$ , still more preferably less than or equal to  $10^{-7} \text{ M}$ , and still more preferably less than or equal to  $10^{-8} \text{ M}$ . Furthermore, antibodies of the present invention preferably block, impair, or inhibit (*e.g.*, decrease with statistical significance) binding of a TGF-beta binding-protein to a TGF-beta family member.

The affinity of an antibody or binding partner, as well as the extent to which an antibody inhibits binding can be readily determined by one of ordinary skill in the art using conventional techniques, for example those described by Scatchard et al. (*Ann. N.Y. Acad. Sci.* 51:660-672, (1949)) or by surface plasmon resonance (SPR; BIAcore, Biosensor, Piscataway, NJ). For surface plasmon resonance, target molecules are immobilized on a solid phase and exposed to ligands in a mobile phase running along a flow cell. If ligand binding to the immobilized target occurs, the local refractive index changes, leading to a change in SPR angle, which can be monitored in real time by detecting changes in the intensity of the reflected light. The rates of change of the SPR signal can be analyzed to yield apparent rate constants for the association and dissociation phases of the binding reaction. The ratio of these values gives the apparent equilibrium constant (affinity) (*see, e.g.*, Wolff et al., *Cancer Res.* 53:2560-65 (1993)).

An antibody according to the present invention may belong to any immunoglobulin class, for example IgG, IgE, IgM, IgD, or IgA. It may be obtained from or derived from an animal, for example, fowl (*e.g.*, chicken) and mammals, which includes but is not limited to a mouse, rat, hamster, rabbit, or other rodent, a cow, horse, sheep, goat, camel, human, or other primate. The antibody may be an internalising antibody.

Methods well known in the art may be used to generate antibodies, polyclonal antisera, or monoclonal antibodies that are specific for a TGF-beta binding protein such as SOST. Antibodies also may be produced as genetically engineered immunoglobulins (Ig) or Ig

fragments designed to have desirable properties. For example, by way of illustration and not limitation, antibodies may include a recombinant IgG that is a chimeric fusion protein having at least one variable (V) region domain from a first mammalian species and at least one constant region domain from a second, distinct mammalian species. Most commonly, a chimeric antibody has murine variable region sequences and human constant region sequences. Such a murine/human chimeric immunoglobulin may be "humanized" by grafting the complementarity determining regions (CDRs) derived from a murine antibody, which confer binding specificity for an antigen, into human-derived V region framework regions and human-derived constant regions. Fragments of these molecules may be generated by proteolytic digestion, or optionally, by proteolytic digestion followed by mild reduction of disulfide bonds and alkylation. Alternatively, such fragments may also be generated by recombinant genetic engineering techniques.

Certain preferred antibodies are those antibodies that inhibit or block a TGF-beta binding protein activity within an *in vitro* assay, as described herein. Binding properties of an antibody to a TGF-beta binding protein may generally be assessed using immunodetection methods including, for example, an enzyme-linked immunosorbent assay (ELISA), immunoprecipitation, immunoblotting, countercurrent immunoelectrophoresis, radioimmunoassays, dot blot assays, inhibition or competition assays, and the like, which may be readily performed by those having ordinary skill in the art (*see, e.g.*, U.S. Patent Nos. 4,376,110 and 4,486,530; Harlow et al., *Antibodies: A Laboratory Manual*, Cold Spring Harbor Laboratory (1988)).

An immunogen may be comprised of cells expressing a TGF-beta binding protein such as a SOST polypeptide, purified or partially purified SOST polypeptide, or variants or fragments (*i.e.*, peptides) thereof, or peptides derived from a SOST polypeptide. Such peptides may be generated by proteolytic cleavage of a larger polypeptide, by recombinant molecular methodologies, or may be chemically synthesized. For instance, nucleic acid sequences encoding SOST polypeptide are provided herein, such that those skilled in the art may routinely prepare SOST polypeptide for use as immunogens. Peptides may be chemically synthesized by methods as described herein and known in the art. Alternatively, peptides may be generated by proteolytic cleavage of a SOST polypeptide, and individual peptides isolated by methods known in the art such as polyacrylamide gel electrophoresis or any number of liquid chromatography or other separation methods. Peptides useful as immunogens typically may have an amino acid sequence of at least 4 or 5 consecutive amino acids from a SOST polypeptide amino acid

sequence such as those described herein, and preferably have at least 6, 7, 8, 9, 10, 11, 12, 14, 15, 16, 18, 19 or 20 consecutive amino acids of a SOST polypeptide. Certain other preferred peptide immunogens comprise at least 6 but no more than 12 or more consecutive amino acids of a SOST polypeptide sequence, and other preferred peptide immunogens comprise at least 20 but no more than 75 consecutive amino acids, and other preferred peptide immunogens comprise at least 21 but no more than 50 consecutive amino acids of a SOST polypeptide. Other preferred peptide immunogens comprise 21-25, 26-30, 31-35, 36-40, 41-50, or any whole integer number of amino acids between and including 21 and 100 consecutive amino acids, and between 100 and 190 consecutive amino acids of a SOST polypeptide sequence.

Polyclonal antibodies that bind specifically to SOST can be prepared using methods described herein and well-known to persons skilled in the art (*see*, for example, Green et al., "Production of Polyclonal Antisera," in *Immunochemical Protocols* (Manson, ed.), pages 1-5 (Humana Press 1992); Harlow et al., *Antibodies: A Laboratory Manual*, Cold Spring Harbor Laboratory (1988); Williams et al., "Expression of foreign proteins in *E. coli* using plasmid vectors and purification of specific polyclonal antibodies," in *DNA Cloning 2: Expression Systems, 2nd Edition*, Glover et al. (eds.), page 15 (Oxford University Press 1995)). Antibodies to a TGF-beta binding-protein can be obtained, for example, by immunizing an animal with the TGF-beta binding-protein product of an expression vector, or by immunizing with an isolated TGF-beta binding protein or a peptide fragment thereof, as described herein. Although polyclonal antibodies are typically raised in animals such as rats, mice, rabbits, goats, cattle, or sheep, an anti-TGF-beta binding-protein antibody of the present invention may also be obtained from a subhuman primate. General techniques for raising diagnostically and therapeutically useful antibodies in baboons may be found, for example, in WO 91/11465 (1991) and in Losman et al., *Int. J. Cancer* 46:310, 1990.

Preparation of an immunogen for injection into animals may include covalent coupling of the TGF-beta binding protein (or variant or fragment thereof), to another immunogenic protein, for example, a carrier protein such as keyhole limpet hemocyanin (KLH) or bovine serum albumin (BSA) or the like. A polypeptide or peptide immunogen may include one or more additional amino acids at either the N-terminal or C-terminal end that facilitate the conjugation procedure (*e.g.*, the addition of a cysteine to facilitate conjugation of a peptide to KLH). Other amino acid residues within a polypeptide or peptide may be substituted to prevent conjugation at that particular amino acid position to a carrier polypeptide (*e.g.*, substituting a serine residue for cysteine at internal positions of a polypeptide/peptide) or may be substituted to

facilitate solubility or to increase immunogenicity.

TGF-beta binding protein peptide, polypeptide, or TGF-beta binding protein-expressing cells to be used as an immunogen may be emulsified in an adjuvant, for example, Freund's complete or incomplete adjuvant or the Ribi Adjuvant System (Corixa Corporation, Seattle, WA). *See also, e.g.,* Harlow et al., *supra*. The immunogen may be injected into the animal via any number of different routes, including intraperitoneally, intramuscularly, intraocularly, intradermally, or subcutaneously. In general, after the first injection, animals receive one or more booster immunizations according to a preferred schedule that may vary according to, *inter alia*, the antigen, the adjuvant (if any), and/or the particular animal species. The immune response may be monitored by periodically bleeding the animal and preparing and analyzing sera in an immunoassay, such as an ELISA or Ouchterlony diffusion assay, or the like, to determine the specific antibody titer. Once an adequate antibody titer is established, the animals may be bled periodically to accumulate the polyclonal antisera or may be exsanguinated. Polyclonal antibodies that bind specifically to the TGF-beta binding protein or peptide may then be purified from such antisera, for example, by affinity chromatography using protein A. Alternatively, affinity chromatography may be performed wherein the TGF-beta binding protein or peptide or an antibody specific for an Ig constant region of the particular immunized animal species is immobilized on a suitable solid support.

Antibodies for use in the invention include monoclonal antibodies that are prepared by conventional immunization and cell fusion procedures as described herein and known in the art. Monoclonal anti-TGF-beta binding-protein antibodies may be generated using a variety of techniques. Monoclonal antibodies that bind to specific antigens may be obtained by methods known to those skilled in the art (*see, for example, Kohler et al., Nature* 256:495, 1975; Coligan et al. (eds.), *Current Protocols in Immunology*, 1:2.5.1-2.6.7 (John Wiley & Sons 1991); U.S. Patent Nos. RE 32,011, 4,902,614, 4,543,439, and 4,411,993; *Monoclonal Antibodies, Hybridomas: A New Dimension in Biological Analyses*, Plenum Press, Kennett, McKearn, and Bechtol (eds.) (1980); and *Antibodies: A Laboratory Manual*, Harlow and Lane (eds.), Cold Spring Harbor Laboratory Press (1988); Picksley et al., "Production of monoclonal antibodies against proteins expressed in *E. coli*," in *DNA Cloning 2: Expression Systems, 2nd Edition*, Glover et al. (eds.), page 93 (Oxford University Press 1995)). Antibody fragments may be derived therefrom using any suitable standard technique such as proteolytic digestion, or optionally, by proteolytic digestion (for example, using papain or pepsin) followed by mild reduction of disulfide bonds and alkylation. Alternatively, such fragments may also be

generated by recombinant genetic engineering techniques.

Briefly, monoclonal antibodies can be obtained by injecting an animal, for example, a rat, hamster, or preferably a mouse, with an immunogen comprising a TGF-beta binding-protein gene product, or peptide fragment thereof, according to methods known in the art and described herein. The presence of specific antibody production may be monitored after the initial injection (injections may be administered by any one of several routes as described herein for generation of polyclonal antibodies) and/or after a booster injection by obtaining a serum sample and detecting the presence of an antibody that binds to a TGF-beta binding-protein or peptide using any one of several immunodetection methods known in the art and described herein. From animals producing antibodies that bind to the TGF-beta binding protein, lymphoid cells, most commonly cells from the spleen or lymph node, are removed to obtain B-lymphocytes. The B-lymphocytes are then fused with a drug-sensitized myeloma cell fusion partner, preferably one that is syngeneic with the immunized animal and that optionally has other desirable properties (*e.g.*, inability to express endogenous Ig gene products, *e.g.*, P3X63 - Ag 8.653 (ATCC No. CRL 1580); NS0, SP20) to produce hybridomas, which are immortal eukaryotic cell lines. The lymphoid (*e.g.*, spleen) cells and the myeloma cells may be combined for a few minutes with a membrane fusion-promoting agent, such as polyethylene glycol or a nonionic detergent, and then plated at low density on a selective medium that supports the growth of hybridoma cells but not unfused myeloma cells. A preferred selection media is HAT (hypoxanthine, aminopterin, thymidine). After a sufficient time, usually about one to two weeks, colonies of cells are observed. Single colonies are isolated, and antibodies produced by the cells may be tested for binding activity to the TGF-beta binding protein, or variant or fragment thereof, using any one of a variety of immunoassays known in the art and described herein. Hybridomas producing monoclonal antibodies with high affinity and specificity for SOST are preferred. Hybridomas that produce monoclonal antibodies that specifically bind to a TGF-beta binding protein or variant or fragment thereof are therefore contemplated by the present invention. The hybridomas are cloned (*e.g.*, by limited dilution cloning or by soft agar plaque isolation) and positive clones that produce an antibody specific to the antigen are selected and cultured. Antibodies that block, inhibit, or impair binding of the TGF-beta binding protein to a TGF-beta family member are preferred.

The monoclonal antibodies from the hybridoma cultures may be isolated from the supernatants of hybridoma cultures. An alternative method for production of a murine monoclonal antibody is to inject the hybridoma cells into the peritoneal cavity of a syngeneic

mouse, for example, a mouse that has been treated (*e.g.*, pristane-primed) to promote formation of ascites fluid containing the monoclonal antibody. Monoclonal antibodies can be isolated and purified by a variety of well-established techniques. Such isolation techniques include affinity chromatography with Protein-A Sepharose, size-exclusion chromatography, and ion-exchange chromatography (*see*, for example, Coligan at pages 2.7.1-2.7.12 and pages 2.9.1-2.9.3; Baines et al., "Purification of Immunoglobulin G (IgG)," in *Methods in Molecular Biology, Vol. 10*, pages 79-104 (The Humana Press, Inc. 1992)). Monoclonal antibodies may be purified by affinity chromatography using an appropriate ligand selected based on particular properties of the antibody (*e.g.*, heavy or light chain isotype, binding specificity, etc.). Examples of a suitable ligand, immobilized on a solid support, include Protein A, Protein G, an anti-constant region (light chain or heavy chain) antibody, an anti-idiotypic antibody, and a TGF-beta binding protein, or fragment or variant thereof.

In addition, an anti-TGF-beta binding-protein antibody of the present invention may be a human monoclonal antibody. Human monoclonal antibodies may be generated by any number of techniques with which those having ordinary skill in the art will be familiar. Such methods include, but are not limited to, Epstein Barr Virus (EBV) transformation of human peripheral blood cells (*e.g.*, containing B lymphocytes), *in vitro* immunization of human B cells, fusion of spleen cells from immunized transgenic mice carrying inserted human immunoglobulin genes, isolation from human immunoglobulin V region phage libraries, or other procedures as known in the art and based on the disclosure herein. For example, human monoclonal antibodies may be obtained from transgenic mice that have been engineered to produce specific human antibodies in response to antigenic challenge. Methods for obtaining human antibodies from transgenic mice are described, for example, by Green et al., *Nature Genet.* 7:13, 1994; Lonberg et al., *Nature* 368:856, 1994; Taylor et al., *Int. Immun.* 6:579, 1994; U.S. Patent No. 5,877,397; Bruggemann et al., 1997 *Curr. Opin. Biotechnol.* 8:455-58; Jakobovits et al., 1995 *Ann. N. Y. Acad. Sci.* 764:525-35. In this technique, elements of the human heavy and light chain locus are introduced into strains of mice derived from embryonic stem cell lines that contain targeted disruptions of the endogenous heavy chain and light chain loci. (*See also* Bruggemann et al., *Curr. Opin. Biotechnol.* 8:455-58 (1997)). For example, human immunoglobulin transgenes may be mini-gene constructs, or transloci on yeast artificial chromosomes, which undergo B cell-specific DNA rearrangement and hypermutation in the mouse lymphoid tissue. Human monoclonal antibodies may be obtained by immunizing the transgenic mice, which may then produce human antibodies specific for the antigen. Lymphoid cells of the immunized transgenic

mice can be used to produce human antibody-secreting hybridomas according to the methods described herein. Polyclonal sera containing human antibodies may also be obtained from the blood of the immunized animals.

Another method for generating human TGF-beta binding protein specific  
5 monoclonal antibodies includes immortalizing human peripheral blood cells by EBV transformation. *See, e.g.*, U.S. Patent No. 4,464,456. Such an immortalized B cell line (or lymphoblastoid cell line) producing a monoclonal antibody that specifically binds to a TGF-beta binding protein (or a variant or fragment thereof) can be identified by immunodetection methods as provided herein, for example, an ELISA, and then isolated by standard cloning techniques.  
10 The stability of the lymphoblastoid cell line producing an anti-TGF-beta binding protein antibody may be improved by fusing the transformed cell line with a murine myeloma to produce a mouse-human hybrid cell line according to methods known in the art (*see, e.g.*, Glasky et al., *Hybridoma* 8:377-89 (1989)). Still another method to generate human monoclonal antibodies is *in vitro* immunization, which includes priming human splenic B cells with antigen,  
15 followed by fusion of primed B cells with a heterohybrid fusion partner. *See, e.g.*, Boerner et al., 1991 *J. Immunol.* 147:86-95.

In certain embodiments, a B cell that is producing an anti-SOST antibody is selected and the light chain and heavy chain variable regions are cloned from the B cell according to molecular biology techniques known in the art (WO 92/02551; US patent  
20 5,627,052; Babcook et al., *Proc. Natl. Acad. Sci. USA* 93:7843-48 (1996)) and described herein. Preferably B cells from an immunized animal are isolated from the spleen, lymph node, or peripheral blood sample by selecting a cell that is producing an antibody that specifically binds to SOST. B cells may also be isolated from humans, for example, from a peripheral blood sample. Methods for detecting single B cells that are producing an antibody with the desired  
25 specificity are well known in the art, for example, by plaque formation, fluorescence-activated cell sorting, *in vitro* stimulation followed by detection of specific antibody, and the like. Methods for selection of specific antibody producing B cells include, for example, preparing a single cell suspension of B cells in soft agar that contains SOST or a peptide fragment thereof. Binding of the specific antibody produced by the B cell to the antigen results in the formation of  
30 a complex, which may be visible as an immunoprecipitate. After the B cells producing the specific antibody are selected, the specific antibody genes may be cloned by isolating and amplifying DNA or mRNA according to methods known in the art and described herein.

For particular uses, fragments of anti-TGF-beta binding protein antibodies may



be desired. Antibody fragments,  $F(ab')_2$ , Fab, Fab', Fv, Fc, Fd, retain the antigen binding site of the whole antibody and therefore bind to the same epitope. These antigen-binding fragments derived from an antibody can be obtained, for example, by proteolytic hydrolysis of the antibody, for example, pepsin or papain digestion of whole antibodies according to conventional methods. As an illustration, antibody fragments can be produced by enzymatic cleavage of antibodies with pepsin to provide a 5S fragment denoted  $F(ab')_2$ . This fragment can be further cleaved using a thiol reducing agent to produce 3.5S Fab' monovalent fragments. Optionally, the cleavage reaction can be performed using a blocking group for the sulfhydryl groups that result from cleavage of disulfide linkages. As an alternative, an enzymatic cleavage using papain produces two monovalent Fab fragments and an Fc fragment directly. These methods are described, for example, by Goldenberg, U.S. patent No. 4,331,647, Nisonoff et al., *Arch. Biochem. Biophys.* 89:230, 1960; Porter, *Biochem. J.* 73:119, 1959; Edelman et al., in *Methods in Enzymology* 1:422 (Academic Press 1967); and by Coligan at pages 2.8.1-2.8.10 and 2.10.-2.10.4. Other methods for cleaving antibodies, such as separating heavy chains to form monovalent light-heavy chain fragments (Fd), further cleaving of fragments, or other enzymatic, chemical, or genetic techniques may also be used, so long as the fragments bind to the antigen that is recognized by the intact antibody.

An antibody fragment may also be any synthetic or genetically engineered protein that acts like an antibody in that it binds to a specific antigen to form a complex. For example, antibody fragments include isolated fragments consisting of the light chain variable region, "Fv" fragments consisting of the variable regions of the heavy and light chains, recombinant single chain polypeptide molecules in which light and heavy variable regions are connected by a peptide linker (scFv proteins), and minimal recognition units consisting of the amino acid residues that mimic the hypervariable region. The antibody of the present invention preferably comprises at least one variable region domain. The variable region domain may be of any size or amino acid composition and will generally comprise at least one hypervariable amino acid sequence responsible for antigen binding and which is adjacent to or in frame with one or more framework sequences. In general terms, the variable (V) region domain may be any suitable arrangement of immunoglobulin heavy ( $V_H$ ) and/or light ( $V_L$ ) chain variable domains. Thus, for example, the V region domain may be monomeric and be a  $V_H$  or  $V_L$  domain, which is capable of independently binding antigen with acceptable affinity. Alternatively, the V region domain may be dimeric and contain  $V_H$ - $V_H$ ,  $V_H$ - $V_L$ , or  $V_L$ - $V_L$  dimers. Preferably, the V region dimer comprises at least one  $V_H$  and at least one  $V_L$  chain that are non-covalently associated (hereinafter referred to

as F<sub>v</sub>). If desired, the chains may be covalently coupled either directly, for example via a disulfide bond between the two variable domains, or through a linker, for example a peptide linker, to form a single chain F<sub>v</sub> (scF<sub>v</sub>).

The variable region domain may be any naturally occurring variable domain or an engineered version thereof. By engineered version is meant a variable region domain that has been created using recombinant DNA engineering techniques. Such engineered versions include those created, for example, from a specific antibody variable region by insertions, deletions, or changes in or to the amino acid sequences of the specific antibody. Particular examples include engineered variable region domains containing at least one CDR and optionally one or more framework amino acids from a first antibody and the remainder of the variable region domain from a second antibody.

The variable region domain may be covalently attached at a C-terminal amino acid to at least one other antibody domain or a fragment thereof. Thus, for example, a V<sub>H</sub> domain that is present in the variable region domain may be linked to an immunoglobulin C<sub>H</sub>1 domain, or a fragment thereof. Similarly a V<sub>L</sub> domain may be linked to a C<sub>K</sub> domain or a fragment thereof. In this way, for example, the antibody may be a Fab fragment wherein the antigen binding domain contains a associated V<sub>H</sub> and V<sub>L</sub> domains covalently linked at their C-termini to a CH1 and C<sub>K</sub> domain, respectively. The CH1 domain may be extended with further amino acids, for example to provide a hinge region or a portion of a hinge region domain as found in a Fab' fragment, or to provide further domains, such as antibody CH2 and CH3 domains.

Another form of an antibody fragment is a peptide comprising for a single complementarity-determining region (CDR). CDR peptides ("minimal recognition units") can be obtained by constructing polynucleotides that encode the CDR of an antibody of interest. Such polynucleotides are prepared, for example, by using the polymerase chain reaction to synthesize the variable region using mRNA of antibody-producing cells as a template (see, for example, Larrick et al., *Methods: A Companion to Methods in Enzymology* 2:106, 1991; Courtenay-Luck, "Genetic Manipulation of Monoclonal Antibodies," in *Monoclonal Antibodies: Production, Engineering and Clinical Application*, Ritter et al. (eds.), page 166 (Cambridge University Press 1995); and Ward et al., "Genetic Manipulation and Expression of Antibodies," in *Monoclonal Antibodies: Principles and Applications*, Birch et al., (eds.), page 137 (Wiley-Liss, Inc. 1995)).

Alternatively, the antibody may be a recombinant or engineered antibody obtained by the use of recombinant DNA techniques involving the manipulation and re-expression of DNA encoding antibody variable and/or constant regions. Such DNA is known and/or is readily

available from DNA libraries including for example phage-antibody libraries (see Chiswell and McCafferty, *Tibtech.* 10:80-84 (1992)) or if desired can be synthesized. Standard molecular biology and/or chemistry procedures may be used to sequence and manipulate the DNA, for example, to introduce codons to create cysteine residues, or to modify, add or delete other amino acids or domains as desired.

Chimeric antibodies, specific for a TGF-beta binding protein, and which include humanized antibodies, may also be generated according to the present invention. A chimeric antibody has at least one constant region domain derived from a first mammalian species and at least one variable region domain derived from a second, distinct mammalian species (see, e.g., Morrison et al., *Proc. Natl. Acad. Sci. USA*, 81:6851-55 (1984)). In preferred embodiments, a chimeric antibody may be constructed by cloning the polynucleotide sequence that encodes at least one variable region domain derived from a non-human monoclonal antibody, such as the variable region derived from a murine, rat, or hamster monoclonal antibody, into a vector containing a nucleotide sequence that encodes at least one human constant region (see, e.g., Shin et al., *Methods Enzymol.* 178:459-76 (1989); Walls et al., *Nucleic Acids Res.* 21:2921-29 (1993)). By way of example, the polynucleotide sequence encoding the light chain variable region of a murine monoclonal antibody may be inserted into a vector containing a nucleotide sequence encoding the human kappa light chain constant region sequence. In a separate vector, the polynucleotide sequence encoding the heavy chain variable region of the monoclonal antibody may be cloned in frame with sequences encoding a human IgG constant region, for example, the human IgG1 constant region. The particular human constant region selected may depend upon the effector functions desired for the particular antibody (e.g., complement fixing, binding to a particular Fc receptor, etc.). Preferably, the constructed vectors will be transfected into eukaryotic cells for stable expression of the chimeric antibody. Another method known in the art for generating chimeric antibodies is homologous recombination (e.g., U.S. Patent No. 5,482,856).

A non-human/human chimeric antibody may be further genetically engineered to create a "humanized" antibody. Such a humanized antibody may comprise a plurality of CDRs derived from an immunoglobulin of a non-human mammalian species, at least one human variable framework region, and at least one human immunoglobulin constant region. Useful strategies for designing humanized antibodies may include, for example by way of illustration and not limitation, identification of human variable framework regions that are most homologous to the non-human framework regions of the chimeric antibody. Without wishing to

be bound by theory, such a strategy may increase the likelihood that the humanized antibody will retain specific binding affinity for a TGF-beta binding protein, which in some preferred embodiments may be substantially the same affinity for a TGF-beta binding protein or variant or fragment thereof, and in certain other preferred embodiments may be a greater affinity for TGF-beta binding protein. *See, e.g., Jones et al., 1986 Nature 321:522-25; Riechmann et al., 1988 Nature 332:323-27.* Designing such a humanized antibody may therefore include determining CDR loop conformations and structural determinants of the non-human variable regions, for example, by computer modeling, and then comparing the CDR loops and determinants to known human CDR loop structures and determinants. *See, e.g., Padlan et al., 1995 FASEB 9:133-39; Chothia et al., 1989 Nature, 342:377-383.* Computer modeling may also be used to compare human structural templates selected by sequence homology with the non-human variable regions. *See, e.g., Bajorath et al., 1995 Ther. Immunol. 2:95-103; EP-0578515-A3.* If humanization of the non-human CDRs results in a decrease in binding affinity, computer modeling may aid in identifying specific amino acid residues that could be changed by site-directed or other mutagenesis techniques to partially, completely or supra-optimally (*i.e.*, increase to a level greater than that of the non-humanized antibody) restore affinity. Those having ordinary skill in the art are familiar with these techniques, and will readily appreciate numerous variations and modifications to such design strategies.

One such method for preparing a humanized antibody is called veneering. As used herein, the terms "veneered FRs" and "recombinantly veneered FRs" refer to the selective replacement of FR residues from, *e.g.*, a rodent heavy or light chain V region, with human FR residues in order to provide a xenogeneic molecule comprising an antigen-binding site that retains substantially all of the native FR polypeptide folding structure. Veneering techniques are based on the understanding that the ligand binding characteristics of an antigen-binding site are determined primarily by the structure and relative disposition of the heavy and light chain CDR sets within the antigen-binding surface. *Davies et al., Ann. Rev. Biochem. 59:439-73, 1990.* Thus, antigen binding specificity can be preserved in a humanized antibody only wherein the CDR structures, their interaction with each other, and their interaction with the rest of the V region domains are carefully maintained. By using veneering techniques, exterior (*e.g.*, solvent-accessible) FR residues that are readily encountered by the immune system are selectively replaced with human residues to provide a hybrid molecule that comprises either a weakly immunogenic, or substantially non-immunogenic veneered surface.

The process of veneering makes use of the available sequence data for human

antibody variable domains compiled by Kabat et al., in *Sequences of Proteins of Immunological Interest*, 4th ed., (U.S. Dept. of Health and Human Services, U.S. Government Printing Office, 1987), updates to the Kabat database, and other accessible U.S. and foreign databases (both nucleic acid and protein). Solvent accessibilities of V region amino acids can be deduced from the known three-dimensional structure for human and murine antibody fragments. Initially, the FRs of the variable domains of an antibody molecule of interest are compared with corresponding FR sequences of human variable domains obtained from the above-identified sources. The most homologous human V regions are then compared residue by residue to corresponding murine amino acids. The residues in the murine FR that differ from the human counterpart are replaced by the residues present in the human moiety using recombinant techniques well known in the art. Residue switching is only carried out with moieties which are at least partially exposed (solvent accessible), and care is exercised in the replacement of amino acid residues that may have a significant effect on the tertiary structure of V region domains, such as proline, glycine, and charged amino acids.

In this manner, the resultant "veneered" antigen-binding sites are thus designed to retain the rodent CDR residues, the residues substantially adjacent to the CDRs, the residues identified as buried or mostly buried (solvent inaccessible), the residues believed to participate in non-covalent (*e.g.*, electrostatic and hydrophobic) contacts between heavy and light chain domains, and the residues from conserved structural regions of the FRs which are believed to influence the "canonical" tertiary structures of the CDR loops. These design criteria are then used to prepare recombinant nucleotide sequences that combine the CDRs of both the heavy and light chain of a antigen-binding site into human-appearing FRs that can be used to transfect mammalian cells for the expression of recombinant human antibodies that exhibit the antigen specificity of the rodent antibody molecule.

An additional method for selecting antibodies that specifically bind to a TGF-beta binding protein or variant or fragment thereof is by phage display. *See, e.g.*, Winter et al., 1994 *Annu. Rev. Immunol.* 12:433-55; Burton et al., 1994 *Adv. Immunol.* 57:191-280. Human or murine immunoglobulin variable region gene combinatorial libraries may be created in phage vectors that can be screened to select Ig fragments (Fab, Fv, sFv, or multimers thereof) that bind specifically to TGF-beta binding protein or variant or fragment thereof. *See, e.g.*, U.S. Patent No. 5,223,409; Huse et al., 1989 *Science* 246:1275-81; Sastry et al., *Proc. Natl. Acad. Sci. USA* 86:5728-32 (1989); Alting-Mees et al., *Strategies in Molecular Biology* 3:1-9 (1990); Kang et al., 1991 *Proc. Natl. Acad. Sci. USA* 88:4363-66; Hoogenboom et al., 1992 *J. Molec. Biol.*

227:381-388; Schlebusch et al., 1997 *Hybridoma* 16:47-52 and references cited therein. For example, a library containing a plurality of polynucleotide sequences encoding Ig variable region fragments may be inserted into the genome of a filamentous bacteriophage, such as M13 or a variant thereof, in frame with the sequence encoding a phage coat protein. A fusion protein  
5 may be a fusion of the coat protein with the light chain variable region domain and/or with the heavy chain variable region domain. According to certain embodiments, immunoglobulin Fab fragments may also be displayed on a phage particle (*see, e.g.*, U.S. Patent No. 5,698,426).

Heavy and light chain immunoglobulin cDNA expression libraries may also be prepared in lambda phage, for example, using  $\lambda$ ImmunoZap<sup>TM</sup>(H) and  $\lambda$ ImmunoZap<sup>TM</sup>(L)  
10 vectors (Stratagene, La Jolla, California). Briefly, mRNA is isolated from a B cell population, and used to create heavy and light chain immunoglobulin cDNA expression libraries in the  $\lambda$  ImmunoZap(H) and  $\lambda$ ImmunoZap(L) vectors. These vectors may be screened individually or co-expressed to form Fab fragments or antibodies (*see* Huse et al., *supra*; *see also* Sastry et al., *supra*). Positive plaques may subsequently be converted to a non-lytic plasmid that allows high  
15 level expression of monoclonal antibody fragments from *E. coli*.

Similarly, portions or fragments, such as Fab and Fv fragments, of antibodies may also be constructed using conventional enzymatic digestion or recombinant DNA techniques to incorporate the variable regions of a gene that encodes a antibody specific for a TGF-beta binding protein. Within one embodiment, in a hybridoma the variable regions of a  
20 gene expressing a monoclonal antibody of interest are amplified using nucleotide primers. These primers may be synthesized by one of ordinary skill in the art, or may be purchased from commercially available sources. (*See, e.g.*, Stratagene (La Jolla, California), which sells primers for mouse and human variable regions including, among others, primers for V<sub>Ha</sub>, V<sub>Hb</sub>, V<sub>Hc</sub>, V<sub>Hd</sub>, C<sub>H1</sub>, V<sub>L</sub> and C<sub>L</sub> regions.) These primers may be used to amplify heavy or light chain  
25 variable regions, which may then be inserted into vectors such as ImmunoZAP<sup>TM</sup> H or ImmunoZAP<sup>TM</sup> L (Stratagene), respectively. These vectors may then be introduced into *E. coli*, yeast, or mammalian-based systems for expression. Large amounts of a single-chain protein containing a fusion of the V<sub>H</sub> and V<sub>L</sub> domains may be produced using these methods (*see* Bird et al., *Science* 242:423-426, 1988). In addition, such techniques may be used to humanize a  
30 non-human antibody V region without altering the binding specificity of the antibody.

In certain particular embodiments of the invention, combinatorial phage libraries may also be used for humanization of non-human variable regions. *See, e.g.*, Rosok et al., 1996 *J. Biol. Chem.* 271:22611-18; Rader et al., 1998 *Proc. Natl. Acad. Sci. USA* 95:8910-15. A

phage library may be screened to select an Ig variable region fragment of interest by immunodetection methods known in the art and described herein, and the DNA sequence of the inserted immunoglobulin gene in the phage so selected may be determined by standard techniques. *See*, Sambrook et al., 2001 *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor Press. The selected Ig-encoding sequence may then be cloned into another suitable vector for expression of the Ig fragment or, optionally, may be cloned into a vector containing Ig constant regions, for expression of whole immunoglobulin chains.

In certain other embodiments, the invention contemplates SOST-specific antibodies that are multimeric antibody fragments. Useful methodologies are described generally, for example in Hayden et al. 1997, *Curr Opin. Immunol.* 9:201-12; Coloma et al., 1997 *Nat. Biotechnol.* 15:159-63. For example, multimeric antibody fragments may be created by phage techniques to form miniantibodies (U.S. Patent No. 5,910 573) or diabodies (Holliger et al., 1997, *Cancer Immunol. Immunother.* 45:128-130).

In certain embodiments of the invention, an antibody specific for SOST may be an antibody that is expressed as an intracellular protein. Such intracellular antibodies are also referred to as intrabodies and may comprise an Fab fragment, or preferably comprise a scFv fragment (*see, e.g.*, Lecerf et al., *Proc. Natl. Acad. Sci. USA* 98:4764-49 (2001)). The framework regions flanking the CDR regions can be modified to improve expression levels and solubility of an intrabody in an intracellular reducing environment (*see, e.g.*, Worn et al., *J. Biol. Chem.* 275:2795-803 (2000)). An intrabody may be directed to a particular cellular location or organelle, for example by constructing a vector that comprises a polynucleotide sequence encoding the variable regions of an intrabody that may be operatively fused to a polynucleotide sequence that encodes a particular target antigen within the cell (*see, e.g.*, Graus-Porta et al., *Mol. Cell Biol.* 15:1182-91 (1995); Lener et al., *Eur. J. Biochem.* 267:1196-205 (2000)). An intrabody may be introduced into a cell by a variety of techniques available to the skilled artisan including via a gene therapy vector, or a lipid mixture (*e.g.*, Provectin™ manufactured by Imgenex Corporation, San Diego, CA), or according to photochemical internalization methods.

Introducing amino acid mutations into an immunoglobulin molecule specific for a TGF-beta binding protein may be useful to increase the specificity or affinity for TGF-beta binding protein or to alter an effector function. Immunoglobulins with higher affinity for TGF-beta binding protein may be generated by site-directed mutagenesis of particular residues. Computer assisted three-dimensional molecular modeling may be employed to identify the amino acid residues to be changed, in order to improve affinity for the TGF-beta binding protein.

See, e.g., Mountain et al., 1992, *Biotechnol. Genet. Eng. Rev.* 10: 1-142. Alternatively, combinatorial libraries of CDRs may be generated in M13 phage and screened for immunoglobulin fragments with improved affinity. See, e.g., Glaser et al., 1992, *J. Immunol.* 149:3903-3913; Barbas et al., 1994 *Proc. Natl. Acad. Sci. USA* 91:3809-13; U.S. Patent No. 5,792,456.

Effector functions may also be altered by site-directed mutagenesis. See, e.g., Duncan et al., 1988 *Nature* 332:563-64; Morgan et al., 1995 *Immunology* 86:319-24; Eghtedarzede-Kondri et al., 1997 *Biotechniques* 23:830-34. For example, mutation of the glycosylation site on the Fc portion of the immunoglobulin may alter the ability of the immunoglobulin to fix complement. See, e.g., Wright et al., 1997 *Trends Biotechnol.* 15:26-32. Other mutations in the constant region domains may alter the ability of the immunoglobulin to fix complement, or to effect antibody-dependent cellular cytotoxicity. See, e.g., Duncan et al., 1988 *Nature* 332:563-64; Morgan et al., 1995 *Immunology* 86:319-24; Sensel et al., 1997 *Mol. Immunol.* 34:1019-29.

According to certain embodiments, non-human, human, or humanized heavy chain and light chain variable regions of any of the Ig molecules described herein may be constructed as single chain Fv (scFv) polypeptide fragments (single chain antibodies). See, e.g., Bird et al., 1988 *Science* 242:423-426; Huston et al., 1988 *Proc. Natl. Acad. Sci. USA* 85:5879-5883. Multi-functional scFv fusion proteins may be generated by linking a polynucleotide sequence encoding an scFv polypeptide in-frame with at least one polynucleotide sequence encoding any of a variety of known effector proteins. These methods are known in the art, and are disclosed, for example, in EP-B1-0318554, U.S. Patent No. 5,132,405, U.S. Patent No. 5,091,513, and U.S. Patent No. 5,476,786. By way of example, effector proteins may include immunoglobulin constant region sequences. See, e.g., Hollenbaugh et al., 1995 *J. Immunol. Methods* 188:1-7. Other examples of effector proteins are enzymes. As a non-limiting example, such an enzyme may provide a biological activity for therapeutic purposes (see, e.g., Siemers et al., 1997 *Bioconjug. Chem.* 8:510-19), or may provide a detectable activity, such as horseradish peroxidase-catalyzed conversion of any of a number of well-known substrates into a detectable product, for diagnostic uses. Still other examples of scFv fusion proteins include Ig-toxin fusions, or immunotoxins, wherein the scFv polypeptide is linked to a toxin.

The scFv or any antibody fragment described herein may, in certain embodiments, be fused to peptide or polypeptide domains that permits detection of specific



binding between the fusion protein and antigen (*e.g.*, a TGF-beta binding protein). For example, the fusion polypeptide domain may be an affinity tag polypeptide for detecting binding of the scFv fusion protein to a TGF-beta binding protein by any of a variety of techniques with which those skilled in the art will be familiar. Examples of a peptide tag, include avidin, streptavidin or His (*e.g.*, polyhistidine). Detection techniques may also include, for example, binding of an avidin or streptavidin fusion protein to biotin or to a biotin mimetic sequence (*see, e.g.*, Luo et al., 1998 *J. Biotechnol.* 65:225 and references cited therein), direct covalent modification of a fusion protein with a detectable moiety (*e.g.*, a labeling moiety), non-covalent binding of the fusion protein to a specific labeled reporter molecule, enzymatic modification of a detectable substrate by a fusion protein that includes a portion having enzyme activity, or immobilization (covalent or non-covalent) of the fusion protein on a solid-phase support. Other useful affinity polypeptides for construction of scFv fusion proteins may include streptavidin fusion proteins, as disclosed, for example, in WO 89/03422, U.S. 5,489,528, U.S. 5,672,691, WO 93/24631, U.S. 5,168,049, U.S. 5,272,254; avidin fusion proteins (*see, e.g.*, EP 511,747); an enzyme such as glutathione-S-transferase; and *Staphylococcus aureus* protein A polypeptide.

The polynucleotides encoding an antibody or fragment thereof that specifically bind a TGF-beta binding protein, as described herein, may be propagated and expressed according to any of a variety of well-known procedures for nucleic acid excision, ligation, transformation, and transfection using any number of known expression vectors. Thus, in certain embodiments expression of an antibody fragment may be preferred in a prokaryotic host, such as *Escherichia coli* (*see, e.g.*, Pluckthun et al., 1989 *Methods Enzymol.* 178:497-515). In certain other embodiments, expression of the antibody or a fragment thereof may be preferred in a eukaryotic host cell, including yeast (*e.g.*, *Saccharomyces cerevisiae*, *Schizosaccharomyces pombe*, and *Pichia pastoris*), animal cells (including mammalian cells) or plant cells. Examples of suitable animal cells include, but are not limited to, myeloma (such as a mouse NSO line), COS, CHO, or hybridoma cells. Examples of plant cells include tobacco, corn, soybean, and rice cells.

One or more replicable expression vectors containing DNA encoding a variable and/or constant region may be prepared and used to transform an appropriate cell line, for example, a non-producing myeloma cell line, such as a mouse NSO line or a bacteria, such as *E.coli*, in which production of the antibody will occur. In order to obtain efficient transcription and translation, the DNA sequence in each vector should include appropriate regulatory sequences, particularly a promoter and leader sequence operatively linked to the variable domain sequence.

Particular methods for producing antibodies in this way are generally well known and routinely used. For example, basic molecular biology procedures are described by Maniatis et al. (*Molecular Cloning, A Laboratory Manual*, 2nd ed., Cold Spring Harbor Laboratory, New York, 1989; *see also* Maniatis et al, 3rd ed., Cold Spring Harbor Laboratory, New York, (2001)). DNA sequencing  
5 can be performed as described in Sanger *et al* (PNAS 74:5463, (1977)) and the Amersham International plc sequencing handbook, and site directed mutagenesis can be carried out according to methods known in the art (Kramer et al., *Nucleic Acids Res.* 12:9441, (1984); Kunkel *Proc. Natl. Acad. Sci. USA* 82:488-92 (1985); Kunkel et al., *Methods in Enzymol.* 154:367-82 (1987); the Anglian Biotechnology Ltd handbook). Additionally, numerous publications describe techniques  
10 suitable for the preparation of antibodies by manipulation of DNA, creation of expression vectors, and transformation of appropriate cells (Mountain A and Adair, J R in *Biotechnology and Genetic Engineering Reviews* (ed. Tombs, M P, 10, Chapter 1, 1992, Intercept, Andover, UK); International Patent Specification No. WO 91/09967).

In certain embodiments, the antibody according to the invention may have one or  
15 more effector or reporter molecules attached to it. A reporter molecule may be a detectable moiety or label such as an enzyme, cytotoxic agent or other reporter molecule, including a dye, radionuclide, luminescent group, fluorescent group, or biotin, or the like. The TGF-beta binding protein-specific immunoglobulin or fragment thereof may be radiolabeled for diagnostic or therapeutic applications. Techniques for radiolabeling of antibodies are known in the art. *See,*  
20 *e.g.*, Adams 1998 *In Vivo* 12:11-21; Hiltunen 1993 *Acta Oncol.* 32:831-9. Therapeutic applications are described in greater detail below and may include use of the TGF-beta binding protein specific antibody (or fragment thereof) in conjunction with other therapeutic agents. The effector or reporter molecules may be attached to the antibody through any available amino acid side-chain, terminal amino acid, or carbohydrate functional group located in the antibody, provided  
25 that the attachment or attachment process does not adversely affect the binding properties such that the usefulness of the molecule is abrogated. Particular functional groups include, for example any free amino, imino, thiol, hydroxyl, carboxyl, or aldehyde group. Attachment of the antibody and the effector and/or reporter molecule(s) may be achieved via such groups and an appropriate functional group in the effector or reporter molecule. The linkage may be direct or indirect through  
30 spacing or bridging groups.

Effector molecules include, for example, antineoplastic agents, toxins (such as enzymatically active toxins of bacterial (such as *Pseudomonas aeruginosa* exotoxin A) or plant origin and fragments thereof (*e.g.*, ricin and fragments thereof, plant gelonin, bryodin from

*Bryonia dioica*, or the like. See, e.g., Thrush et al., 1996 *Annu. Rev. Immunol.* 14:49-71; Frankel et al., 1996 *Cancer Res.* 56:926-32). Additional effector molecules include biologically active proteins (for example, enzymes), nucleic acids and fragments thereof, naturally occurring and synthetic polymers, for example, polysaccharides and polyalkylene polymers such as poly(ethylene glycol) and derivatives thereof, radionuclides, particularly radioiodide, and chelated metals. Suitable reporter groups include chelated metals, fluorescent compounds or compounds that may be detected by NMR or ESR spectroscopy. Particularly useful effector groups are calichaemicin and derivatives thereof (see, for example, South African Patent Specifications Nos. 85/8794, 88/8127 and 90/2839).

Numerous other toxins, including chemotherapeutic agents, anti-mitotic agents, antibiotics, inducers of apoptosis (or "apoptogens", see, e.g., Green and Reed, 1998, *Science* 281:1309-1312), or the like, are known to those familiar with the art, and the examples provided herein are intended to be illustrative without limiting the scope and spirit of the invention. Particular antineoplastic agents include cytotoxic and cytostatic agents, for example alkylating agents, such as nitrogen mustards (e.g., chlorambucil, melphalan, mechlorethamine, cyclophosphamide, or uracil mustard) and derivatives thereof, triethylenephosphoramidate, triethylenethiophosphoramide, busulphan, or cisplatin; antimetabolites, such as methotrexate, fluorouracil, floxuridine, cytarabine, mercaptopurine, thioguanine, fluoroacetic acid or fluorocitric acid, antibiotics, such as bleomycins (e.g., bleomycin sulphate), doxorubicin, daunorubicin, mitomycins (e.g., mitomycin C), actinomycins (e.g., dactinomycin), plicamycin, calichaemicin and derivatives thereof, or esperamicin and derivatives thereof; mitotic inhibitors, such as etoposide, vincristine or vinblastine and derivatives thereof; alkaloids, such as ellipticine; polyols such as taxicin-I or taxicin-II; hormones, such as androgens (e.g., dromostanolone or testolactone), progestins (e.g., megestrol acetate or medroxyprogesterone acetate), estrogens (e.g., dimethylstilbestrol diphosphate, polyestradiol phosphate or estramustine phosphate) or antiestrogens (e.g., tamoxifen); anthraquinones, such as mitoxantrone, ureas, such as hydroxyurea; hydrazines, such as procarbazine; or imidazoles, such as dacarbazine.

Chelated metals useful as effector molecules include chelates of di- or tripositive metals having a coordination number from 2 to 8 inclusive. Particular examples of such metals include technetium (Tc), rhenium (Re), cobalt (Co), copper (Cu), gold (Au), silver (Ag), lead (Pb), bismuth (Bi), indium (In), gallium (Ga), yttrium (Y), terbium (Tb), gadolinium (Gd), and scandium (Sc). In general the metal is preferably a radionuclide. Particular radionuclides include  $^{99m}\text{Tc}$ ,  $^{186}\text{Re}$ ,  $^{188}\text{Re}$ ,  $^{58}\text{Co}$ ,  $^{60}\text{Co}$ ,  $^{67}\text{Cu}$ ,  $^{195}\text{Au}$ ,  $^{199}\text{Au}$ ,  $^{110}\text{Ag}$ ,  $^{203}\text{Pb}$ ,  $^{206}\text{Bi}$ ,  $^{207}\text{Bi}$ ,  $^{111}\text{In}$ ,  $^{67}\text{Ga}$ ,  $^{68}\text{Ga}$ ,

<sup>88</sup>Y, <sup>90</sup>Y, <sup>160</sup>Tb, <sup>153</sup>Gd, and <sup>47</sup>Sc. The chelated metal may be for example one of the above types of metal chelated with any suitable polydentate chelating agent, for example acyclic or cyclic polyamines, polyethers, (e.g., crown ethers and derivatives thereof), polyamides; porphyrins; and carbocyclic derivatives. In general, the type of chelating agent will depend on the metal in use.

5 One particularly useful group of chelating agents in conjugates according to the invention, however, comprises acyclic and cyclic polyamines, especially polyaminocarboxylic acids, for example diethylenetriaminepentaacetic acid and derivatives thereof, and macrocyclic amines, e.g. cyclic tri-aza and tetra-aza derivatives (for example, as described in International Patent Specification No. WO 92/22583); and polyamides, especially desferrioxamine and derivatives  
10 thereof.

When a thiol group in the antibody is used as the point of attachment, the effector molecule may be attached according to a reaction that occurs with a thiol-reactive group present in the effector or reporter molecule. Examples of such groups include an  $\alpha$ -halocarboxylic acid or ester, such as iodoacetamide, an imide, such as maleimide, a vinyl sulphone, or a disulphide. These  
15 and other suitable linking procedures are generally and more particularly described in International Patent Specifications Nos. WO 93/06231, WO 92/22583, WO 90/091195, and WO 89/01476.

The invention also contemplates the generation of anti-idiotypic antibodies that recognize an antibody (or antigen-binding fragment thereof) that specifically binds to a SOST polypeptide as provided herein, or a variant or fragment thereof. Anti-idiotypic antibodies may  
20 be generated as polyclonal antibodies or as monoclonal antibodies by the methods described herein, using an anti-TGF-beta binding protein antibody (or antigen-binding fragment thereof) as immunogen. Anti-idiotypic antibodies or fragments thereof may also be generated by any of the recombinant genetic engineering methods described above, or by phage display selection. An anti-idiotypic antibody may react with the antigen binding site of the anti-SOST antibody  
25 such that binding of the antibody to a SOST polypeptide is competitively inhibited. Alternatively, an anti-idiotypic antibody as provided herein may not competitively inhibit binding of an anti-TGF-beta binding protein antibody to a TGF-beta binding protein.

In certain embodiments of the invention, an antibody that specifically binds to SOST may be used to detect expression of the SOST polypeptide. In certain particular  
30 embodiments, one antibody or a panel of antibodies may be exposed to cells that express a SOST polypeptide and expression of the SOST polypeptide may be determined by detection using another SOST specific antibody that binds to a different epitope than the antibody or antibodies first used in the assay.

## ASSAYS FOR SELECTING AGENTS THAT INCREASE BONE DENSITY

As discussed above, the present invention provides methods for selecting and/or isolating agents that are capable of increasing bone density. In certain preferred embodiments, the agent is an antibody that specifically binds to SOST (a TGF-beta binding protein) or a  
5 variant or a fragment thereof. For example, within one embodiment of the present invention, methods are provided for determining whether an agent is capable of increasing bone mineral content, comprising the steps of (a) contacting or mixing a candidate agent with a TGF-beta binding protein and a member of the TGF-beta family of proteins and (b) determining whether the candidate agent stimulates signaling by the TGF-beta family of proteins, or impairs or  
10 inhibits the binding of the TGF-beta binding protein to one or more members of the TGF-beta family of proteins. Within certain embodiments, the agent enhances the ability of TGF-beta to function as a positive regulator of mesenchymal cell differentiation.

In a preferred embodiment of the invention, a method is provided for identifying an antibody that modulates a TGF-beta signaling pathway comprising contacting an antibody  
15 that specifically binds to a SOST polypeptide with a SOST peptide, including but not limited to the peptides disclosed herein, under conditions and for a time sufficient to permit formation of an antibody plus (+) SOST (antibody/SOST) complex and then detecting the level (*e.g.*, quantifying the amount) of the SOST/antibody complex to determine the presence of an antibody that modulates a TGF-beta signaling pathway. The method may be performed using  
20 SPR or any number of different immunoassays known in the art and disclosed herein, including an ELISA, immunoblot, or the like. A TGF-beta signaling pathway includes a signaling pathway by which a BMP binds to a type I and a type II receptor on a cell to stimulate or induce the pathway that modulates bone mineral content. In certain preferred embodiments of the invention, an antibody that specifically binds to SOST stimulates or enhances the pathway for  
25 increasing bone mineral content. Such an antibody may be identified using the methods disclosed herein to detect binding of an antibody to SOST specific peptides.

The subject invention methods may also be used for identifying antibodies that impair, inhibit (including competitively inhibit), or prevent binding of a BMP to a SOST polypeptide by detecting whether an antibody binds to SOST peptides that are located in regions  
30 or portions of regions on SOST to which a BMP binds, such as peptides at the amino terminal end of SOST and peptides that include amino terminal amino acid residues and a portion of the core region (docking core) of SOST (*e.g.*, SEQ ID NOs:2-19, 21-28, and 47-50). The methods of the present invention may also be used to identify an antibody that impairs, prevents, or

inhibits, formation of SOST homodimers. Such an antibody that binds specifically to SOST may be identified by detecting binding of the antibody to peptides that are derived from the core or the carboxy terminal region of SOST (e.g., SEQ ID NOs: 29-46 and 51-54).

Within other embodiments of the invention, methods are provided for  
5 determining whether an agent is capable of increasing bone mineral content, comprising the steps of (a) contacting a candidate agent to cells that express a TGF-beta binding-protein and (b) determining whether the expression of the TGF-beta binding-protein in the exposed cells decreases or whether an activity of the TGF-beta binding protein decreases, and thereby determining whether the compound is capable of increasing bone mineral content. Within one  
10 embodiment, the cells may include spontaneously transformed or untransformed normal human bone from bone biopsies or rat parietal bone osteoblasts.

Immunoassays may be used for detecting and quantifying the expression of a TGF-beta binding protein and include, for example, Countercurrent Immuno-Electrophoresis (CIEP), radioimmunoassays, radioimmunoprecipitations, Enzyme-Linked Immuno-Sorbent  
15 Assays (ELISA), immunoblot assays such as dot blot assays and Western blots, inhibition or competition assays, and sandwich assays (see U.S. Patent Nos. 4,376,110 and 4,486,530; see also *Antibodies: A Laboratory Manual, supra*). Such immunoassays may use an antibody that is specific for a TGF-beta binding protein such as the anti-sclerostin antibodies described herein, or may use an antibody that is specific for a reporter molecule that is attached to the TGF-beta  
20 binding protein. The level of polypeptide expression may also be determined by quantifying the amount of TGF-beta binding protein that binds to a TGF-beta binding protein ligand. By way of example, binding of sclerostin in a sample to a BMP may be detected by surface plasmon resonance. Alternatively, the level of expression of mRNA encoding the specific TGF-beta binding protein may be quantified.

25 In other embodiments, an antibody specific for SOST may be used in a method, such as a competition assay, in which candidate agents are screened to identify an agent that will compete with the antibody for binding to the TGF-binding protein. The interaction between an antibody and a TGF-beta protein may be determined using immunoassay methods known in the art and described herein.

30 By way of example, a family member of the TGF-beta superfamily, such as a TGF-beta binding protein or a TGF-beta superfamily member that is a ligand of TGF-beta binding protein, is first bound to a solid phase, followed by addition of a candidate agent. A ligand of the TGF-beta superfamily member that is bound to the solid phase is added

concurrently or added in a subsequent step, the solid phase washed, and the quantity of ligand that bound to the family member determined, which indicates whether the candidate agent blocks binding of the ligand to the TGF-beta member. Alternatively, the ligand of the TGF-beta superfamily member may be attached to the solid phase, and the TGF-beta superfamily member may be added concurrently or subsequently to the addition of the candidate molecule. The amount of ligand that binds to the TGF-beta superfamily member can be measured by labeling the ligand with a detectable molecule according to methods known in the art and described herein. Alternatively, the binding of the ligand to the TGF-beta superfamily member may be measured by adding one or more detecting molecules, such as an antibody or fragment thereof that specifically binds to the ligand and quantifying the amount of the molecule that binds to the ligand. Agents that are suitable for use in increasing bone mineral content as described herein are those agents that decrease the binding of TGF-beta binding protein to its ligand that is a member of the TGF-beta superfamily in a statistically significant manner.

Within other embodiments of the invention, methods are provided for determining whether an agent is capable of increasing bone mineral content, comprising the steps of (a) contacting a candidate agent to cells that express a TGF-beta protein and (b) determining whether the activity of TGF-beta protein from the exposed cells is altered, and therefrom determining whether the compound is capable of increasing bone mineral content. Similar to the methods described herein, a wide variety of methods may be used to assess changes in TGF-beta binding-protein expression when a cell producing the TGF-beta binding protein is exposed to a selected test compound.

Within another embodiment of the present invention, methods are provided for determining whether an agent is capable of increasing bone mineral content, comprising the steps of (a) contacting a candidate agent with a TGF-beta-binding-protein and a selected member of the TGF-beta family of proteins, and (b) determining whether the selected agent up-regulates the signaling of the TGF-beta family of proteins or inhibits the binding of the TGF-beta binding-protein to the TGF-beta family member. Within certain embodiments, the molecule enhances the ability of a TGF-beta family member to function as a positive regulator of mesenchymal cell differentiation.

Similar to the above-described methods, a variety of methods may be used to assess stimulation of a TGF-beta family member by a test compound, such as an antibody that specifically binds to SOST. One such representative method (*see also* Durham et al., *Endo.* 136:1374-1380) comprises attaching a TGF-beta family member, such as a BMP (e.g., BMP-2,

BMP-4, BMP-5, BMP-6, BMP-7) to the solid phase at a concentration that is equivalent to its  $K_d$ . The dissociation constant ( $K_d$ ) may be determined according to methods for measuring binding rate constants known in the art, such as surface plasmon resonance using a BIAcore instrument according to techniques known in the art and described by the manufacturer (Biosensor, Piscataway, NJ). A collection of antagonist candidate agents is then added at a fixed concentration (typically, 20  $\mu$ M for small organic molecule collections (an isolated organic small molecule as used herein means that the molecule is greater than 90% pure as determined according to methods well known in the art (*e.g.*, NMR, melting point) and 1  $\mu$ M for candidate antibodies). An agent is considered an antagonist of this interaction when the agent inhibits binding of the BMP to SOST, that is, a decrease in the level of binding is observed, by at least 40%, preferably 50% or 60%, and more preferably 80% or 90% compared to the level of BMP binding to SOST that is observed in the absence of the agent. Such an agent may be further evaluated as a potential inhibitor on the basis of titration studies according to which its inhibition constant and its influence on TGF-beta binding-protein binding affinity is determined. Comparable specificity control assays may also be conducted to establish the selectivity profile for the identified antagonist according to studies using assays dependent on the BMP ligand action (*e.g.* BMP/BMP receptor competition study).

Within yet other embodiments of the present invention, methods are provided for determining whether a candidate agent is capable of increasing bone mineral content, comprising the step of determining whether a candidate agent inhibits the binding of TGF-beta binding-protein to bone, or an analogue thereof. As used herein, it should be understood that bone or analogues thereof refers to hydroxyapatite, or a surface composed of a powdered form of bone, crushed bone or intact bone. Similar to the methods described herein, a wide variety of methods may be used to assess the inhibition of TGF-beta binding-protein localization to bone matrix (*see, e.g.*, Nicolas et al., *Calcif. Tissue Int.* 47:206-12 (1995)).

While the methods described herein may refer to the analysis of an individual candidate agent, the present invention should not be so limited. In particular, the agent may be contained within a mixture of candidate agents. As described herein, the candidate agents may be small organic molecules or may be antibodies or fragments thereof or other peptides or polypeptides. Antibodies or fragments may be identified by screening a library of antibodies or antibody fragments or prepared by methods described herein. Hence, the recited methods may further comprise the step of isolating an agent that inhibits the binding of TGF-beta binding-protein to a TGF-beta family member.



In one embodiment of the invention, an antibody or antigen-binding fragment thereof that specifically binds to a SOST polypeptide is capable of competitively inhibiting binding of a TGF-beta family member to the SOST polypeptide. The capability of the antibody or antibody fragment to impair or blocking binding of a TGF-beta family member, such as a BMP, to SOST may be determined according to any of the methods described herein. The antibody or fragment thereof that specifically binds to SOST may impair, block, or prevent binding of a TGF-beta family member to SOST by impairing SOST homodimer formation. An antibody that specifically binds to SOST may also be used to identify an activity of SOST by inhibiting or impairing SOST from binding to a BMP. Alternatively, the antibody or fragment thereof may be incorporated in a cell-based assay or in an animal model in which SOST has a defined activity to determine whether the antibody alters (increases or decreases in a statistically significant manner) that activity. An antibody or fragment thereof that specifically binds to SOST may be used to examine the effect of such an antibody in a signal transduction pathway and thereby modulate (stimulate or inhibit) the signaling pathway. Preferably, binding of an antibody to SOST results in a stimulation or induction of a signaling pathway.

#### PRODUCTION OF PROTEINS

Polypeptides described herein include the TGF binding protein sclerostin and variants thereof and antibodies or fragments thereof that specifically bind to sclerostin. The polynucleotides that encode these polypeptides include derivatives that are substantially similar to the polynucleotides (and, when appropriate, to the proteins including peptides and polypeptides that are encoded by the polynucleotides and their derivatives). As used herein, a nucleotide sequence is deemed to be "substantially similar" if (a) the nucleotide sequence is derived from the coding region of the polynucleotides described herein and includes, for example, portions of the sequence or allelic variations of the sequences discussed above, or alternatively, encodes a molecule that inhibits the binding of TGF-beta binding-protein to a member of the TGF-beta family; (b) the polynucleotide is capable of hybridizing to polynucleotide as provided herein under moderate, high or very high stringency (*see* Sambrook et al., *Molecular Cloning: A Laboratory Manual*, 3rd ed., Cold Spring Harbor Laboratory Press, NY, 2001); and/or (c) the polynucleotide sequences are degenerate with respect to codon sequences for a particular amino acid of the polynucleotide sequences described in (a) or (b). Further, the nucleic acid molecule disclosed herein includes both complementary and non-complementary sequences, provided the sequences otherwise meet the criteria set forth herein.

Within the context of the present invention, high stringency means standard hybridization conditions (e.g., 5XSSPE, 0.5% SDS at 55-65° C, or the equivalent; 5xSSPE (1xSSPE = 180 mM sodium chloride; 10 mM sodium phosphate; 1 mM EDTA (pH 7.7); 5xDenhardt's solution (100x Denhardt's = 2% (w/v) bovine serum albumin, 2% (w/v) Ficoll, 2% (w/v) polyvinylpyrrolidone); and 0.5% SDS). Post-hybridization washes at high stringency are typically performed in 0.5xSSC (1xSSC = 150 mM sodium chloride, 15 mM trisodium citrate) or in 0.5xSSPE at 55-60° C.).

Polynucleotides encoding SOST may be isolated from genomic or cDNA libraries, from cells in a biological sample, tissues, or cell lines, and prepared using any variety of techniques (see, e.g., Sambrook, *supra*). For example, a polynucleotide may be amplified from cDNA prepared from a suitable cell or tissue. These polynucleotides may be amplified by polymerase chain reaction (PCR) using sequence specific primers that are designed on the basis of the sequences provided herein, and which may be purchased or synthesized. Polynucleotides may also be obtained by screening cDNA or genomic DNA libraries using sequence specific probes and primers. General methods for screening libraries by PCR are provided by, for example, Yu et al., "Use of the Polymerase Chain Reaction to Screen Phage Libraries," in *Methods in Molecular Biology, Vol. 15: PCR Protocols: Current Methods and Applications*, White (ed.), pages 211-215 (Humana Press, Inc. 1993). Moreover, techniques for using PCR to isolate related genes are described by, for example, Preston, "Use of Degenerate Oligonucleotide Primers and the Polymerase Chain Reaction to Clone Gene Family Members," in *Methods in Molecular Biology, Vol. 15: PCR Protocols: Current Methods and Applications*, White (ed.), pages 317-337 (Humana Press, Inc. 1993).

The sequence of a TGF-beta binding-protein cDNA or TGF-beta binding-protein genomic fragment can be determined using standard methods. Moreover, the identification of genomic fragments containing a TGF-beta binding-protein promoter or regulatory element can be achieved using well-established techniques, such as deletion analysis (see, generally, Ausubel (1995)).

As an alternative, a polynucleotide encoding a TGF-beta binding-protein can be obtained by synthesizing DNA molecules using mutually priming long oligonucleotides and the nucleotide sequences described herein (see, e.g., Ausubel (1995) at pages 8-8 to 8-9). If PCR is incorporated into the method DNA molecules at least two kilobases in length can be synthesized (Adang et al., *Plant Molec. Biol.* 21:1131, 1993; Bambot et al., *PCR Methods and Applications* 2:266, 1993; Dillon et al., "Use of the Polymerase Chain Reaction for the Rapid Construction of

Synthetic Genes," in *Methods in Molecular Biology, Vol. 15: PCR Protocols: Current Methods and Applications*, White (ed.), pages 263-268, (Humana Press, Inc. 1993); Holowachuk et al., *PCR Methods Appl.* 4:299, 1995).

Nucleic acid molecules encoding a variant TGF-beta binding-protein can be  
5 obtained by screening various cDNA or genomic libraries with polynucleotide probes having  
nucleotide sequences based upon SEQ ID NO:55-57, 59, 61, 63, 65, 67, and 69, using  
procedures described herein. TGF-beta binding-protein polynucleotide variants can also be  
constructed synthetically. For example, a nucleic acid molecule can be designed that encodes a  
polypeptide having a conservative amino acid change when compared with the amino acid  
10 sequence of SEQ ID NOs: 1, 20, 58, 60, 62, 64, 66, 68, or 70. That is, variants can be obtained  
that contain one or more amino acid substitutions of SEQ ID NOs: 1, 20, 58, 60, 62, 64, 66, 68,  
or 70, in which an alkyl amino acid is substituted for an alkyl amino acid in a TGF-beta binding-  
protein amino acid sequence; an aromatic amino acid is substituted for an aromatic amino acid  
in a TGF-beta binding-protein amino acid sequence; a sulfur-containing amino acid is  
15 substituted for a sulfur-containing amino acid in a TGF-beta binding-protein amino acid  
sequence; a hydroxy-containing amino acid is substituted for a hydroxy-containing amino acid  
in a TGF-beta binding-protein amino acid sequence; an acidic amino acid is substituted for an  
acidic amino acid in a TGF-beta binding-protein amino acid sequence; a basic amino acid is  
substituted for a basic amino acid in a TGF-beta binding-protein amino acid sequence; or a  
20 dibasic monocarboxylic amino acid is substituted for a dibasic monocarboxylic amino acid in a  
TGF-beta binding-protein amino acid sequence.

Among the common amino acids, for example, a conservative amino acid  
substitution is illustrated by a substitution among amino acids within each of the following  
groups: (1) glycine, alanine, valine, leucine, and isoleucine (non-polar alkyl group-containing  
25 side chain); (2) phenylalanine, tyrosine, and tryptophan (aromatic side chain) (3) serine and  
threonine (hydroxyl group side chain), (4) aspartate and glutamate (carboxylic acid group side  
chain); (5) glutamine and asparagines (amide group-containing side chain) and (6) lysine,  
arginine and histidine (amino group side chain). In making such substitutions, it is important to,  
where possible, maintain the cysteine backbone outlined in Figure 1.

30 Conservative amino acid changes in SOST can be introduced by substituting  
nucleotides for the nucleotides recited in any one of SEQ ID NOs: 55-57, 59, 61, 63, 65, 67, or  
69. Such conservative amino acid variants can be obtained, for example, by oligonucleotide-  
directed mutagenesis, linker-scanning mutagenesis, mutagenesis using the polymerase chain

reaction, and the like (*see* Ausubel (1995) at pages 8-10 to 8-22; and McPherson (ed.), *Directed Mutagenesis: A Practical Approach* (IRL Press 1991)). The functional activity of such variants can be determined using standard methods and assays described herein. Alternatively, a variant TGF-beta binding-protein polypeptide retains the ability to specifically bind to anti-SOST antibodies.

Routine deletion analyses of nucleic acid molecules can be performed to obtain "functional fragments" of a nucleic acid molecule that encodes a TGF-beta binding-protein polypeptide. By way of illustration, DNA molecules having the nucleotide sequence of any one of SEQ ID NOs:55-57, 59, 61, 63, 65, 67, or 69 can be digested with *Bal31* nuclease to obtain a series of nested deletions. The fragments are then inserted into expression vectors in proper reading frame, and the expressed polypeptides are isolated and tested for activity, or for the ability to bind anti-TGF-beta binding-protein antibodies. One alternative to exonuclease digestion is to use oligonucleotide-directed mutagenesis to introduce deletions or stop codons to specify production of a desired fragment. Alternatively, particular fragments of a TGF-beta binding-protein encoding polynucleotide can be synthesized using the polymerase chain reaction. In addition to assays and methods disclosed herein, standard techniques for functional analysis of proteins are described by, for example, Treuter et al., *Molec. Gen. Genet.* 240:113, 1993; Content et al., "Expression and preliminary deletion analysis of the 42 kDa 2-5A synthetase induced by human interferon," in *Biological Interferon Systems, Proceedings of ISIR-TNO Meeting on Interferon Systems*, Cantell (ed.), pages 65-72 (Nijhoff 1987); Herschman, "The EGF Receptor," in *Control of Animal Cell Proliferation, Vol. 1*, Boynton et al., (eds.) pages 169-199 (Academic Press 1985); Coumailleau et al., *J. Biol. Chem.* 270:29270, 1995; Fukunaga et al., *J. Biol. Chem.* 270:25291, 1995; Yamaguchi et al., *Biochem. Pharmacol.* 50:1295, 1995; and Meisel et al., *Plant Molec. Biol.* 30:1, 1996.

The structure of the polypeptides encoded by the nucleic acid molecules described herein may be predicted from the primary translation products using the hydrophobicity plot function of, for example, P/C Gene or Intelligenetics Suite (Intelligenetics, Mountain View, California), or according to the methods described by Kyte and Doolittle (*J. Mol. Biol.* 157:105-132, 1982). Polypeptides may be prepared in the form of acidic or basic salts or in neutral form. In addition, individual amino acid residues may be modified by oxidation or reduction. Furthermore, various substitutions, deletions, or additions may be made to the amino acid or nucleic acid sequences, the net effect of which is to retain or further enhance or decrease the biological activity of the mutant or wild-type protein. Preferably, a

SOST variant, or a fragment thereof retains or has enhanced ability to bind to a SOST specific antibody. Moreover, due to degeneracy in the genetic code, for example, the nucleotide sequences encoding the same amino acid sequence may vary considerably.

Other derivatives of the proteins provided herein include conjugates of the  
5 proteins with other proteins or polypeptides. This may be accomplished, for example, by the synthesis of N-terminal or C-terminal fusion proteins that may be added to facilitate purification or identification of proteins (*see* U.S. Patent No. 4,851,341, *see also*, Hopp et al., *Bio/Technology* 6:1204, 1988.) Alternatively, fusion proteins such as FLAG®/TGF-beta binding-protein may be constructed in order to assist in the identification, expression, and  
10 analysis of the protein.

Proteins described herein may be constructed using a wide variety of techniques described herein. Further, mutations may be introduced at particular loci by synthesizing oligonucleotides that contain a mutant sequence that are flanked by restriction sites, enabling ligation to fragments of the native sequence. Following ligation, the resulting reconstructed  
15 sequence encodes a derivative having the desired amino acid insertion, substitution, or deletion.

Alternatively, oligonucleotide-directed site-specific (or segment specific) mutagenesis procedures may be employed to provide an altered polynucleotide having particular codons altered according to the substitution, deletion, or insertion. Exemplary methods of making the alterations set forth above are disclosed by Walder et al. (*Gene* 42:133, 1986); Bauer  
20 et al. (*Gene* 37:73, 1985); Craik (*BioTechniques*, January 1985, 12-19); Smith et al. (*Genetic Engineering: Principles and Methods*, Plenum Press, 1981); and Sambrook et al. (*supra*). Deletion or truncation derivatives of proteins (*e.g.*, a soluble extracellular portion) may also be constructed by using convenient restriction endonuclease sites adjacent to the desired deletion. Subsequent to restriction, overhangs may be filled in and the DNA religated. Exemplary  
25 methods of making the alterations set forth above are disclosed by Sambrook et al. (*Molecular Cloning: A Laboratory Manual*, 3d Ed., Cold Spring Harbor Laboratory Press (2001)).

Mutations that are made in the nucleic acid molecules preferably preserve the reading frame of the coding sequences. Furthermore, the mutations will preferably not create complementary regions that when transcribed could hybridize to produce secondary mRNA  
30 structures, such as loops or hairpins, which would adversely affect translation of the mRNA. Although a mutation site may be predetermined, it is not necessary that the nature of the mutation *per se* be predetermined. For example, in order to select for optimum characteristics of mutants at a given site, random mutagenesis may be conducted at the target codon and the

expressed mutants screened for gain, loss, or retention of biological activity. Alternatively, mutations may be introduced at particular loci by synthesizing oligonucleotides containing a mutant sequence, flanked by restriction sites enabling ligation to fragments of the native sequence. Following ligation, the resulting reconstructed sequence encodes a derivative having the desired amino acid insertion, substitution, or deletion. Nucleic acid molecules that encode proteins of the present invention may also be constructed using techniques such as polymerase chain reaction (PCR) mutagenesis, chemical mutagenesis (Drinkwater and Klinedinst, *PNAS* 83:3402-3406, 1986), forced nucleotide misincorporation (*e.g.*, Liao and Wise *Gene* 88:107-111, 1990), or use of randomly mutagenized oligonucleotides (Horwitz et al., *Genome* 3:112-117, 1989).

The present invention also provides for the manipulation and expression of the polynucleotides provided herein and of polynucleotides encoding the subject invention antibodies by culturing host cells containing a vector capable of expressing such polynucleotides. An expression vector refers to recombinant nucleic acid construct that is capable of directing the expression of desired protein. The vector may be composed of deoxyribonucleic acids (DNA), synthetic or cDNA-derived, ribonucleic acids (RNA), or a combination of the two (*e.g.*, a DNA-RNA chimera). A cDNA polynucleotide is generally understood to mean a DNA polynucleotide prepared by transcribing an RNA molecule, such as mRNA. These vectors or vector constructs that include a polynucleotide sequence encoding the desired protein preferably are operably linked to suitable transcriptional or translational regulatory elements. Suitable regulatory elements may be derived from a variety of sources, including bacterial, fungal, viral, mammalian, insect, or plant genes. Selection of appropriate regulatory elements is dependent on the host cell chosen, and may be readily accomplished by one of ordinary skill in the art. Examples of regulatory elements include a transcriptional promoter and enhancer or RNA polymerase binding sequence, a transcriptional terminator, and a ribosomal binding sequence including a translation initiation signal. Optionally, the vector may include a polyadenylation sequence, one or more restriction sites, as well as one or more selectable markers such as neomycin phosphotransferase or hygromycin phosphotransferase or any other markers known in the art. Additionally, depending on the host cell chosen and the vector employed, other genetic elements such as an origin of replication, additional nucleic acid restriction sites, enhancers, sequences conferring inducibility of transcription, and selectable markers, may also be incorporated into the vectors described herein.

Nucleic acid molecules that encode any of the proteins described above may be

readily expressed by a wide variety of prokaryotic and eukaryotic host cells, including bacterial, mammalian, yeast or other fungi, viral, insect, or plant cells. Methods for transforming or transfecting such cells to express foreign DNA are well known in the art (*see, e.g., Itakura et al., U.S. Patent No. 4,704,362; Hinnen et al., Proc. Natl. Acad. Sci. USA 75:1929-1933, 1978; Murray et al., U.S. Patent No. 4,801,542; Upshall et al., U.S. Patent No. 4,935,349; Hagen et al., U.S. Patent No. 4,784,950; Axel et al., U.S. Patent No. 4,399,216; Goeddel et al., U.S. Patent No. 4,766,075; and Sambrook et al. Molecular Cloning: A Laboratory Manual, 3rd ed., Cold Spring Harbor Laboratory Press, 2001; for plant cells see Czako and Marton, Plant Physiol. 104:1067-1071, 1994; and Paszkowski et al., Biotech. 24:387-392, 1992).*

Bacterial host cells suitable for carrying out the present invention include *E. coli* BL21(DE3), BL21(DE3)pLysS, BL21(DE3)pLysE, DH1, DH4I, DH5, DH5I, DH5IF', DH5IMCR, DH10B, DH10B/p3, DH11S, C600, HB101, JM101, JM105, JM109, JM110, K38, RR1, Y1088, Y1089, CSH18, ER1451, and ER1647 (*see, for example, Brown (Ed.), Molecular Biology Labfax (Academic Press 1991); B. subtilis (BR151, YB886, MI119, MI120, and B170 (see, for example, Hardy, "Bacillus Cloning Methods," in DNA Cloning: A Practical Approach, Glover (Ed.) (IRL Press 1985); Salmonella typhimurium; and various species within the genera Pseudomonas, Streptomyces, and Staphylococcus, as well as many other bacterial species well known to one of ordinary skill in the art. A representative example of a bacterial host cell includes E. coli DH5 $\alpha$  (Stratagene, LaJolla, California).*

Bacterial expression vectors preferably comprise a promoter that functions in the host cell, one or more selectable phenotypic markers, and a bacterial origin of replication. Representative promoters include the  $\beta$ -lactamase (penicillinase) and lactose promoter system (*see Chang et al., Nature 275:615, 1978*), the T7 RNA polymerase promoter (*Studier et al., Meth. Enzymol. 185:60-89, 1990*), the lambda promoter (*Elvin et al., Gene 87:123-126, 1990*), the *trp* promoter (*Nichols and Yanofsky, Meth. in Enzymology 101:155, 1983*), and the *tac* promoter (*Russell et al., Gene 20:231, 1982*). Additional promoters include promoters capable of recognizing the T4, T3, Sp6 and T7 polymerases, the P<sub>R</sub> and P<sub>L</sub> promoters of bacteriophage lambda, the *recA*, heat shock, *lacUV5*, *tac*, *lpp-lacSpr*, *phoA*, and *lacZ* promoters of *E. coli*, promoters of *B. subtilis*, the promoters of the bacteriophages of *Bacillus*, *Streptomyces* promoters, the *int* promoter of bacteriophage lambda, the *bla* promoter of pBR322, and the CAT promoter of the chloramphenicol acetyl transferase gene. Prokaryotic promoters have been reviewed by Glick, *J. Ind. Microbiol. 1:277, 1987*, Watson et al., *Molecular Biology of the Gene, 4th Ed. (Benjamin Cummins 1987)*, and by Ausubel et al. (1995). Representative

selectable markers include various antibiotic resistance markers such as the kanamycin or ampicillin resistance genes. Many plasmids suitable for transforming host cells are well known in the art, including among others, pBR322 (see Bolivar et al., *Gene* 2:95, 1977), the pUC plasmids pUC18, pUC19, pUC118, pUC119 (see Messing, *Meth. in Enzymology* 101:20-77, 1983 and Vieira and Messing, *Gene* 19:259-268, 1982), and pNH8A, pNH16a, pNH18a, and Bluescript M13 (Stratagene, La Jolla, California).

Yeast and fungi host cells suitable for carrying out the present invention include, among others, *Saccharomyces pombe*, *Saccharomyces cerevisiae*, the genera *Pichia* or *Kluyveromyces* and various species of the genus *Aspergillus* (McKnight et al., U.S. Patent No. 4,935,349). Suitable expression vectors for yeast and fungi include, among others, YCp50 (ATCC No. 37419) for yeast, and the amdS cloning vector pV3 (Turnbull, *Bio/Technology* 7:169, 1989), YRp7 (Struhl et al., *Proc. Natl. Acad. Sci. USA* 76:1035-1039, 1978), YEpl3 (Broach et al., *Gene* 8:121-133, 1979), pJDB249 and pJDB219 (Beggs, *Nature* 275:104-108, 1978) and derivatives thereof.

Preferred promoters for use in yeast include promoters from yeast glycolytic genes (Hitzeman et al., *J. Biol. Chem.* 255:12073-12080, 1980; Alber and Kawasaki, *J. Mol. Appl. Genet.* 1:419-434, 1982) or alcohol dehydrogenase genes (Young et al., in *Genetic Engineering of Microorganisms for Chemicals*, Hollaender et al. (eds.), p. 355, Plenum, New York, 1982; Ammerer, *Meth. Enzymol.* 101:192-201, 1983). Examples of useful promoters for fungi vectors include those derived from *Aspergillus nidulans* glycolytic genes, such as the *adh3* promoter (McKnight et al., *EMBO J.* 4:2093-2099, 1985). The expression units may also include a transcriptional terminator. An example of a suitable terminator is the *adh3* terminator (McKnight et al., *supra*, 1985).

As with bacterial vectors, the yeast vectors will generally include a selectable marker, which may be one of any number of genes that exhibit a dominant phenotype for which a phenotypic assay exists to enable transformants to be selected. Preferred selectable markers are those that complement host cell auxotrophy, provide antibiotic resistance, or enable a cell to utilize specific carbon sources, and include *leu2* (Broach et al., *ibid.*), *ura3* (Botstein et al., *Gene* 8:17, 1979), or *his3* (Struhl et al., *ibid.*). Another suitable selectable marker is the *cat* gene, which confers chloramphenicol resistance on yeast cells. Many yeast cloning vectors have been designed and are readily available. These vectors include YIp-based vectors such as YIp5, YRp vectors such as YRp17, YEpl vectors such as YEpl3, and YCp vectors such as YCp19. One skilled in the art will appreciate that a wide variety of suitable vectors are available for



expression in yeast cells.

Techniques for transforming fungi are well known in the literature and have been described, for instance, by Beggs (*ibid.*), Hinnen et al. (*Proc. Natl. Acad. Sci. USA* 75:1929-1933, 1978), Yelton et al. (*Proc. Natl. Acad. Sci. USA* 81:1740-1747, 1984), and Russell (*Nature* 301:167-169, 1983). The genotype of the host cell may contain a genetic defect that is complemented by the selectable marker present on the expression vector. Choice of a particular host and selectable marker is well within the level of ordinary skill in the art.

Protocols for the transformation of yeast are also well known to those of ordinary skill in the art. For example, transformation may be readily accomplished either by preparation of spheroplasts of yeast with DNA (*see* Hinnen et al., *PNAS USA* 75:1929, 1978) or by treatment with alkaline salts such as LiCl (*see* Itoh et al., *J. Bacteriology* 153:163, 1983). Transformation of fungi may also be carried out using polyethylene glycol as described by Cullen et al. (*Bio/Technology* 5:369, 1987).

Viral vectors include those that comprise a promoter that directs the expression of an isolated nucleic acid molecule encoding a desired protein as described above. A wide variety of promoters may be used within the context of the present invention, including for example, promoters such as MoMLV LTR; RSV LTR (*e.g.*, Gorman et al., *Proc. Natl. Acad. Sci. USA* 79:6777, 1982); Friend MuLV LTR; adenoviral promoter (Ohno et al., *Science* 265:781-784, 1994); neomycin phosphotransferase promoter/enhancer; late parvovirus promoter (Koering et al., *Hum. Gene Therap.* 5:457-463, 1994); Herpes TK promoter (*see, e.g.*, McKnight, *Cell* 31:355, 1982); SV40 promoter (*e.g.*, SV40 early promoter (Benoist et al., *Nature* 290:304, 1981); metallothionein IIa gene enhancer/promoter; mouse metallothionein I gene (Hamer et al., *J. Molec. Appl. Genet.* 1:273, 1982); mouse mammary tumor virus promoter (*see, generally*, Etcheverry, "Expression of Engineered Proteins in Mammalian Cell Culture," in *Protein Engineering: Principles and Practice*, Cleland et al. (eds.), pages 163-181 (John Wiley & Sons, Inc. 1996)); cytomegalovirus immediate early promoter, and the cytomegalovirus immediate late promoter. Alternatively, a prokaryotic promoter, such as the bacteriophage T3 RNA polymerase promoter, can be used to control TGF-beta binding-protein gene expression in mammalian cells if the prokaryotic promoter is regulated by a eukaryotic promoter (Zhou et al., *Mol. Cell. Biol.* 10:4529, 1990; Kaufman et al., *Nucl. Acids Res.* 19:4485, 1991).

Within particularly preferred embodiments of the invention, the promoter is a tissue-specific promoter (*see e.g.*, WO 91/02805; EP 0,415,731; and WO 90/07936). Representative examples of suitable tissue specific promoters include neural specific enolase

promoter, platelet derived growth factor beta promoter, bone morphogenic protein promoter, human alpha1-chimaerin promoter, synapsin I promoter and synapsin II promoter. In addition to the above-noted promoters, other viral-specific promoters (e.g., retroviral promoters (including those noted above, as well as others such as HIV promoters), hepatitis, herpes (e.g., EBV), and bacterial, fungal or parasitic (e.g., malarial) -specific promoters may be utilized in order to target a specific cell or tissue that is infected with a virus, bacteria, fungus, or parasite.

Mammalian cells suitable for carrying out the present invention include, among others COS, CHO, SaOS, osteosarcomas, KS483, MG-63, primary osteoblasts, and human or mammalian bone marrow stroma. Additional mammalian host cells include African green monkey kidney cells (Vero; ATCC CRL 1587), human embryonic kidney cells (293-HEK; ATCC CRL 1573), baby hamster kidney cells (BHK-21; ATCC CRL 8544), canine kidney cells (MDCK; ATCC CCL 34), Chinese hamster ovary cells (CHO-K1; ATCC CCL61), rat pituitary cells (GH1; ATCC CCL82), HeLa S3 cells (ATCC CCL2.2), rat hepatoma cells (H-4-II-E; ATCC CRL 1548) SV40-transformed monkey kidney cells (COS-1; ATCC CRL 1650) and murine embryonic cells (NIH-3T3; ATCC CRL 1658).

Mammalian expression vectors for use in carrying out the present invention will include a promoter capable of directing the transcription of a cloned gene or cDNA. Preferred promoters include viral promoters and cellular promoters. Bone specific promoters include the promoter for bone sialo-protein and the promoter for osteocalcin. Viral promoters include the cytomegalovirus immediate early promoter (Boshart et al., *Cell* 41:521-530, 1985), cytomegalovirus immediate late promoter, SV40 promoter (Subramani et al., *Mol. Cell. Biol.* 1:854-864, 1981), MMTV LTR, RSV LTR, metallothionein-1, adenovirus E1a. Cellular promoters include the mouse metallothionein-1 promoter (Palmiter et al., U.S. Patent No. 4,579,821), a mouse V<sub>K</sub> promoter (Bergman et al., *Proc. Natl. Acad. Sci. USA* 81:7041-7045, 1983; Grant et al., *Nucleic Acids Res.* 15:5496, 1987) and a mouse V<sub>H</sub> promoter (Loh et al., *Cell* 33:85-93, 1983). The choice of promoter will depend, at least in part, upon the level of expression desired or the recipient cell line to be transfected.

Such expression vectors may also contain a set of RNA splice sites located downstream from the promoter and upstream from the DNA sequence encoding the peptide or protein of interest. Preferred RNA splice sites may be obtained from adenovirus and/or immunoglobulin genes. Also contained in the expression vectors is a polyadenylation signal located downstream of the coding sequence of interest. Suitable polyadenylation signals include the early or late polyadenylation signals from SV40 (Kaufman and Sharp, *ibid.*), the

polyadenylation signal from the Adenovirus 5 E1B region and the human growth hormone gene terminator (DeNoto et al., *Nucleic Acids Res.* 9:3719-3730, 1981). The expression vectors may include a noncoding viral leader sequence, such as the Adenovirus 2 tripartite leader, located between the promoter and the RNA splice sites. Preferred vectors may also include enhancer sequences, such as the SV40 enhancer. Expression vectors may also include sequences encoding the adenovirus VA RNAs. Suitable expression vectors can be obtained from commercial sources (e.g., Stratagene, La Jolla, California).

Vector constructs comprising cloned DNA sequences can be introduced into cultured mammalian cells by, for example, liposome-mediated transfection, calcium phosphate-mediated transfection (Wigler et al., *Cell* 14:725, 1978; Corsaro and Pearson, *Somatic Cell Genetics* 7:603, 1981; Graham and Van der Eb, *Virology* 52:456, 1973), electroporation (Neumann et al., *EMBO J.* 1:841-845, 1982), or DEAE-dextran mediated transfection (Ausubel et al. (eds.), *Current Protocols in Molecular Biology*, John Wiley and Sons, Inc., NY, 1987); retroviral, adenoviral and protoplast fusion-mediated transfection (see Sambrook et al., *supra*).

Naked vector constructs can also be taken up by muscular cells or other suitable cells subsequent to injection into the muscle of a mammal (or other animal). To identify cells that have been stably transfected with the vector containing the cloned DNA, a selectable marker is generally introduced into the cells along with the gene or cDNA of interest. Preferred selectable markers for use in cultured mammalian cells include genes that confer resistance to drugs, such as neomycin, hygromycin, and methotrexate. The selectable marker may be an amplifiable selectable marker. Preferred amplifiable selectable markers are the DHFR gene and the neomycin resistance gene. Selectable markers are reviewed by Thilly (*Mammalian Cell Technology*, Butterworth Publishers, Stoneham, Massachusetts).

Mammalian cells containing a suitable vector are allowed to grow for a period of time, typically 1-2 days, to begin expressing the DNA sequence(s) of interest. Drug selection is then applied to select for growth of cells that are expressing the selectable marker in a stable fashion. For cells that have been transfected with an amplifiable, selectable marker, the drug concentration may be increased in a stepwise manner to select for increased copy number of the cloned sequences, thereby increasing expression levels. Cells expressing the introduced sequences are selected and screened for production of the protein of interest in the desired form or at the desired level. Cells that satisfy these criteria can then be cloned and scaled up for production.

Numerous insect host cells known in the art can also be useful within the present

invention. For example, baculoviruses as vectors for expressing heterologous DNA sequences in insect cells may be used (Atkinson et al., *Pestic. Sci.* 28:215-224 (1990)). The baculovirus system provides an efficient means to introduce cloned TGF-beta binding-protein encoding polynucleotides into insect cells. Suitable expression vectors are based upon the *Autographa californica* multiple nuclear polyhedrosis virus (AcMNPV), which contain well known promoters such as *Drosophila* heat shock protein (hsp) 70 promoter, *Autographa californica* nuclear polyhedrosis virus immediate-early gene promoter (*ie-1*), and the delayed early 39K promoter, baculovirus p10 promoter, and the *Drosophila* metallothionein promoter. Suitable insect host cells include cell lines derived from IPLB-Sf-21, a *Spodoptera frugiperda* pupal ovarian cell line, such as Sf9 (ATCC CRL 1711), Sf21AE, and Sf21 (Invitrogen Corporation; San Diego, CA), as well as *Drosophila* Schneider-2 cells. Established techniques for producing recombinant proteins in baculovirus systems are provided by Bailey et al., "Manipulation of Baculovirus Vectors," in *Methods in Molecular Biology, Volume 7: Gene Transfer and Expression Protocols*, Murray (ed.), pages 147-168 (The Humana Press, Inc. 1991); Patel et al., "The baculovirus expression system," in *DNA Cloning 2: Expression Systems, 2nd Edition*, Glover et al. (eds.), pages 205-244 (Oxford University Press 1995); Ausubel (1995) at pages 16-37 to 16-57, by Richardson (ed.), *Baculovirus Expression Protocols* (The Humana Press, Inc. 1995); and Lucknow, "Insect Cell Expression Technology," in *Protein Engineering: Principles and Practice*, Cleland et al. (eds.), pages 183-218 (John Wiley & Sons, Inc. 1996).

Alternatively, numerous plant host cells are known in the art and can also be useful for expressing polynucleotides, for example, *Agrobacterium rhizogenes* (Sinkar et al. *J. Biosci. (Bangalore)* 11:47-58, 1987). Expression vectors can also be introduced into plant protoplasts, intact plant tissues, or isolated plant cells. General methods of culturing plant tissues are provided, for example, by Miki et al., "Procedures for Introducing Foreign DNA into Plants," in *Methods in Plant Molecular Biology and Biotechnology*, Glick et al. (eds.), pages 67-88 (CRC Press, 1993).

Within related embodiments of the present invention, proteins of the present invention may be expressed in a transgenic animal whose germ cells and somatic cells contain a gene that encodes the desired protein and that is operatively linked to a promoter effective for the expression of the gene. Alternatively, in a similar manner, transgenic animals may be prepared that lack the desired gene (e.g., "knock-out" mice). Such transgenics may be prepared in a variety of non-human animals, including mice, rats, rabbits, sheep, dogs, goats, and pigs (see Hammer et al., *Nature* 315:680-683, 1985, Palmiter et al., *Science* 222:809-814, 1983,

Brinster et al., *Proc. Natl. Acad. Sci. USA* 82:4438-4442, 1985, Palmiter and Brinster, *Cell* 41:343-345, 1985, and U.S. Patent Nos. 5,175,383, 5,087,571, 4,736,866, 5,387,742, 5,347,075, 5,221,778, and 5,175,384). Briefly, an expression vector, including a nucleic acid molecule to be expressed together with appropriately positioned expression control sequences, is introduced into pronuclei of fertilized eggs, for example, by microinjection. Integration of the injected DNA is detected by blot analysis of DNA from tissue samples. It is preferred that the introduced DNA be incorporated into the germ line of the animal so that it is passed on to the animal's progeny. Tissue-specific expression may be achieved through the use of a tissue-specific promoter, or through the use of an inducible promoter, such as the metallothionein gene promoter (Palmiter et al., 1983, *supra*), which allows regulated expression of the transgene.

Proteins can be isolated by, among other methods, culturing suitable host and vector systems to produce the recombinant translation products of the present invention. Supernatants from such cell lines, or protein inclusion bodies, or whole cells from which the protein is not secreted into the supernatant, can then be treated by a variety of purification procedures in order to isolate the desired proteins. For example, the supernatant may be first concentrated using commercially available protein concentration filters, such as an Amicon or Millipore Pellicon ultrafiltration unit. Following concentration, the concentrate may be applied to a suitable purification matrix such as an affinity matrix, for example, a specific antibody bound to a suitable support. Alternatively, anion or cation exchange resins or size-exclusion matrices may be employed in order to purify the protein. As a further alternative, one or more reverse-phase high performance liquid chromatography (RP-HPLC) steps may be used to further purify the protein. Other methods of isolating the proteins of the present invention are well known in the art. The purity of an isolated protein or polypeptide may be determined by SDS-PAGE analysis followed by Coomassie blue staining or by silver staining.

General methods for expressing and recovering foreign protein produced by a mammalian cell system is provided by, for example, Etcheverry, "Expression of Engineered Proteins in Mammalian Cell Culture," in *Protein Engineering: Principles and Practice*, Cleland et al. (eds.), pages 163 (Wiley-Liss, Inc. 1996). Standard techniques for recovering protein produced by a bacterial system is provided by, for example, Grishammer et al., "Purification of over-produced proteins from *E. coli* cells," in *DNA Cloning 2: Expression Systems, 2nd Edition*, Glover et al. (eds.), pages 59-92 (Oxford University Press 1995). Established methods for isolating recombinant proteins from a baculovirus system are described by Richardson (ed.), *Baculovirus Expression Protocols* (The Humana Press, Inc., 1995).

More generally, a TGF-beta binding-protein can be isolated by standard techniques, such as affinity chromatography, size exclusion chromatography, ion exchange chromatography, HPLC, and the like. Additional variations in TGF-beta binding-protein isolation and purification can be devised by those of skill in the art. For example, anti-TGF-beta binding-protein antibodies, obtained as described herein, can be used to isolate large quantities of protein by immunoaffinity purification. Purification of antibodies of the present invention are further described herein.

#### DETECTABLE LABELS

A detectable label is a molecule or atom that can be conjugated to a polypeptide (including an antibody or fragment thereof) or a polynucleotide to produce a molecule useful for diagnosis. Examples of detectable labels include chelators, photoactive agents, radioisotopes, fluorescent agents, paramagnetic ions, enzymes, and other marker moieties. A TGF binding protein or an antibody that specifically binds to the TGF binding protein or candidate molecules described above may be labeled with a variety of compounds including, for example, fluorescent molecules, toxins, and radionuclides. Representative examples of fluorescent molecules include fluorescein, *Phycobili* proteins, such as phycoerythrin, rhodamine, Texas red, and luciferase. Representative examples of toxins include ricin, abrin diphtheria toxin, cholera toxin, gelonin, pokeweed antiviral protein, tritin, *Shigella* toxin, and *Pseudomonas* exotoxin A. Representative examples of radionuclides include Cu-64, Ga-67, Ga-68, Zr-89, Ru-97, Tc-99m, Rh-105, Pd-109, In-111, I-123, I-125, I-131, Re-186, Re-188, Au-198, Au-199, Pb-203, At-211, Pb-212 and Bi-212. In addition, the antibodies described above may also be labeled or conjugated to one partner of a ligand binding pair. Representative examples include avidin-biotin, streptavidin-biotin, and riboflavin-riboflavin binding protein.

Methods for conjugating or labeling the molecules described herein with the representative labels set forth above may be readily accomplished by one of ordinary skill in the art (see Trichothecene Antibody Conjugate, U.S. Patent No. 4,744,981; Antibody Conjugate, U.S. Patent No. 5,106,951; Fluorogenic Materials and Labeling Techniques, U.S. Patent No. 4,018,884; Metal Radionuclide Labeled Proteins for Diagnosis and Therapy, U.S. Patent No. 4,897,255; and Metal Radionuclide Chelating Compounds for Improved Chelation Kinetics, U.S. Patent No. 4,988,496; see also Inman, *Methods In Enzymology*, Vol. 34, *Affinity Techniques, Enzyme Purification: Part B*, Jakoby and Wilchek (eds.), Academic Press, New York, p. 30, 1974; see also Wilchek and Bayer, "The Avidin-Biotin Complex in Bioanalytical

Applications," *Anal. Biochem.* 171:1-32, 1988). An immunoconjugate is a composition comprising an antibody, such as an anti-TGF-beta binding protein antibody, or an antibody fragment thereof, and a detectable label. Preferably the antibody moiety of the immunoconjugate specifically binds to its cognate antigen with the same or slightly reduced binding affinity after conjugation as before conjugation.

#### PHARMACEUTICAL COMPOSITIONS

As noted above, the present invention also provides a variety of pharmaceutical compositions, comprising one of the above-described molecules that inhibits binding of the TGF-beta binding-protein to a member of the TGF-beta family along with a pharmaceutically or physiologically acceptable carrier, excipient, or diluent. Any suitable carrier known to those of ordinary skill in the art may be employed in the pharmaceutical compositions of the present invention. Carriers for therapeutic use are well known, and are described, for example, in *Remingtons Pharmaceutical Sciences*, Mack Publishing Co. (A.R. Gennaro ed. (1985)). Generally, such carriers should be nontoxic to recipients at the dosages and concentrations employed. Ordinarily, the preparation of such compositions entails combining the therapeutic agent with buffers; antioxidants such as ascorbic acid, low molecular weight (less than about 10 residues) polypeptides; proteins; amino acids; carbohydrates including maltose, glucose, sucrose, or dextrans; chelating agents such as EDTA; glutathione; and other stabilizers and excipients. Neutral buffered saline or saline mixed with nonspecific serum albumin are exemplary diluents. Alternatively, compositions of the present invention may be formulated as a lyophilizate.

In general, the type of carrier is selected based on the mode of administration. Pharmaceutical compositions may be formulated for any appropriate manner of administration, including, for example, topical, oral, nasal, intrathecal, rectal, vaginal, sublingual or parenteral administration, including subcutaneous, intravenous, intramuscular, intrasternal, intracavernous, intrameatal, or intraurethral injection or infusion. A pharmaceutical composition (e.g., for oral administration or delivery by injection) may be in the form of a liquid (e.g., an elixir, syrup, solution, emulsion or suspension). A liquid pharmaceutical composition may include, for example, one or more of the following: sterile diluents such as water for injection, saline solution, preferably physiological saline, Ringer's solution, isotonic sodium chloride, fixed oils that may serve as the solvent or suspending medium, polyethylene glycols, glycerin, propylene glycol or other solvents; antibacterial agents; antioxidants; chelating agents; buffers such as

acetates, citrates or phosphates and agents for the adjustment of tonicity such as sodium chloride or dextrose. A parenteral preparation can be enclosed in ampoules, disposable syringes or multiple dose vials made of glass or plastic. The use of physiological saline is preferred, and an injectable pharmaceutical composition is preferably sterile.

5           The compositions described herein may be formulated for sustained release (*i.e.*, a formulation such as a capsule or sponge that effects a slow release of compound following administration). Such compositions may generally be prepared using well known technology and administered by, for example, oral, rectal or subcutaneous implantation, or by implantation at the desired target site. Sustained-release formulations may contain an agent dispersed in a carrier matrix and/or contained within a reservoir surrounded by a rate controlling membrane. Carriers for use within such formulations are biocompatible, and may also be biodegradable; preferably the formulation provides a relatively constant level of active component release. The amount of active compound contained within a sustained release formulation depends upon the site of implantation, the rate and expected duration of release and the nature of the condition to be treated or prevented. Illustrative carriers useful in this regard include microparticles of poly(lactide-co-glycolide), polyacrylate, latex, starch, cellulose, dextran and the like. Other illustrative delayed-release carriers include supramolecular biovectors, which comprise a non-liquid hydrophilic core (*e.g.*, a cross-linked polysaccharide or oligosaccharide) and, optionally, an external layer comprising an amphiphilic compound, such as a phospholipid (*see e.g.*, U.S. Patent No. 5,151,254 and PCT applications WO 94/20078, WO/94/23701 and WO 96/06638).

          In another illustrative embodiment, biodegradable microspheres (*e.g.*, polylactate polyglycolate) are employed as carriers for the compositions of this invention. Suitable biodegradable microspheres are disclosed, for example, in U.S. Patent Nos. 4,897,268; 5,075,109; 5,928,647; 5,811,128; 5,820,883; 5,853,763; 5,814,344, 5,407,609 and 5,942,252. Modified hepatitis B core protein carrier systems, such as described in WO/99 40934, and references cited therein, will also be useful for many applications. Another illustrative carrier/delivery system employs a carrier comprising particulate-protein complexes, such as those described in U.S. Patent No. 5,928,647, which are capable of inducing a class I-restricted cytotoxic T lymphocyte responses in a host.

30           In another illustrative embodiment, calcium phosphate core particles are employed as carriers or as controlled release matrices for the compositions of this invention. Exemplary calcium phosphate particles are disclosed, for example, in published patent application No. WO/0046147.



For pharmaceutical compositions comprising a polynucleotide encoding an anti-SOST antibody and/or modulating agent (such that the polypeptide and/or modulating agent is generated *in situ*), the polynucleotide may be present within any of a variety of delivery systems known to those of ordinary skill in the art, including nucleic acid, and bacterial, viral and mammalian expression systems. Techniques for incorporating DNA into such expression systems are well known to those of ordinary skill in the art. The DNA may also be "naked," as described, for example, in Ulmer et al., *Science* 259:1745-1749, 1993 and reviewed by Cohen, *Science* 259:1691-1692, 1993. The uptake of naked DNA may be increased by coating the DNA onto biodegradable beads, which are efficiently transported into the cells.

The development of suitable dosing and treatment regimens for using the particular compositions described herein in a variety of treatment regimens, including *e.g.*, oral, parenteral, intravenous, intranasal, and intramuscular administration and formulation, is well known in the art, some of which are briefly discussed below for general purposes of illustration.

In certain applications, the pharmaceutical compositions disclosed herein may be delivered *via* oral administration to an animal. As such, these compositions may be formulated with an inert diluent or with an assimilable edible carrier, or they may be enclosed in hard- or soft-shell gelatin capsule, or they may be compressed into tablets, or they may be incorporated directly with the food of the diet.

In certain circumstances it will be desirable to deliver the pharmaceutical compositions disclosed herein parenterally, intravenously, intramuscularly, or even intraperitoneally. Such approaches are well known to the skilled artisan, some of which are further described, for example, in U.S. Patent No. 5,543,158; U.S. Patent No. 5,641,515 and U.S. Patent No. 5,399,363. In certain embodiments, solutions of the active compounds as free base or pharmacologically acceptable salts may be prepared in water suitably mixed with a surfactant, such as hydroxypropylcellulose. Dispersions may also be prepared in glycerol, liquid polyethylene glycols, and mixtures thereof and in oils. Under ordinary conditions of storage and use, these preparations generally will contain a preservative to prevent the growth of microorganisms.

Illustrative pharmaceutical forms suitable for injectable use include sterile aqueous solutions or dispersions and sterile powders for the extemporaneous preparation of sterile injectable solutions or dispersions (for example, see U.S. Patent No. 5,466,468). In all cases the form must be sterile and must be fluid to the extent that easy syringability exists. It must be stable under the conditions of manufacture and storage and must be preserved against

the contaminating action of microorganisms, such as bacteria and fungi. The carrier can be a solvent or dispersion medium containing, for example, water, ethanol, polyol (e.g., glycerol, propylene glycol, and liquid polyethylene glycol, and the like), suitable mixtures thereof, and/or vegetable oils. Proper fluidity may be maintained, for example, by the use of a coating, such as lecithin, by the maintenance of the required particle size in the case of dispersion and/or by the use of surfactants. The prevention of the action of microorganisms can be facilitated by various antibacterial and antifungal agents, for example, parabens, chlorobutanol, phenol, sorbic acid, thimerosal, and the like. In many cases, it will be preferable to include isotonic agents, for example, sugars or sodium chloride. Prolonged absorption of the injectable compositions can be brought about by the use in the compositions of agents delaying absorption, for example, aluminum monostearate and gelatin.

In one embodiment, for parenteral administration in an aqueous solution, the solution should be suitably buffered if necessary and the liquid diluent first rendered isotonic with sufficient saline or glucose. These particular aqueous solutions are especially suitable for intravenous, intramuscular, subcutaneous and intraperitoneal administration. In this connection, a sterile aqueous medium that can be employed will be known to those of skill in the art in light of the present disclosure. For example, one dosage may be dissolved in 1 ml of isotonic NaCl solution and either added to 1000 ml of hypodermoclysis fluid or injected at the proposed site of infusion, (see for example, *Remington's Pharmaceutical Sciences*, 15th ed., pp. 1035-1038 and 1570-1580). Some variation in dosage will necessarily occur depending on the condition of the subject being treated. Moreover, for human administration, preparations will of course preferably meet sterility, pyrogenicity, and the general safety and purity standards as required by FDA Office of Biologics standards.

In another embodiment of the invention, the compositions disclosed herein may be formulated in a neutral or salt form. Illustrative pharmaceutically-acceptable salts include the acid addition salts (formed with the free amino groups of the protein) and which are formed with inorganic acids such as, for example, hydrochloric or phosphoric acids, or such organic acids as acetic, oxalic, tartaric, mandelic, and the like. Salts formed with the free carboxyl groups can also be derived from inorganic bases such as, for example, sodium, potassium, ammonium, calcium, or ferric hydroxides, and such organic bases as isopropylamine, trimethylamine, histidine, procaine and the like. Upon formulation, solutions will be administered in a manner compatible with the dosage formulation and in such amount as is therapeutically effective.

The carriers can further comprise any and all solvents, dispersion media,

vehicles, coatings, diluents, antibacterial and antifungal agents, isotonic and absorption delaying agents, buffers, carrier solutions, suspensions, colloids, and the like. The use of such media and agents for pharmaceutical active substances is well known in the art. Except insofar as any conventional media or agent is incompatible with the active ingredient, its use in the therapeutic compositions is contemplated. Supplementary active ingredients can also be incorporated into the compositions. The phrase "pharmaceutically-acceptable" refers to molecular entities and compositions that do not produce an allergic or similar untoward reaction when administered to a human.

In certain embodiments, liposomes, nanocapsules, microparticles, lipid particles, vesicles, and the like, are used for the introduction of the compositions of the present invention into suitable host cells/organisms. In particular, the compositions of the present invention may be formulated for delivery either encapsulated in a lipid particle, a liposome, a vesicle, a nanosphere, or a nanoparticle or the like. Alternatively, compositions of the present invention can be bound, either covalently or non-covalently, to the surface of such carrier vehicles.

The formation and use of liposome and liposome-like preparations as potential drug carriers is generally known to those of skill in the art (see for example, Lasic, *Trends Biotechnol.* 16(7):307-21, 1998; Takakura, *Nippon Rinsho* 56(3):691-95, 1998; Chandran et al., *Indian J. Exp. Biol.* 35(8):801-09, 1997; Margalit, *Crit. Rev. Ther. Drug Carrier Syst.* 12(2-3):233-61, 1995; U.S. Patent No. 5,567,434; U.S. Patent No. 5,552,157; U.S. Patent No. 5,565,213; U.S. Patent No. 5,738,868 and U.S. Patent No. 5,795,587, each specifically incorporated herein by reference in its entirety). Liposomes have been used successfully with a number of cell types that are normally difficult to transfect by other procedures, including T cell suspensions, primary hepatocyte cultures and PC 12 cells (Renneisen et al., *J. Biol. Chem.* 265(27):16337-42, 1990; Muller et al., *DNA Cell Biol.* 9(3):221-29, 1990). In addition, liposomes are free of the DNA length constraints that are typical of viral-based delivery systems. Liposomes have been used effectively to introduce genes, various drugs, radiotherapeutic agents, enzymes, viruses, transcription factors, allosteric effectors and the like, into a variety of cultured cell lines and animals. Furthermore, the use of liposomes does not appear to be associated with autoimmune responses or unacceptable toxicity after systemic delivery. In certain embodiments, liposomes are formed from phospholipids that are dispersed in an aqueous medium and spontaneously form multilamellar concentric bilayer vesicles (also termed multilamellar vesicles (MLVs)).

Alternatively, in other embodiments, the invention provides for

pharmaceutically-acceptable nanocapsule formulations of the compositions of the present invention. Nanocapsules can generally entrap compounds in a stable and reproducible way (see, for example, Quintanar-Guerrero et al., *Drug Dev. Ind. Pharm.* 24(12):1113-28, 1998). To avoid side effects due to intracellular polymeric overloading, such ultrafine particles (sized  
5 around 0.1  $\mu\text{m}$ ) may be designed using polymers able to be degraded *in vivo*. Such particles can be made as described, for example, by Couvreur et al., *Crit. Rev. Ther. Drug Carrier Syst.* 5(1):1-20, 1988; zur Muhlen et al., *Eur. J. Pharm. Biopharm.* 45(2):149-55, 1998; Zambaux et al., *J. Controlled Release* 50(1-3):31-40, 1998; and U.S. Patent No. 5,145,684.

In addition, pharmaceutical compositions of the present invention may be placed  
10 within containers, along with packaging material that provides instructions regarding the use of such pharmaceutical compositions. Generally, such instructions will include a tangible expression describing the reagent concentration, as well as within certain embodiments, relative amounts of excipient ingredients or diluents (e.g., water, saline or PBS) that may be necessary to reconstitute the pharmaceutical composition.

#### METHODS OF TREATMENT

The present invention also provides methods for increasing the mineral content and mineral density of bone. Briefly, numerous conditions result in the loss of bone mineral content, including for example, disease, genetic predisposition, accidents that result in the lack  
20 of use of bone (e.g., due to fracture), therapeutics that effect bone resorption or that kill bone forming cells, and normal aging. Through use of the molecules described herein that inhibit binding of the TGF-beta binding-protein to a TGF-beta family member, such conditions may be treated or prevented. As used herein, that bone mineral content is understood to have increased if bone mineral content has increased in a statistically significant manner at a selected site.

A wide variety of conditions that result in loss of bone mineral content may be  
25 treated with the molecules described herein. Patients with such conditions may be identified through clinical diagnosis using well known techniques (see, e.g., Harrison's Principles of Internal Medicine, McGraw-Hill, Inc.). Representative examples of diseases that may be treated included dysplasias, wherein growth or development of bone is abnormal. Representative  
30 examples of such conditions include achondroplasia, cleidocranial dysostosis, enchondromatosis, fibrous dysplasia, Gaucher's Disease, hypophosphatemic rickets, Marfan's syndrome, multiple hereditary exostoses, neurofibromatosis, osteogenesis imperfecta, osteopetrosis, osteopoikilosis, sclerotic lesions, fractures, periodontal disease, pseudoarthrosis,

and pyogenic osteomyelitis.

Other conditions that may be treated or prevented include a wide variety of causes of osteopenia (*i.e.*, a condition that causes greater than one standard deviation of bone mineral content or density below peak skeletal mineral content at youth). Representative  
5 examples of such conditions include anemic states, conditions caused by steroids, conditions caused by heparin, bone marrow disorders, scurvy, malnutrition, calcium deficiency, idiopathic osteoporosis, congenital osteopenia or osteoporosis, alcoholism, chronic liver disease, senility, postmenopausal state, oligomenorrhea, amenorrhea, pregnancy, diabetes mellitus, hyperthyroidism, Cushing's disease, acromegaly, hypogonadism, immobilization or disuse,  
10 reflex sympathetic dystrophy syndrome, transient regional osteoporosis, and osteomalacia.

Within one embodiment of the present invention, bone mineral content or density may be increased by administering to a warm-blooded animal a therapeutically effective amount of a molecule that inhibits a SOST polypeptide from binding to a TGF-beta family member. Examples of warm-blooded animals that may be treated include both vertebrates and mammals,  
15 including for example humans, horses, cows, pigs, sheep, dogs, cats, rats, and mice. Representative examples of therapeutic molecules include ribozymes, ribozyme genes, antisense oligonucleotides, and antibodies as described herein.

Within other embodiments of the present invention, methods are provided for increasing bone density comprising the steps of introducing into cells that home to bone, a  
20 vector that directs the expression of a molecule that inhibits binding of SOST to a member of the TGF-beta family, and administering the vector containing cells to a warm-blooded animal. Briefly, cells that home to bone may be obtained directly from the bone of patients (*e.g.*, cells obtained from the bone marrow such as CD34+, osteoblasts, osteocytes, and the like), from peripheral blood, or from cultures. Representative examples of suitable vectors include viral  
25 vectors such as herpes viral vectors (*e.g.*, U.S. Patent No. 5,288,641); adenoviral vectors (*e.g.*, WO 94/26914, WO 93/9191; Kolls et al., *Proc. Natl. Acad. Sci. USA* 91(1):215-219, 1994; Kass-Eisler et al., *Proc. Natl. Acad. Sci. USA* 90(24):11498-502, 1993; Guzman et al., *Circulation* 88(6):2838-48, 1993; Guzman et al., *Cir. Res.* 73(6):1202-1207, 1993; Zabner et al., *Cell* 75(2):207-216, 1993; Li et al., *Hum. Gene Ther.* 4(4):403-409, 1993; Caillaud et al., *Eur. J.*  
30 *Neurosci.* 5(10):1287-1291, 1993; Vincent et al., *Nat. Genet.* 5(2):130-134, 1993; Jaffe et al., *Nat. Genet.* 1(5):372-378, 1992; and Levrero et al., *Gene* 101(2):195-202, 1991); adeno-associated viral vectors (WO 95/13365; Flotte et al., *PNAS* 90(22):10613-10617, 1993); baculovirus vectors; parvovirus vectors (Koering et al., *Hum. Gene Therap.* 5:457-463, 1994);

pox virus vectors (Panicali and Paoletti, *Proc. Natl. Acad. Sci. USA* 79:4927-4931, 1982; and Ozaki et al., *Biochem. Biophys. Res. Comm.* 193(2):653-660, 1993), and retroviruses (e.g., EP 0,415,731; WO 90/07936; WO 91/0285, WO 94/03622; WO 93/25698; WO 93/25234; U.S. Patent No. 5,219,740; WO 93/11230; WO 93/10218). Viral vectors may likewise be  
5 constructed that contain a mixture of different elements (e.g., promoters, envelope sequences, and the like) from different viruses or from non-viral sources. Within various embodiments, either the viral vector itself, or a viral particle that contains the viral vector may be used in the methods and compositions described below.

Within other embodiments of the invention, nucleic acid molecules that encode a  
10 molecule that inhibits binding of a SOST polypeptide to a member of the TGF-beta family themselves may be administered by a variety of techniques including, for example, administration of asialosomucoid (ASOR) conjugated with poly-L-lysine DNA complexes (Cristano et al., *PNAS* 92:122-92126, 1993); DNA linked to killed adenovirus (Curiel et al., *Hum. Gene Ther.* 3(2):147-154, 1992); cytofectin-mediated introduction (DMRIE-DOPE, Vical,  
15 California); direct DNA injection (Acsadi et al., *Nature* 352:815-818, 1991); DNA ligand (Wu et al., *J. of Biol. Chem.* 264:16985-16987, 1989); lipofection (Felgner et al., *Proc. Natl. Acad. Sci. USA* 84:7413-7417, 1989); liposomes (Pickering et al., *Circ.* 89(1):13-21, 1994; and Wang et al., *PNAS* 84:7851-7855, 1987); microprojectile bombardment (Williams et al., *Proc. Natl. Acad. Sci. USA* 88:2726-2730, 1991); and direct delivery of nucleic acids which encode the  
20 protein itself either alone (Vile and Hart, *Cancer Res.* 53: 3860-3864, 1993), or utilizing PEG-nucleic acid complexes.

Determination of increased bone mineral content may be determined directly through the use of X-rays (e.g., Dual Energy X-ray Absorptometry or "DEXA"), or by inference through bone turnover markers such as osteoblast specific alkaline phosphatase, osteocalcin,  
25 type 1 procollagen C' propeptide (PICP), and total alkaline phosphatase (see Comier, *Curr. Opin. in Rheu.* 7:243 (1995)), or by markers of bone resorption including, but not limited to, pyridinoline, deoxypryridinoline, N-telopeptide, urinary hydroxyproline, plasma tartrate-resistant acid phosphatases, and galactosyl hydroxylysine (see Comier, *id.*). The amount of bone mass may also be calculated from body weights or by using other methods (see Guinness-Hey,  
30 *Metab. Bone Dis. Relat. Res.* 5:177-181, 1984).

As will be evident to one of skill in the art, the amount and frequency of administration will depend, of course, on such factors as the nature and severity of the indication being treated, the desired response, the condition of the patient, and so forth. Typically, the

compositions may be administered by a variety of techniques, as noted above.

The following examples are offered by way of illustration, and not by way of limitation.

5

## EXAMPLES

### EXAMPLE 1

#### MODELING OF SCLEROSTIN CORE REGION

Homology recognition techniques (*e.g.*, PSI-BLAST (Altschul et al., *Nucleic*  
10 *Acids Res.* 25:3389-402 (1997)), FUGUE (Shi et al., *J. Mol. Biol.* 310:243-57 (2001)) suggested that the core region of SOST (SOST\_Core) adopts a cystine-knot fold. FUGUE is a sensitive method for detecting homology between sequences and structures. Human Chorionic Gonadotropin  $\beta$  (hCG- $\beta$ ), for which an experimentally determined 3D structure is known, was identified by FUGUE (Shi et al., *supra*) as the closest homologue of SOST\_Core. Therefore,  
15 hCG- $\beta$  was used as the structural template to build 3D models for SOST\_Core.

An alignment of SOST\_Core and its close homologues is shown in Figure 1. Among the homologues shown in the alignment, only hCG- $\beta$  (CGHB) had known 3D structure. The sequence identity between SOST\_Core and hCG- $\beta$  was approximately 25%. Eight CYS residues were conserved throughout the family, emphasizing the overall structural similarity  
20 between SOST\_Core and hCG- $\beta$ . Three pairs of cystines (1-5, 3-7, 4-8) formed disulfide bonds (shown with solid lines in Figure 1) in a "knot" configuration, which is characteristic to the cystine-knot fold. An extra disulfide bond (2-6), shown as a dotted line in Figure 1, was unique to this family and distinguished the family of proteins from other cystine-knot families (*e.g.*, TGF- $\beta$ , BMP).

25 SOST\_Core was modeled using PDB (Berman et al., *Acta Crystallogr. D. Biol. Crystallogr.* 58(Pt 6 Pt1):899-907 (2002)) entry 1HCN, the 3D structure of hCG- $\beta$  (Wu et al., *Structure* 2:545-58 (1994)), as the structural template. Models were calculated with MODELER (Sali & Blundell, *J. Mol. Biol.* 234:779-815 (1993)). A snapshot of the best model is shown in Figure 2.

30 Most of the cystine-knot proteins form dimers because of the lack of hydrophobic core in a monomer (Scheufler et al., *supra*; Schlunegger and Grutter, *J. Mol. Biol.* 231:445-58 (1993)); Wu et al., *supra*). SOST likely follows the same rule and forms a homodimer to increase its stability. Constructing a model for the dimerized SOST\_Core region presented

several challenges because (1) the sequence similarity between SOST\_Core and hCG- $\beta$  was low (25%); (2) instead of a homodimer, hCG- $\beta$  formed a heterodimer with hCG- $\alpha$ ; and (3) a number of different relative conformations of monomers have been observed in dimerized cystine-knot proteins from different families (*e.g.*, PDGF, TGF- $\beta$ , Neurotrophin, IL-17F, Gonadotropin), which suggested that the dimer conformation of SOST could deviate significantly from the hCG- $\alpha/\beta$  heterodimer conformation. In constructing the model, hCG- $\alpha$  was replaced with hCG- $\beta$  from the heterodimer structure (1HCN) using structure superimposition techniques combined with manual adjustment, and then a SOST\_Core homodimer model was built according to the pseudo hCG- $\beta$  homodimer structure. The final model is shown in Figure 3.

## EXAMPLE 2

### MODELING SOST-BMP INTERACTION

This example describes protein modeling of type I and type II receptor binding sites on BMP that are involved with interaction between BMP and SOST.

Competition studies demonstrated that SOST competed with both type I and type II receptors for binding to BMP. In an ELISA-based competition assay, BMP-6 selectively interacted with the sclerostin-coated surface (300 ng/well) with high affinity ( $K_D = 3.4$  nM). Increasing amounts of BMP receptor IA (FC fusion construct) competed with sclerostin for binding to BMP-6 (11 nM) ( $IC_{50} = 114$  nM). A 10-fold molar excess of the BMP receptor was sufficient to reduce binding of sclerostin to BMP-6 by approximately 50%. This competition was also observed with a BMP receptor II-FC fusion protein ( $IC_{50} = 36$  nM) and DAN ( $IC_{50} = 43$  nM). Specificity of the assay was shown by lack of competition for binding to BMP-6 between sclerostin and a rActivin R1B-FC fusion protein, a TGF- $\beta$  receptor family member that did not bind BMP.

The type I and type II receptor binding sites on a BMP polypeptide have been mapped and were spatially separated (Scheufler et al., *supra*; Innis et al., *supra*; Nickel et al., *supra*; Hart et al. *supra*). Noggin, another BMP antagonist that binds to BMP with high affinity, contacts BMP at both type I and type II receptor binding sites via the N-terminal portion of Noggin (Groppe et al., *supra*). The two  $\beta$ -strands in the core region near the C-terminal also contact BMP at the type II receptor binding site.

A manually tuned alignment of Noggin and SOST indicated that the two polypeptides shared sequence similarity between the N-terminal portions of the proteins and between the core regions. An amino acid sequence alignment is presented in Figure 4. The



cysteine residues that form the characteristic cys-knot were conserved between Noggin and SOST. The overall sequence identity was 24%, and the sequence identity within the N-terminal binding region (alignment positions 1-45) was 33%. Two residues in the Noggin N-terminal binding region, namely Leu (L) at alignment position 21 and Glu (E) at position 23, were reported to play important roles in BMP binding (Groppe et al., *supra*). Both residues were conserved in SOST as well. The sequence similarity within the core region (alignment positions 131-228) was about 20%, but the cys-knot scaffold was maintained and a sufficient number of key residues was conserved, supporting the homology between Noggin and SOST.

The Noggin structure was compared to SOST also to understand how two SOST monomers dimerize. As shown in Figure 5, the Noggin structure suggested that the linker between the N-terminal region and the core region not only played a role in connecting the two regions, but also formed part of the dimerization interface between two Noggin monomers. One major difference between Noggin and SOST was that the linker between the N-terminal region and the core region was much shorter in SOST.

The C-terminal region of SOST may play a role in SOST dimerization. The sequence of Noggin ended with the core region, while SOST had an extra C-terminal region. In the Noggin structure a disulfide bond connected the C-termini of two Noggin monomers. Thus, the C-terminal region of SOST started close to the interface of two monomers and could contribute to dimerization. In addition, secondary structure prediction shows that some portions of the C-terminal region of SOST had a tendency to form helices. This region in SOST may be responsible for the dimerization activity, possibly through helix-helix packing, which mimicked the function of the longer linker in Noggin. Another difference between the structure of Noggin and SOST was the amino acid insertion in the SOST core region at alignment positions 169-185 (see Figure 4). This insertion extended a  $\beta$ -hairpin, which pointed towards the dimerization interface in the Noggin structure (shown in Figure 5 as a loop region in the middle of the monomers and above the C-terminal Cys residue). This elongated  $\beta$ -hairpin could also contribute to SOST dimerization.

### EXAMPLE 3

#### DESIGN AND PREPARATION OF SOST PEPTIDE IMMUNOGENS

This Example describes the design of SOST peptide immunogens that are used for immunizing animals and generating antibodies that block interactions between BMP and SOST and prevent dimer formation of SOST monomers.

BMP Binding Fragments

The overall similarity between SOST and Noggin and the similarity between the N-terminal regions of the two polypeptides suggest that SOST may interact with BMP in a similar manner to Noggin. That is, the N-terminal region of SOST may interact with both the type I and type II receptor binding sites on BMP, and a portion of the core region (amino acid alignment positions 190-220 in Figure 4) may interact with the type II receptor binding site such that antibodies specific for these SOST regions may block or impair binding of BMP to SOST.

The amino acid sequences of these SOST polypeptide fragments for rat and human SOST are provided as follows.

10 SOST\_N\_Linker: The N-terminal region (includes the short linker that connects to the core region)

Human: QGWQAFKNDATEIIPELGEYPEPPPELENNKTMNRAE  
NGGRPPHHPFETKDVSEYS (SEQ ID NO:47)

15 Rat: QGWQAFKNDATEIIPGLREYPEPPQELENNQTMNRAEN  
GGRPPHHPYDTKDVSEYS (SEQ ID NO:48)

SOST\_Core\_Bind: Portion of the core region that is likely to contact BMP at its type II receptor binding site (extended slightly at both termini to include the CYS residue anchors):

20 Human: CIPDRYRAQRVQLLCPGGEAPRARKVRLVASC (SEQ ID NO:49)

Rat: CIPDRYRAQRVQLLCPGGAAPRSRKVRLVASC (SEQ ID NO:50)

SOST Dimerization Fragments

25 The C-terminal region of SOST is likely to be involved in the formation of SOST homodimers (see Example 2). The elongated  $\beta$ -hairpin may also play a role in homodimer formation. Antibodies that specifically bind to such regions may prevent or impair dimerization of SOST monomers, which may in turn interfere with interaction between SOST and BMP. Polypeptide fragments in rat and human SOST corresponding to these regions are as follows.

SOST\_C: the C-terminal region

30 Human: LTRFHNQSELKDFGTEAARPQKGRKPRPRARSAKANQA  
ELENAY (SEQ ID NO:51)

Rat: LTRFHNQSELKDFGPETARPQKGRKPRPRARGAKANQAE

## LENAY (SEQ ID NO:52)

SOST\_Core\_Dimer: Portion of the core region that is likely involved in SOST dimerization (extended slightly at both termini to include the Cys residue anchors):

Human: CGPARLLPNAIGRGKWWRPSPGPDFRC (SEQ ID NO:53)

5 Rat: CGPARLLPNAIGRVKWWRPNGPDFRC (SEQ ID NO:54)

BMP Binding Fragment at SOST N-terminus

The key N-terminal binding region of SOST (alignment positions 1-35 in Figure 4) was modeled on the basis of the Noggin/BMP-7 complex structure (Protein Data Bank Entry No: 1M4U) and the amino acid sequence alignment (see Figure 4) to identify amino acid residues of the SOST N-terminus that likely interact with BMP. The model of SOST is presented in Figure 6. In the comparative model, phenylalanine (Phe, F) at alignment position 8 (see arrow and accompanying text) in the SOST sequence projects into a hydrophobic pocket on the surface of the BMP dimer. The same "knob-into-hole" feature has been observed in the BMP and type I receptor complex structure (Nickel et al., *supra*), where Phe85 of the receptor fits into the same pocket, which is a key feature in ligand-type I receptor recognition for TGF- $\beta$  superfamily members (including, for example, TGF- $\beta$  family, BMP family, and the like). According to the model, a proline (Pro) directed turn is also conserved, which allows the N-terminal binding fragment to thread along the BMP dimer surface, traveling from type I receptor binding site to type II receptor binding site on the other side of the complex. Also conserved is another Pro-directed turn near the carboxy end of the binding fragment, which then connects to the linker region. Extensive contacts between SOST and BMP are evident in Figure 6.

Peptide Immunogens

25 Peptides were designed to encompass the SOST N-terminal region predicted to make contact with BMP proteins. The peptide sequences are presented below. For immunizing animals, the peptide sequences were designed to overlap, and an additional cysteine was added to the C-terminal end to facilitate crosslinking to KLH. The peptides were then used for immunization. The peptide sequences of the immunogens are as follows.

30 Human SOST:

QGWQAFKNDATETIPELGEY (SEQ ID NO:2)

TEIPELGEYPEPPPELENN (SEQ ID NO:3)

PEPPPELENNKTMNRAENG (SEQ ID NO:4)

KTMNRAENGGRPPHHPFETK (SEQ ID NO:5)

RPPHHPFETKDVSEYS (SEQ ID NO:6)

Human SOST Peptides with Additional Cys:

5 QGWQAFKNDATEIIPELGEY-C (SEQ ID NO:7)

TEIIPELGEYPEPPPELENN-C (SEQ ID NO:8)

PEPPPELENNKTMNRAENG-C (SEQ ID NO:9)

KTMNRAENGGRPPHHPFETK-C (SEQ ID NO:10)

RPPHHPFETKDVSEYS-C (SEQ ID NO:11)

10

Rat SOST:

QGWQAFKNDATEIIPGLREYPEPP (SEQ ID NO:12)

PEPPQELENNQTMNRAENG (SEQ ID NO:13)

15 ENGGRPPHHPYDTKDVSEYS (SEQ ID NO:14)

TEIIPGLREYPEPPQELENN (SEQ ID NO:15)

Rat SOST Peptides with Additional Cys:

QGWQAFKNDATEIIPGLREYPEPP-C (SEQ ID NO:16)

20 PEPPQELENNQTMNRAENG-C (SEQ ID NO:17)

ENGGRPPHHPYDTKDVSEYS-C (SEQ ID NO:18)

TEIIPGLREYPEPPQELENN-C (SEQ ID NO:19)

25 The following peptides were designed to contain the amino acid portion of core region that was predicted to make contact with BMP proteins. Cysteine was added at the C-terminal end of each peptide for conjugation to KLH, and the conjugated peptides were used for immunization. In the Docking Core N-terminal Peptide an internal cysteine was changed to a serine to avoid double conjugation to KLH.

For Human SOST:

30 Amino acid sequence without Cys residues added:

Docking\_Core\_N-terminal\_Peptide: IPDRYRAQRVQLLCPGGEAP (SEQ ID NO:21)

Docking\_Core\_Cterm\_Peptide: QLLCPGGEAPRARKVRLVAS (SEQ ID

NO:22)

Docking\_Core\_N-terminal\_Peptide: IPDRYRAQQRVQLLCPGGEAP-C (SEQ ID NO:23)

5 Docking\_Core\_Cterm\_Peptide: QLLCPGGEAPRARKVRLVAS-C (SEQ ID NO:24)

For Rat SOST:

10 Amino acid sequence without Cys residues added or substituted:

Docking\_Core\_N-terminal\_Peptide: IPDRYRAQQRVQLLCPGG (SEQ ID NO:25)

Docking\_Core\_Cterm\_Peptide: PGGAAPRSRKVRLVAS (SEQ ID NO:26)

15 Peptide immunogens with Cys added and substituted:

Docking\_Core\_N-terminal\_Peptide: IPDRYRAQQRVQLLSPGG-C (SEQ ID NO:27)

Docking\_Core\_Cterm\_Peptide: PGGAAPRSRKVRLVAS-C (SEQ ID NO:28)

20 Two regions within SOST that potentially interact to form SOST homodimers include the amino acids with the SOST core region that are not present in Noggin. Human SOST peptides designed to contain this sequence had a C-terminal or N-terminal Cys that was conjugated to KLH. For the rat SOST peptide, a cysteine was added to the carboxy terminus of the sequence (SEQ ID NO:31). The KLH conjugated peptides were used for immunization.

25 For Human SOST:

CGPARLLPNAIGRGKWWRPS (SEQ ID NO:29)

IGRGKWWRPSGPDFRC (SEQ ID NO:30)

For Rat SOST:

30 PNAIGRVKWWRPNGPDFR (SEQ ID NO:31)

Rat SOST peptide with cysteine added

PNAIGRVKWWRPNGPDFR-C (SEQ ID NO:32)

The second region within SOST that potentially interacts to form SOST homodimers includes the C-terminal region. Peptide immunogens were designed to include amino acid sequences within this region (see below). For conjugation to KLH, a cysteine residue was added to the C-terminal end, and the conjugated peptides were used for immunization.

For Human SOST:

KRLTRFHNQS ELKDFGTEAA (SEQ ID NO:33)

ELKDFGTEAA RPQKGRKPRP (SEQ ID NO:34)

RPQKGRKPRP RARSAKANQA (SEQ ID NO:35)

RARSAKANQA ELENAY (SEQ ID NO:36)

Peptide immunogens with Cys added at C-terminus:

KRLTRFHNQS ELKDFGTEAA-C (SEQ ID NO:37)

ELKDFGTEAA RPQKGRKPRP-C (SEQ ID NO:38)

RPQKGRKPRP RARSAKANQA-C (SEQ ID NO:39)

RARSAKANQA ELENAY-C (SEQ ID NO:40)

For Rat SOST:

KRLTRFHNQSELKDFGPETARPQ (SEQ ID NO:41)

KGRKPRPRARGAKANQAELENAY (SEQ ID NO:42)

SELKDFGPETARPQKGRKPRPRAR (SEQ ID NO:43)

Peptide immunogens with Cys added at C-terminus:

KRLTRFHNQSELKDFGPETARPQ-C (SEQ ID NO:44)

KGRKPRPRARGAKANQAELENAY-C (SEQ ID NO:45)

SELKDFGPETARPQKGRKPRPRAR-C (SEQ ID NO:46)

#### EXAMPLE 4

##### ASSAY FOR DETECTING BINDING OF ANTIBODIES TO A TGF-BETA BINDING-PROTEIN

This example describes an assay for detecting binding of a ligand, for example, an antibody or antibody fragment thereof, to sclerostin.

A FLAG®-sclerostin fusion protein was prepared according to protocols

provided by the manufacturer (Sigma Aldrich, St. Louis, MO) and as described in U.S. Patent No. 6,395,511. Each well of a 96 well microtiter plate is coated with anti-FLAG® monoclonal antibody (Sigma Aldrich) and then blocked with 10% BSA in PBS. The fusion protein (20 ng) is added to 100 µl PBS/0.2% BSA and adsorbed onto the 96-well plate for 60 minutes at room temperature. This protein solution is removed and the wells are washed to remove unbound fusion protein. A BMP, for example, BMP-4, BMP-5, BMP-6, or BMP-7, is diluted in PBS/0.2% BSA and added to each well at concentrations ranging from 10 pM to 500 nM. After an incubation for 2 hours at room temperature, the binding solution is removed and the plate is washed three times with 200 µl volumes of PBS/0.2% BSA. Binding of the BMP to sclerostin is detected using polyclonal antiserum or monoclonal antibody specific for the BMP and an appropriate enzyme-conjugated second step reagent according to standard ELISA techniques (see, e.g., Ausubel et al., *Current Protocols in Mol Biol.* Vol 2 11.2.1-11.2.22 (1998)). Specific binding is calculated by subtracting non-specific binding from total binding and analyzed using the LIGAND program (Munson and Podbard, *Anal. Biochem.* 107:220-39 (1980)).

Binding of sclerostin to a BMP is also detected by homogeneous time resolved fluorescence detection (Mellor et al., *J Biomol. Screening*, 3:91-99 (1998)). A polynucleotide sequence encoding sclerostin is operatively linked to a human immunoglobulin constant region in a recombinant nucleic acid construct and expressed as a human Fc-sclerostin fusion protein according to methods known in the art and described herein. Similarly, a BMP ligand is engineered and expressed as a BMP-mouse Fc fusion protein. These two fusion proteins are incubated together and the assay conducted as described by Mellor et al.

#### EXAMPLE 5

##### SCREENING ASSAY FOR ANTIBODIES THAT INHIBIT BINDING OF TGF-BETA FAMILY MEMBERS TO TGF-BETA BINDING PROTEIN

This example describes a method for detecting an antibody that inhibits binding of a TGF-beta family member to sclerostin. An ELISA is performed essentially as described in Example 4 except that the BMP concentration is held fixed at its K<sub>d</sub> (determined, for example, by BIAcore analysis). In addition, an antibody or a library or collection of antibodies is added to the wells to a concentration of 1 µM. Antibodies are incubated for 2 hours at room temperature with the BMP and sclerostin, the solution removed, and the bound BMP is quantified as described (see Example 4). Antibodies that inhibit 40% of the BMP binding observed in the

absence of antibody are considered antagonists of this interaction. These antibodies are further evaluated as potential inhibitors by performing titration studies to determine their inhibition constants and their effect on TGF-beta binding-protein binding affinity. Comparable specificity control assays may also be conducted to establish the selectivity profile for the identified antagonist using assays dependent on the BMP ligand action (e.g., a BMP/BMP receptor competition study).

### EXAMPLE 6

#### INHIBITION OF TGF-BETA BINDING-PROTEIN LOCALIZATION TO BONE MATRIX

Evaluation of inhibition of localization to bone matrix (hydroxyapatite) is conducted using modifications to the method of Nicolas (*Calcif. Tissue Int.* 57:206-12 (1995)). Briefly, <sup>125</sup>I-labelled TGF-beta binding-protein is prepared as described by Nicolas (*supra*). Hydroxyapatite is added to each well of a 96-well microtiter plate equipped with a polypropylene filtration membrane (Polyfritroninc, Weymouth MA). TGF-beta binding-protein diluted in 0.2% albumin in PBS buffer is then added to the wells. The wells containing matrix are washed 3 times with 0.2% albumin in PBS buffer. Adsorbed TGF-beta binding-protein is eluted using 0.3 M NaOH and then quantified.

An antibody or other agent that inhibits or impairs binding of the sclerostin TGF-beta binding protein to the hydroxyapatite is identified by incubating the TGF-beta binding protein with the antibody and applying the mixture to the matrix as described above. The matrix is washed 3 times with 0.2% albumin in PBS buffer. Adsorbed sclerostin is eluted with 0.3 M NaOH and then quantified. An antibody that inhibits the level of binding of sclerostin to the hydroxyapatite by at least 40% compared to the level of binding observed in the absence of antibody is considered a bone localization inhibitor. Such an antibody is further characterized in dose response studies to determine its inhibition constant and its effect on TGF-beta binding-protein binding affinity.

From the foregoing, although specific embodiments of the invention have been described herein for purposes of illustration, various modifications may be made without deviating from the spirit and scope of the invention. Accordingly, the invention is not limited except as by the appended claims.



## CLAIMS

We claim the following:

1. An isolated antibody, or an antigen-binding fragment thereof, that binds specifically to a SOST polypeptide, said SOST polypeptide comprising an amino acid sequence set forth in SEQ ID NO:1, 20, 58, 60, 62, or 68 wherein the antibody competitively inhibits binding of the SOST polypeptide to at least one of (i) a bone morphogenic protein (BMP) Type I Receptor binding site and (ii) a BMP Type II Receptor binding site, wherein the BMP Type I Receptor binding site is capable of binding to a BMP Type I Receptor polypeptide comprising an amino acid sequence set forth in a sequence selected from the group consisting of GenBank Acc. Nos. NM\_004329 (SEQ ID NO:71); D89675 (SEQ ID NO:72); NM\_001203 (SEQ ID NO:73); S75359 (SEQ ID NO:74); NM\_030849 (SEQ ID NO:75); D38082 (SEQ ID NO:76); NP\_001194 (SEQ ID NO:77); BAA19765 (SEQ ID NO:78); and AAB33865 (SEQ ID NO:79), and wherein the BMP Type II Receptor binding site is capable of binding to a BMP Type II Receptor polypeptide comprising the amino acid sequence set forth in a sequence selected from the group consisting of GenBank Acc. NOs. U25110 (SEQ ID NO:80); NM\_033346 (SEQ ID NO:81); Z48923 (SEQ ID NO:83); CAA88759 (SEQ ID NO:84); and NM\_001204 (SEQ ID NO:82).
2. The antibody, or antigen-binding fragment thereof, according to claim 1 wherein the antibody, or the antigen-binding fragment thereof, binds specifically to a polypeptide that comprises an amino acid sequence selected from the group consisting of SEQ ID NOs:2, 3, 4, 5, 6, 12, 13, 14, 15, 21, 22, 25, 26, 47, 48, 49, and 50.
3. An isolated antibody, or an antigen-binding fragment thereof, that is produced by immunizing a non-human animal with a peptide of at least 20 amino acids and no more than 75 amino acids that comprises an amino acid sequence selected from the group consisting of SEQ ID NOs:2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 21, 22, 23, 24, 25, 26, 27, 28, 47, 48, 49, and 50.
4. The antibody, or an antigen-binding fragment thereof, according to claim 3 wherein the antibody binds specifically to a SOST polypeptide, said SOST polypeptide comprising an amino acid sequence set forth in SEQ ID NOs:1, 20, 58, 60, 62, or 68.
5. The antibody, or an antigen-binding fragment thereof, according to claim 4 wherein the antibody competitively inhibits binding of the SOST polypeptide to at least one of (i) a bone morphogenic protein (BMP) Type I Receptor binding site and (ii) a BMP Type II Receptor

binding site, wherein the BMP Type I Receptor binding site is capable of binding to a BMP Type I Receptor polypeptide comprising an amino acid sequence set forth in a sequence selected from the group consisting of GenBank Acc. Nos. NM\_004329 (SEQ ID NO:71); D89675 (SEQ ID NO:72); NM\_001203 (SEQ ID NO:73); S75359 (SEQ ID NO:74); NM\_030849 (SEQ ID NO:75); D38082 (SEQ ID NO:76); NP\_001194 (SEQ ID NO:77); BAA19765 (SEQ ID NO:78); and AAB33865 (SEQ ID NO:79), and wherein the BMP Type II Receptor binding site is capable of binding to a BMP Type II Receptor polypeptide comprising the amino acid sequence set forth in a sequence selected from the group consisting of GenBank Acc. NOs. U25110 (SEQ ID NO:80); NM\_033346 (SEQ ID NO:81); Z48923 (SEQ ID NO:83); CAA88759 (SEQ ID NO:84); and NM\_001204 (SEQ ID NO:82).

6. An antibody, or an antigen-binding fragment thereof, that is produced by immunizing a non-human animal with a peptide of at least 20 amino acids and no more than 75 amino acids that comprises an amino acid sequence selected from the group consisting of SEQ ID NO:2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 21, 22, 23, 24, 25, 26, 27, 28, 47, 48, 49, and 50 wherein the antibody binds specifically to a SOST polypeptide, said SOST polypeptide comprising an amino acid sequence set forth in SEQ ID NOs:1, 20, 58, 60, 62, or 68, and wherein the antibody competitively inhibits binding of the SOST polypeptide to at least one of (i) a bone morphogenic protein (BMP) Type I Receptor binding site and (ii) a BMP Type II Receptor binding site, wherein the BMP Type I Receptor binding site is capable of binding to a BMP Type I Receptor polypeptide comprising an amino acid sequence set forth in a sequence selected from the group consisting of GenBank Acc. Nos. NM\_004329 (SEQ ID NO:71); D89675 (SEQ ID NO:72); NM\_001203 (SEQ ID NO:73); S75359 (SEQ ID NO:74); NM\_030849 (SEQ ID NO:75); D38082 (SEQ ID NO:76); NP\_001194 (SEQ ID NO:77); BAA19765 (SEQ ID NO:78); and AAB33865 (SEQ ID NO:79), and wherein the BMP Type II Receptor binding site is capable of binding to a BMP Type II Receptor polypeptide comprising the amino acid sequence set forth in a sequence selected from the group consisting of GenBank Acc. NOs. U25110 (SEQ ID NO:80); NM\_033346 (SEQ ID NO:81); Z48923 (SEQ ID NO:83); CAA88759 (SEQ ID NO:84); and NM\_001204 (SEQ ID NO:82).

7. An isolated antibody, or an antigen-binding fragment thereof, that binds specifically to a SOST polypeptide and that impairs formation of a SOST homodimer, wherein the SOST polypeptide comprises an amino acid sequence set forth in SEQ ID NOs:1, 20, 58, 60, 62, or 68.

8. The antibody of claim 7 wherein the antibody binds to a polypeptide that comprises an

amino acid sequence selected from the group consisting of SEQ ID NOs:29, 30, 31, 33, 34, 35, 36, 41, 42, 43, 51, 52, 53, and 54.

9. An antibody, or an antigen-binding fragment thereof, that is produced by immunizing a non-human animal with a peptide of at least 20 amino acids and no more than 75 amino acids comprising an amino acid sequence selected from the group consisting of SEQ ID NOs: 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 51, 52, 53, and 54.

10. The antibody, or an antigen-binding fragment thereof, according to claim 9, wherein the antibody, or the antigen-binding fragment thereof, binds specifically to a SOST polypeptide that comprises an amino acid sequence set forth in SEQ ID NO:1, 20, 58, 60, 62, or 68.

11. The antibody, or an antigen-binding fragment thereof, according to claim 10, wherein the antibody, or the antigen-binding fragment thereof, impairs formation of a SOST homodimer.

12. An antibody, or an antigen-binding fragment thereof, that is produced by immunizing a non-human animal with a peptide of at least 20 amino acids and no more than 75 amino acids comprising an amino acid sequence selected from the group consisting of SEQ ID NOs: 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 51, 52, 53, and 54, wherein the antibody, or the antigen-binding fragment thereof, binds specifically to a SOST polypeptide that comprises an amino acid sequence set forth in SEQ ID NO:1, 20, 58, 60, 62, or 68, and wherein the antibody impairs formation of a SOST homodimer.

13. The antibody of any one of claims 1-12, wherein the antibody is a polyclonal antibody.

14. The antibody of any one of claims 1-12, wherein the antibody is a monoclonal antibody.

15. The antibody of claim 14 wherein the monoclonal antibody is selected from the group consisting of a mouse monoclonal antibody, a human monoclonal antibody, a rat monoclonal antibody, and a hamster monoclonal antibody.

16. A hybridoma cell producing the antibody of claim 14.

17. A host cell that is capable of expressing the antibody of claim 14.

18. The antibody of any one of claims 1-12 wherein the antibody is a humanized antibody or a chimeric antibody.

19. A host cell that is capable of expressing the antibody of claim 18.

20. The antibody of any one of claims 1-12, wherein the antigen-binding fragment is selected from the group consisting of F(ab')<sub>2</sub>, Fab', Fab, Fd, and Fv.

21. The antibody of any one of claims 1-12 that comprises a single chain antibody.

22. A host cell that is capable of expressing the antibody of claim 21.

23. A composition comprising an antibody, or antigen-binding fragment thereof, according to any one of claims 1-12 and a physiologically acceptable carrier.

24. An immunogen comprising a peptide comprising 6, 7, 8, 9, 10, 11, or 12 consecutive amino acids of a SOST polypeptide, said SOST polypeptide comprising an amino acid sequence set forth in SEQ ID NOs:1, 20, 58, 60, 62, or 68, wherein the peptide is capable of eliciting in a non-human animal an antibody that binds specifically to the SOST polypeptide and that competitively inhibits binding of the SOST polypeptide to at least one of (i) a bone morphogenic protein (BMP) Type I Receptor binding site and (ii) a BMP Type II Receptor binding site, wherein the BMP Type I Receptor binding site is capable of binding to a BMP Type I Receptor polypeptide comprising an amino acid sequence set forth in a sequence selected from the group consisting of GenBank Acc. Nos. NM\_004329 (SEQ ID NO:71); D89675 (SEQ ID NO:72); NM\_001203 (SEQ ID NO:73); S75359 (SEQ ID NO:74); NM\_030849 (SEQ ID NO:75); D38082 (SEQ ID NO:76); NP\_001194 (SEQ ID NO:77); BAA19765 (SEQ ID NO:78); and AAB33865 (SEQ ID NO:79), and wherein the BMP Type II Receptor binding site is capable of binding to a BMP Type II Receptor polypeptide comprising the amino acid sequence set forth in a sequence selected from the group consisting of GenBank Acc. NOs. U25110 (SEQ ID NO:80); NM\_033346 (SEQ ID NO:81); Z48923 (SEQ ID NO:83); CAA88759 (SEQ ID NO:84); and NM\_001204 (SEQ ID NO:82).

25. An immunogen comprising a peptide comprising at least 21 consecutive amino acids and no more than 50 consecutive amino acids of a SOST polypeptide, said SOST polypeptide comprising an amino acid sequence set forth in SEQ ID NOs:1, 20, 58, 60, 62, or 68, wherein the peptide is capable of eliciting in a non-human animal an antibody that binds specifically to the SOST polypeptide and that competitively inhibits binding of the SOST polypeptide to at least one of (i) a bone morphogenic protein (BMP) Type I Receptor binding site and (ii) a BMP Type II Receptor binding site, wherein the BMP Type I Receptor binding site is capable of binding to a BMP Type I Receptor polypeptide comprising an amino acid sequence set forth in a sequence selected from the group consisting of GenBank Acc. Nos. NM\_004329 (SEQ ID NO:71); D89675 (SEQ ID NO:72); NM\_001203 (SEQ ID NO:73); S75359 (SEQ ID NO:74); NM\_030849 (SEQ ID NO:75); D38082 (SEQ ID NO:76); NP\_001194 (SEQ ID NO:77); BAA19765 (SEQ ID NO:78); and AAB33865 (SEQ ID NO:79), and wherein the BMP Type II Receptor binding site is capable of binding to a BMP Type II Receptor polypeptide comprising the amino acid sequence set forth in a sequence selected from the group consisting of GenBank Acc. NOs. U25110 (SEQ ID NO:80); NM\_033346 (SEQ ID NO:81); Z48923 (SEQ ID NO:83);

CAA88759 (SEQ ID NO:84); and NM\_001204 (SEQ ID NO:82).

26. An immunogen comprising a peptide of at least 20 amino acids and no more than 75 amino acids comprising an amino acid sequence selected from the group consisting of SEQ ID NOs: 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 21, 22, 23, 24, 25, 26, 27, 28, 47, 48, 49, and 50.

27. The immunogen according to claim 26 wherein the peptide is capable of eliciting in a non-human animal an antibody that binds specifically to a SOST polypeptide, said SOST polypeptide comprising an amino acid sequence set forth in SEQ ID NO:1, 20, 58, 60, 62, or 68.

28. The immunogen according to claim 27 wherein the antibody competitively inhibits binding of the SOST polypeptide to at least one of (i) a bone morphogenic protein (BMP) Type I Receptor binding site and (ii) a BMP Type II Receptor binding site, wherein the BMP Type I Receptor binding site is capable of binding to a BMP Type I Receptor polypeptide comprising an amino acid sequence set forth in a sequence selected from the group consisting of GenBank Acc. Nos. NM\_004329 (SEQ ID NO:71); D89675 (SEQ ID NO:72); NM\_001203 (SEQ ID NO:73); S75359 (SEQ ID NO:74); NM\_030849 (SEQ ID NO:75); D38082 (SEQ ID NO:76); NP\_001194 (SEQ ID NO:77); BAA19765 (SEQ ID NO:78); and AAB33865 (SEQ ID NO:79), and wherein the BMP Type II Receptor binding site is capable of binding to a BMP Type II Receptor polypeptide comprising the amino acid sequence set forth in a sequence selected from the group consisting of GenBank Acc. NOs. U25110 (SEQ ID NO:80); NM\_033346 (SEQ ID NO:81); Z48923 (SEQ ID NO:83); CAA88759 (SEQ ID NO:84); and NM\_001204 (SEQ ID NO:82).

29. An immunogen comprising a peptide of at least 20 amino acids and no more than 75 amino acids comprising an amino acid sequence selected from the group consisting of SEQ ID NOs: 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 21, 22, 23, 24, 25, 26, 27, 28, 47, 48, 49, and 50, wherein the peptide is capable of eliciting in a non-human animal an antibody that binds specifically to a SOST polypeptide, said SOST polypeptide comprising an amino acid sequence set forth in SEQ ID NO:1, 20, 58, 60, 62, or 68, wherein the antibody competitively inhibits binding of the SOST polypeptide to at least one of (i) a bone morphogenic protein (BMP) Type I Receptor binding site and (ii) a BMP Type II Receptor binding site, wherein the BMP Type I Receptor binding site is capable of binding to a BMP Type I Receptor polypeptide comprising an amino acid sequence set forth in a sequence selected from the group consisting of GenBank Acc. Nos. NM\_004329 (SEQ ID NO:71); D89675 (SEQ ID NO:72); NM\_001203 (SEQ ID NO:73); S75359 (SEQ ID NO:74); NM\_030849 (SEQ ID NO:75); D38082 (SEQ ID

NO:76); NP\_001194 (SEQ ID NO:77); BAA19765 (SEQ ID NO:78); and AAB33865 (SEQ ID NO:79), and wherein the BMP Type II Receptor binding site is capable of binding to a BMP Type II Receptor polypeptide comprising the amino acid sequence set forth in a sequence selected from the group consisting of GenBank Acc. NOs. U25110 (SEQ ID NO:80);  
5 NM\_033346 (SEQ ID NO:81); Z48923 (SEQ ID NO:83); CAA88759 (SEQ ID NO:84); and NM\_001204 (SEQ ID NO:82).

30. An immunogen comprising a peptide of at least 20 amino acids and no more than 75 amino acids comprising an amino acid sequence selected from the group consisting of SEQ ID NOs: 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 51, 52, 53, and 54.

10 31. The immunogen according to claim 30 wherein the peptide is capable of eliciting in a non-human animal an antibody that binds specifically to a SOST polypeptide, said SOST polypeptide comprising an amino acid sequence set forth in SEQ ID NO:1, 20, 58, 60, 62, or 68.

32. The immunogen according to claim 31 wherein the antibody impairs formation of a SOST homodimer.

15 33. An immunogen comprising a peptide of at least 20 amino acids and no more than 75 amino acids comprising an amino acid sequence selected from the group consisting of SEQ ID NOs: 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 51, 52, 53, and 54, wherein the peptide is capable of eliciting in a non-human animal an antibody that binds specifically to a SOST polypeptide, said SOST polypeptide comprising an amino acid sequence  
20 set forth in SEQ ID NO:1, 20, 58, 60, 62, or 68, and wherein the antibody impairs formation of a SOST homodimer.

34. An immunogen comprising a peptide comprising 6, 7, 8, 9, 10, 11, or 12 consecutive amino acids of a SOST polypeptide, said SOST polypeptide comprising an amino acid sequence set forth in SEQ ID NO:1, 20, 58, 60, 62, or 68, wherein the peptide is capable of eliciting in a  
25 non-human animal an antibody that binds specifically to the SOST polypeptide and that impairs formation of a SOST homodimer.

35. An immunogen comprising a peptide that comprises at least 21 consecutive amino acids and no more than 50 consecutive amino acids of a SOST polypeptide, said SOST polypeptide comprising an amino acid sequence set forth in SEQ ID NO:1, 20, 58, 60, 62, or 68, wherein the  
30 peptide is capable of eliciting in a non-human animal an antibody that binds specifically to the SOST polypeptide and that impairs formation of a SOST homodimer.

36. The immunogen of any one of claims 24-35 wherein the peptide is associated with a carrier molecule.

37. The immunogen of claim 36 wherein the carrier molecule is a carrier polypeptide.

38. The immunogen of claim 37 wherein the carrier polypeptide is keyhole limpet  
5 hemocyanin.

39. A method for producing an antibody that specifically binds to a SOST polypeptide, comprising immunizing a non-human animal with an immunogen according to any one of claims 24-29, wherein (a) the SOST polypeptide comprises an amino acid sequence set forth in SEQ ID NO:1, 20, 58, 60, 62, or 68; (b) the antibody competitively inhibits binding of the SOST  
10 polypeptide to at least one of (i) a bone morphogenic protein (BMP) Type I Receptor binding site and (ii) a BMP Type II Receptor binding site; (c) the BMP Type I Receptor binding site is capable of binding to a BMP Type I Receptor polypeptide comprising the amino acid sequence set forth in a sequence selected from the group consisting of GenBank Acc. Nos. NM\_004329 (SEQ ID NO:71); D89675 (SEQ ID NO:72); NM\_001203 (SEQ ID NO:73); S75359 (SEQ ID  
15 NO:74); NM\_030849 (SEQ ID NO:75); D38082 (SEQ ID NO:76); NP\_001194 (SEQ ID NO:77); BAA19765 (SEQ ID NO:78); and AAB33865 (SEQ ID NO:79); and (d) the BMP Type II Receptor binding site is capable of binding to a BMP Type II Receptor polypeptide comprising the amino acid sequence set forth in a sequence selected from the group consisting of GenBank Acc. NOs. U25110 (SEQ ID NO:80); NM\_033346 (SEQ ID NO:81); Z48923 (SEQ  
20 ID NO:83); CAA88759 (SEQ ID NO:84); and NM\_001204 (SEQ ID NO:82).

40. A method for producing an antibody that specifically binds to a SOST polypeptide, said SOST polypeptide comprising an amino acid sequence set forth in SEQ ID NO:1, 20, 58, 60, 62, or 68, comprising immunizing a non-human animal with an immunogen according to any one of claims 30-35, wherein the antibody impairs formation of a SOST homodimer.

25 41. A method for identifying an antibody that modulates a TGF-beta signaling pathway comprising:

(a) contacting an antibody that specifically binds to a SOST polypeptide which comprises an amino acid sequence set forth in any one of SEQ ID NOS: 1, 20, 58, 60, 62, and 68 with at least one SOST peptide comprising an amino acid sequence selected from the  
30 group consisting of SEQ ID NO: 2-19, and 21-54, under conditions and for a time sufficient to permit formation of an antibody/SOST peptide complex; and

(b) detecting a level of antibody/SOST peptide complex, and thereby

detecting the presence of an antibody that modulates a TGF-beta signaling pathway.

42. A method for identifying an antibody that impairs binding of a BMP to a SOST polypeptide comprising:

(a) contacting (i) an antibody that specifically binds to a SOST polypeptide which comprises an amino acid sequence set forth in any one of SEQ ID NOS: 1, 20, 58, 60, 62, and 68 with (ii) at least one SOST peptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO: 2-19, 21-28, and 47-50 under conditions and for a time sufficient to permit formation of an antibody/SOST peptide complex; and

(b) detecting a level of antibody/SOST peptide complex, and thereby detecting the presence of an antibody that impairs binding of a BMP to a SOST polypeptide.

43. A method for identifying an antibody that impairs SOST homodimer formation comprising:

(a) contacting (i) an antibody that specifically binds to a SOST polypeptide which comprises an amino acid sequence set forth in any one of SEQ ID NOS: 1, 20, 58, 60, 62, and 68, with (ii) at least one SOST peptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO: 29-46 and 51-54 under conditions and for a time sufficient to permit formation of an antibody/SOST peptide complex; and

(b) detecting a level of antibody/SOST peptide complex, and thereby detecting the presence of an antibody that that impairs SOST homodimer formation.

44. A method for identifying an antibody that increases bone mineral content comprising:

(a) contacting (i) an antibody that specifically binds to a SOST polypeptide which comprises an amino acid sequence set forth in any one of SEQ ID NOS: 1, 20, 58, 60, 62, and 68 with (ii) at least one SOST peptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO: 2-19, and 21-54 under conditions and for a time sufficient to permit formation of an antibody/SOST peptide complex; and

(b) detecting a level of antibody/SOST peptide complex, and thereby detecting the presence of an antibody that increases bone mineral content.

45. The method according to any one of claims 41-44 wherein the antibody is selected from the group consisting of an antibody that is present in a biological sample and a purified antibody.

46. The method according to claim 45 wherein the purified antibody is selected from the group consisting of a polyclonal antibody, a monoclonal antibody, a chimeric antibody, a humanized antibody, and an antigen-binding fragment of an antibody.

47. The antibody according to claim 2 wherein the polypeptide comprises an amino acid



sequence selected from the group consisting of SEQ ID NO:5, 6, and 14.

48. The antibody according to any one of claims 3-6 wherein the peptide comprises an amino acid sequence selected from the group consisting of SEQ ID NO:5, 6, 10, 11, 14, and 18.

49. The immunogen of any one of claims 26-29 wherein the peptide comprises an amino acid sequence selected from the group consisting of SEQ ID NO:5, 6, 10, 11, 14, and 18.

50. A method for producing an antibody that specifically binds to a SOST polypeptide, comprising immunizing a non-human animal with an immunogen according to claim 49, wherein the SOST polypeptide comprises an amino acid sequence set forth in SEQ ID NO:1, 20, 28, 60, 62, or 68.

51. The method of any one of claims 41, 42, and 44 wherein the SOST peptide comprises an amino acid sequence selected from the group consisting of SEQ ID NO:5, 6, 10, 11, 14, and 18.




FIG. 1




FIG. 2




FIG. 3

4/7

	20	40	60	
NOGG_HUMAN :	QHYLHIRPA--PSDNLPLVDLIEHPPYFDPKEDLNELRLSLGGSHYDPSFMATSP	EDRPG :	62	
NOGG_CHICK :	QHYLHIRPA--PSDNLPLVDLIEHPPYFDPKEDLNELRLSLGGSHYDPSFMATSP	EDRLG :	62	
NOGG_XENLA :	QHYLHIRPA--PSDNLPLVDLIEHPPYFDPKEDLNELRLSLGGSHYDPSFMATILP	EERLG :	62	
NOGG_FUGRU :	QPYVLLRPY--PSDSLPIVELKEDPPYFDPKEDLNELKSVLG-DFDSRFLSVLPPAEDGHA	:	62	
NOGG_ZEBRA :	QHYVLLRPY--PSDSLPIVELKEDPPYFDPKEDLNELRAILGGSHFQNFMSINPP	EDKHA :	62	
SOST_HUMAN :	QEWDA--FKNDATETIP--ELGEMPEP--PPELENNOTMRAENGSRP-PHPFETKDV	----	52	
SOST_RAT :	QEWDA--FKNDATETIP--ELGEMPEP--PPELENNOTMRAENGSRP-PHPYDTKDV	----	52	
SOST_MOUSE :	QEWDA--FKNDATETIP--ELGEMPEP--PPELENNOTMRAENGSRP-PHPYDAKDV	----	50	

	80	100	120	
NOGG_HUMAN :	GGGGAAGGAEDLAELDQLLRORPSGAMPSEIKGLEFSEGLAQGKKQRLSKLRRKLOMLWSQTF	:	127	
NOGG_CHICK :	-----VDLAELDQLLRORPSGAMPSEIKGLEFYDGLQPGKKHRLSKLRRKLOMLWSQTF	:	119	
NOGG_XENLA :	-----VEDLGELDQLLRORPSGAMPSEIKGLEFYEGLOS-KKHRLSKLRRKLOMLWSQTF	:	118	
NOGG_FUGRU :	G-----NDELDDFD-AQR--WEGALPKETRAVDG-DAPQLGKKHKPSKKLRRKLOMLWYSF	:	116	
NOGG_ZEBRA :	G-----QDELNESE-LMKQRPNGIMPKEIKAMEF-DIQ-HGKKHKPSKKLRRKLOMLWSYTF	:	117	
SOST_HUMAN :	-----	SEYS :	56	
SOST_RAT :	-----	SEYS :	56	
SOST_MOUSE :	-----	SEYS :	54	

	140	160	180	
NOGG_HUMAN :	CP-VLYA--INDLGSRFWRFRVKGSCFSKRSCSVPEGM-----VCKPKSVHL	:	173	
NOGG_CHICK :	CP-VLYT--INDLGSRFWRFRVKGSCFSKRSCSVPEGM-----VCKPKSVHL	:	165	
NOGG_XENLA :	CP-VLYT--INDLGSRFWRFRVKGSCFSKRSCSVPEGM-----VCKAAGSMHL	:	164	
NOGG_FUGRU :	CP-LAHA--INDLGSRFWRFRVKGSCFSKRSCSVPEGM-----TCKPATSTHL	:	162	
NOGG_ZEBRA :	CP-VVHT--INDLGSRFWRFRVKGSCYINKRSCSVPEGM-----VCKPKSSHL	:	163	
SOST_HUMAN :	CRRLHFTRYVTTGPCRSAGVITELVCS--GQCEPARILPNAIGRGKWRPSPGPDFRCIPDRYRAQ	:	119	
SOST_RAT :	CRRLHYTFVTTGPCRSAGVITELVCS--GQCEPARILPNAIGRVKWRPSPGPDFRCIPDRYRAQ	:	119	
SOST_MOUSE :	CRRLHYTFELTTGPCRSAGVITELVCS--GQCEPARILPNAIGRVKWRPSPGPDFRCIPDRYRAQ	:	117	

FIG. 4A

	200	220	240	260	
NOGG_HUMAN :	TVLRWRCQ-RRGSRGWIPIQYHILSEKQSC	-----			: 205
NOGG_CHICK :	TVLRWRCQ-RRGSRCTWIPQYHILAEKQSC	-----			: 197
NOGG_XENLA :	TVLRWRCQ-RRVQKCAWITIQYHILSEKQSC	-----			: 196
NOGG_FUGRU :	TVLRWRCVQRKVELKCAWIPMQYHIVTDKQSC	-----			: 195
NOGG_ZEBRA :	TVLRWRCVQRKVELKCAWIPVQYHIVSEKQSC	-----			: 196
SOST_HUMAN :	RVQLUCP--GG--EAPRARKVRLVASQKQRLTRFHNQSELKDFGTEARPQKGRKPRPRARS				: 178
SOST_RAT :	RVQLUCP--GG--AAPRSRKVRLVASQKQRLTRFHNQSELKDFGPETARPQKGRKPRPRARG				: 178
SOST_MOUSE :	RVQLUCP--GG--AAPRSRKVRLVASQKQRLTRFHNQSELKDFGPETARPQKGRKPRPGARG				: 176

NOGG\_HUMAN : ----- : -  
 NOGG\_CHICK : ----- : -  
 NOGG\_XENLA : ----- : -  
 NOGG\_FUGRU : ----- : -  
 NOGG\_ZEBRA : ----- : -  
 SOST\_HUMAN : AKANQAELENAY : 190  
 SOST\_RAT : AKANQAELENAY : 190  
 SOST\_MOUSE : AKANQAELENAY : 188

FIG. 4B




FIG. 5




FIG. 6



**1**

<110> Winkler, David G.  
Shi, Jiye  
Latham, John

<212> PRT

2

&lt;213&gt; Homo sapiens

&lt;400&gt; 2

Gln Gly Trp Gln Ala Phe Lys Asn Asp Ala Thr Glu Ile Ile Pro Glu  
 1 5 10 15  
 Leu Gly Glu Tyr  
 20

&lt;210&gt; 3

&lt;211&gt; 20

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 3

Thr Glu Ile Ile Pro Glu Leu Gly Glu Tyr Pro Glu Pro Pro Pro Glu  
 1 5 10 15  
 Leu Glu Asn Asn  
 20

&lt;210&gt; 4

&lt;211&gt; 20

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 4

Pro Glu Pro Pro Pro Glu Leu Glu Asn Asn Lys Thr Met Asn Arg Ala  
 1 5 10 15  
 Glu Asn Gly Gly  
 20

&lt;210&gt; 5

&lt;211&gt; 20

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 5

Lys Thr Met Asn Arg Ala Glu Asn Gly Gly Arg Pro Pro His His Pro  
 1 5 10 15  
 Phe Glu Thr Lys  
 20

&lt;210&gt; 6

&lt;211&gt; 16

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 6

Arg Pro Pro His His Pro Phe Glu Thr Lys Asp Val Ser Glu Tyr Ser  
 1 5 10 15

&lt;210&gt; 7

&lt;211&gt; 21

&lt;212&gt; PRT

<213> Artificial Sequence

**<220>**

<223> Human SOST peptide fragment with additional  
cysteine added

<400> 7

Gln Gly Trp Gln Ala Phe Lys Asn Asp Ala Thr Glu Ile Ile Pro Glu  
1 5 10 15  
Leu Gly Glu Tyr Cys  
20

<210> 8

<211> 21

<212> PRT

<213> Artificial Sequence

**<220>**

<223> Human SOST peptide fragment with additional  
cysteine added

<400> 8

Thr Glu Ile Ile Pro Glu Leu Gly Glu Tyr Pro Glu Pro Pro Pro Glu  
1 5 10 15  
Leu Glu Asn Asn Cys  
20

<210> 9

<211> 21

<212> PRT

<213> Artificial Sequence

**<220>**

<223> Human SOST peptide fragment with additional  
cysteine added

<400> 9

Pro Glu Pro Pro Pro Glu Leu Glu Asn Asn Lys Thr Met Asn Arg Ala  
1 5 10 15  
Glu Asn Gly Gly Cys  
20

<210> 10

<211> 21

<212> PRT

<213> Artificial Sequence

**<220>**

<223> Human SOST peptide fragment with additional cysteine added

<400> 10

Lys Thr Met Asn Arg Ala Glu Asn Gly Gly Arg Pro Pro His His Pro  
1 5 10 15  
Phe Glu Thr Lys Cys

20

<210> 11  
 <211> 17  
 <212> PRT  
 <213> Artificial Sequence

<220>  
 <223> Human SOST peptide fragment with additional  
 cysteine added

<400> 11  
 Arg Pro Pro His His Pro Phe Glu Thr Lys Asp Val Ser Glu Tyr Ser  
 1 5 10 15  
 Cys

<210> 12  
 <211> 24  
 <212> PRT  
 <213> Rattus norvegicus

<400> 12  
 Gln Gly Trp Gln Ala Phe Lys Asn Asp Ala Thr Glu Ile Ile Pro Gly  
 1 5 10 15  
 Leu Arg Glu Tyr Pro Glu Pro Pro  
 20

<210> 13  
 <211> 20  
 <212> PRT  
 <213> Rattus morvegicus

<400> 13  
 Pro Glu Pro Pro Gln Glu Leu Glu Asn Asn Gln Thr Met Asn Arg Ala  
 1 5 10 15  
 Glu Asn Gly Gly  
 20

<210> 14  
 <211> 20  
 <212> PRT  
 <213> Rattus norvegicus

<400> 14  
 Glu Asn Gly Gly Arg Pro Pro His His Pro Tyr Asp Thr Lys Asp Val  
 1 5 10 15  
 Ser Glu Tyr Ser  
 20

<210> 15  
 <211> 20  
 <212> PRT

5

&lt;213&gt; Rattus norvegicus

&lt;400&gt; 15

Thr	Glu	Ile	Ile	Pro	Gly	Leu	Arg	Glu	Tyr	Pro	Glu	Pro	Pro	Gln	Glu
1				5				10						15	
Leu	Glu	Asn	Asn												
			20												

&lt;210&gt; 16

&lt;211&gt; 25

&lt;212&gt; PRT

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; Rat SOST peptide fragment with additional cysteine added

&lt;400&gt; 16

Gln	Gly	Trp	Gln	Ala	Phe	Lys	Asn	Asp	Ala	Thr	Glu	Ile	Ile	Pro	Gly
1				5				10						15	
Leu	Arg	Glu	Tyr	Pro	Glu	Pro	Pro	Cys							
			20					25							

&lt;210&gt; 17

&lt;211&gt; 21

&lt;212&gt; PRT

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; Rat SOST peptide fragment with additional cysteine added

&lt;400&gt; 17

Pro	Glu	Pro	Pro	Gln	Glu	Leu	Glu	Asn	Asn	Gln	Thr	Met	Asn	Arg	Ala
1				5				10						15	
Glu	Asn	Gly	Gly	Cys											
			20												

&lt;210&gt; 18

&lt;211&gt; 21

&lt;212&gt; PRT

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; Rat SOST peptide fragment with additional cysteine added

&lt;400&gt; 18

Glu	Asn	Gly	Gly	Arg	Pro	Pro	His	His	Pro	Tyr	Asp	Thr	Lys	Asp	Val
1				5				10						15	
Ser	Glu	Tyr	Ser	Cys											
			20												

&lt;210&gt; 19

6

<211> 21  
 <212> PRT  
 <213> Artificial Sequence

<220>  
 <223> Rat SOST peptide fragment with additional cysteine added

<400> 19  
 Thr Glu Ile Ile Pro Gly Leu Arg Glu Tyr Pro Glu Pro Pro Gln Glu  
 1 5 10 15  
 Leu Glu Asn Asn Cys  
 20

<210> 20  
 <211> 190  
 <212> PRT  
 <213> Rattus norvegicus

<400> 20  
 Gln Gly Trp Gln Ala Phe Lys Asn Asp Ala Thr Glu Ile Ile Pro Gly  
 1 5 10 15  
 Leu Arg Glu Tyr Pro Glu Pro Pro Gln Glu Leu Glu Asn Asn Gln Thr  
 20 25 30  
 Met Asn Arg Ala Glu Asn Gly Gly Arg Pro Pro His His Pro Tyr Asp  
 35 40 45  
 Thr Lys Asp Val Ser Glu Tyr Ser Cys Arg Glu Leu His Tyr Thr Arg  
 50 55 60  
 Phe Val Thr Asp Gly Pro Cys Arg Ser Ala Lys Pro Val Thr Glu Leu  
 65 70 75 80  
 Val Cys Ser Gly Gln Cys Gly Pro Ala Arg Leu Leu Pro Asn Ala Ile  
 85 90 95  
 Gly Arg Val Lys Trp Trp Arg Pro Asn Gly Pro Asp Phe Arg Cys Ile  
 100 105 110  
 Pro Asp Arg Tyr Arg Ala Gln Arg Val Gln Leu Leu Cys Pro Gly Gly  
 115 120 125  
 Ala Ala Pro Arg Ser Arg Lys Val Arg Leu Val Ala Ser Cys Lys Cys  
 130 135 140  
 Lys Arg Leu Thr Arg Phe His Asn Gln Ser Glu Leu Lys Asp Phe Gly  
 145 150 155 160  
 Pro Glu Thr Ala Arg Pro Gln Lys Gly Arg Lys Pro Arg Pro Arg Ala  
 165 170 175  
 Arg Gly Ala Lys Ala Asn Gln Ala Glu Leu Glu Asn Ala Tyr  
 180 185 190

<210> 21  
 <211> 20  
 <212> PRT  
 <213> Homo sapiens

<400> 21  
 Ile Pro Asp Arg Tyr Arg Ala Gln Arg Val Gln Leu Leu Cys Pro Gly  
 1 5 10 15  
 Gly Glu Ala Pro  
 20

7

<210> 22  
<211> 20  
<212> PRT  
<213> Homo sapiens

<400> 22  
Gln Leu Leu Cys Pro Gly Gly Glu Ala Pro Arg Ala Arg Lys Val Arg  
1 5 10 15  
Leu Val Ala Ser  
20

<210> 23  
<211> 21  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> Human SOST peptide fragment with additional  
cysteine added

<400> 23  
Ile Pro Asp Arg Tyr Arg Ala Gln Arg Val Gln Leu Leu Cys Pro Gly  
1 5 10 15  
Gly Glu Ala Pro Cys  
20

<210> 24  
<211> 21  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> Human SOST peptide fragment with additional  
cysteine added

<400> 24  
Gln Leu Leu Cys Pro Gly Gly Glu Ala Pro Arg Ala Arg Lys Val Arg  
1 5 10 15  
Leu Val Ala Ser Cys  
20

<210> 25  
<211> 17  
<212> PRT  
<213> Rattus Norvegicus

<400> 25  
Ile Pro Asp Arg Tyr Arg Ala Gln Arg Val Gln Leu Leu Cys Pro Gly  
1 5 10 15  
Gly

<210> 26

<211> 16  
<212> PRT  
<213> Rattus norvegicus.

<400> 26  
Pro Gly Gly Ala Ala Pro Arg Ser Arg Lys Val Arg Leu Val Ala Ser  
1 5 10 15

<210> 27  
<211> 18  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> Rat SOST peptide fragment with additional cysteine  
added

<400> 27  
Ile Pro Asp Arg Tyr Arg Ala Gln Arg Val Gln Leu Leu Ser Pro Gly  
1 5 10 15  
Gly Cys

<210> 28  
<211> 17  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> Rat SOST peptide fragment with additional cysteine  
added

<400> 28  
Pro Gly Gly Ala Ala Pro Arg Ser Arg Lys Val Arg Leu Val Ala Ser  
1 5 10 15  
Cys

<210> 29  
<211> 20  
<212> PRT  
<213> Homo sapiens

<400> 29  
Cys Gly Pro Ala Arg Leu Leu Pro Asn Ala Ile Gly Arg Gly Lys Trp  
1 5 10 15  
Trp Arg Pro Ser  
20

<210> 30  
<211> 16  
<212> PRT  
<213> Homo sapiens



&lt;400&gt; 30

Ile	Gly	Arg	Gly	Lys	Trp	Trp	Arg	Pro	Ser	Gly	Pro	Asp	Phe	Arg	Cys
1				5					10					15	

&lt;210&gt; 31

&lt;211&gt; 18

&lt;212&gt; PRT

&lt;213&gt; Rattus norvegicus

&lt;400&gt; 31

Pro	Asn	Ala	Ile	Gly	Arg	Val	Lys	Trp	Trp	Arg	Pro	Asn	Gly	Pro	Asp
1				5					10					15	

Phe Arg

&lt;210&gt; 32

&lt;211&gt; 19

&lt;212&gt; PRT

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; Rat SOST peptide fragment with additional cysteine added

&lt;400&gt; 32

Pro	Asn	Ala	Ile	Gly	Arg	Val	Lys	Trp	Trp	Arg	Pro	Asn	Gly	Pro	Asp
1				5					10					15	

Phe Arg Cys

&lt;210&gt; 33

&lt;211&gt; 20

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 33

Lys	Arg	Leu	Thr	Arg	Phe	His	Asn	Gln	Ser	Glu	Leu	Lys	Asp	Phe	Gly
1				5					10					15	

Thr Glu Ala Ala

20

&lt;210&gt; 34

&lt;211&gt; 20

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 34

Glu	Leu	Lys	Asp	Phe	Gly	Thr	Glu	Ala	Ala	Arg	Pro	Gln	Lys	Gly	Arg
1				5					10					15	

Lys Pro Arg Pro

20

&lt;210&gt; 35

10

&lt;211&gt; 20

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 35

Arg	Pro	Gln	Lys	Gly	Arg	Lys	Pro	Arg	Pro	Arg	Ala	Arg	Ser	Ala	Lys
1				5				10						15	
Ala	Asn	Gln	Ala												
				20											

&lt;210&gt; 36

&lt;211&gt; 16

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 36

Arg	Ala	Arg	Ser	Ala	Lys	Ala	Asn	Gln	Ala	Glu	Leu	Glu	Asn	Ala	Tyr
1				5				10						15	

&lt;210&gt; 37

&lt;211&gt; 21

&lt;212&gt; PRT

&lt;213&gt; Artificial Sequence

&lt;220&gt;

<223> Human SOST peptide fragment with additional  
cysteine added

&lt;400&gt; 37

Lys	Arg	Leu	Thr	Arg	Phe	His	Asn	Gln	Ser	Glu	Leu	Lys	Asp	Phe	Gly
1				5				10						15	
Thr	Glu	Ala	Ala	Cys											
				20											

&lt;210&gt; 38

&lt;211&gt; 21

&lt;212&gt; PRT

&lt;213&gt; Artificial Sequence

&lt;220&gt;

<223> Human SOST peptide fragment with additional  
cysteine added

&lt;400&gt; 38

Glu	Leu	Lys	Asp	Phe	Gly	Thr	Glu	Ala	Ala	Arg	Pro	Gln	Lys	Gly	Arg
1				5				10						15	
Lys	Pro	Arg	Pro	Cys											
				20											

&lt;210&gt; 39

&lt;211&gt; 21

&lt;212&gt; PRT

&lt;213&gt; Artificial Sequence

&lt;220&gt;

<223> Human SOST peptide fragment with additional  
cysteine added

&lt;400&gt; 39

Arg	Pro	Gln	Lys	Gly	Arg	Lys	Pro	Arg	Pro	Arg	Ala	Arg	Ser	Ala	Lys
1				5				10						15	
Ala	Asn	Gln	Ala	Cys											
			20												

&lt;210&gt; 40

&lt;211&gt; 17

&lt;212&gt; PRT

&lt;213&gt; Artificial Sequence

&lt;220&gt;

<223> Human SOST peptide fragment with additional  
cysteine added

&lt;400&gt; 40

Arg	Ala	Arg	Ser	Ala	Lys	Ala	Asn	Gln	Ala	Glu	Leu	Glu	Asn	Ala	Tyr
1				5				10						15	
Cys															

&lt;210&gt; 41

&lt;211&gt; 23

&lt;212&gt; PRT

&lt;213&gt; Rattus norvegicus

&lt;400&gt; 41

Lys	Arg	Leu	Thr	Arg	Phe	His	Asn	Gln	Ser	Glu	Leu	Lys	Asp	Phe	Gly
1				5				10						15	
Pro	Glu	Thr	Ala	Arg	Pro	Gln									
			20												

&lt;210&gt; 42

&lt;211&gt; 23

&lt;212&gt; PRT

&lt;213&gt; Rattus norvegicus

&lt;400&gt; 42

Lys	Gly	Arg	Lys	Pro	Arg	Pro	Arg	Ala	Arg	Gly	Ala	Lys	Ala	Asn	Gln
1				5				10						15	
Ala	Glu	Leu	Glu	Glu	Asn	Ala	Tyr								
			20												

&lt;210&gt; 43

&lt;211&gt; 24

&lt;212&gt; PRT

&lt;213&gt; Rattus norvegicus

&lt;400&gt; 43

Ser Glu Leu Lys Asp Phe Gly Pro Glu Thr Ala Arg Pro Gln Lys Gly

Arg Lys Pro Arg Pro Arg Ala Arg

20

```
<210> 44
<211> 24
<212> PRT
<213> Artificial Sequence
```

<220>  
<223> Rat SOST peptide fragment with additional cysteine added

```
<400> 44
Lys Arg Leu Thr Arg Phe His Asn Gln Ser Glu Leu Lys Asp Phe Gly
 1             5             10             15
Pro Glu Thr Ala Arg Pro Gln Cys
      20
```

```
<210> 45
<211> 24
<212> PRT
<213> Artificial Sequence
```

```
<220>
<223> Rat SOST peptide fragment with additional cysteine
      added
```

```

<400> 45
Lys Gly Arg Lys Pro Arg Pro Arg Ala Arg Gly Ala Lys Ala Asn Gln
 1          5          10          15
Ala Glu Leu Glu Asn Ala Tyr Cys
      20

```

```
<210> 46
<211> 25
<212> PRT
<213> Artificial Sequence
```

```
<220>
<223> Rat SOST peptide fragment with additional cysteine
      added
```

```
<400> 46
Ser Glu Leu Lys Asp Phe Gly Pro Glu Thr Ala Arg Pro Gln Lys Gly
 1          5          10          15
Arg Lys Pro Arg Pro Arg Ala Arg Cys
      20          25
```

```
<210> 47
<211> 56
<212> PRT
<213> Homo sapiens
```

13

&lt;400&gt; 47

Gln Gly Trp Gln Ala Phe Lys Asn Asp Ala Thr Glu Ile Ile Pro Glu  
 1 5 10 15  
 Leu Gly Glu Tyr Pro Glu Pro Pro Pro Glu Leu Glu Asn Asn Lys Thr  
 20 25 30  
 Met Asn Arg Ala Glu Asn Gly Gly Arg Pro Pro His His Pro Phe Glu  
 35 40 45  
 Thr Lys Asp Val Ser Glu Tyr Ser  
 50 55

&lt;210&gt; 48

&lt;211&gt; 56

&lt;212&gt; PRT

&lt;213&gt; Rattus norvegicus

&lt;400&gt; 48

Gln Gly Trp Gln Ala Phe Lys Asn Asp Ala Thr Glu Ile Ile Pro Gly  
 1 5 10 15  
 Leu Arg Glu Tyr Pro Glu Pro Pro Gln Glu Leu Glu Asn Asn Gln Thr  
 20 25 30  
 Met Asn Arg Ala Glu Asn Gly Gly Arg Pro Pro His His Pro Tyr Asp  
 35 40 45  
 Thr Lys Asp Val Ser Glu Tyr Ser  
 50 55

&lt;210&gt; 49

&lt;211&gt; 32

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 49

Cys Ile Pro Asp Arg Tyr Arg Ala Gln Arg Val Gln Leu Leu Cys Pro  
 1 5 10 15  
 Gly Gly Glu Ala Pro Arg Ala Arg Lys Val Arg Leu Val Ala Ser Cys  
 20 25 30

&lt;210&gt; 50

&lt;211&gt; 32

&lt;212&gt; PRT

&lt;213&gt; Rattus norvegicus

&lt;400&gt; 50

Cys Ile Pro Asp Arg Tyr Arg Ala Gln Arg Val Gln Leu Leu Cys Pro  
 1 5 10 15  
 Gly Gly Ala Ala Pro Arg Ser Arg Lys Val Arg Leu Val Ala Ser Cys  
 20 25 30

&lt;210&gt; 51

&lt;211&gt; 44

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 51

Leu Thr Arg Phe His Asn Gln Ser Glu Leu Lys Asp Phe Gly Thr Glu

14

1 5 10 15  
 Ala Ala Arg Pro Gln Lys Gly Arg Lys Pro Arg Pro Arg Ala Arg Ser  
 20 25 30  
 Ala Lys Ala Asn Gln Ala Glu Leu Glu Asn Ala Tyr  
 35 40

<210> 52  
 <211> 44  
 <212> PRT  
 <213> Rattus norvegicus

<400> 52  
 Leu Thr Arg Phe His Asn Gln Ser Glu Leu Lys Asp Phe Gly Pro Glu  
 1 5 10 15  
 Thr Ala Arg Pro Gln Lys Gly Arg Lys Pro Arg Pro Arg Ala Arg Gly  
 20 25 30  
 Ala Lys Ala Asn Gln Ala Glu Leu Glu Asn Ala Tyr  
 35 40

<210> 53  
 <211> 26  
 <212> PRT  
 <213> Homo sapiens

<400> 53  
 Cys Gly Pro Ala Arg Leu Leu Pro Asn Ala Ile Gly Arg Gly Lys Trp  
 1 5 10 15  
 Trp Arg Pro Ser Gly Pro Asp Phe Arg Cys  
 20 25

<210> 54  
 <211> 26  
 <212> PRT  
 <213> Rattus norvegicus

<400> 54  
 Cys Gly Pro Ala Arg Leu Leu Pro Asn Ala Ile Gly Arg Val Lys Trp  
 1 5 10 15  
 Trp Arg Pro Asn Gly Pro Asp Phe Arg Cys  
 20 25

<210> 55  
 <211> 570  
 <212> DNA  
 <213> Homo sapiens

<400> 55  
 caggggtggc aggcgttcaa gaatgatgcc acggaaatca tccccgagct cggagagtac 60  
 cccgagcctc caccggagct ggagaacaac aagaccatga accgggcgga gaacggaggg 120  
 cggcctcccc accacccctt tgagaccaa gacgtgtccg agtacagctg ccgcgagctg 180  
 cacttcaccc gctacgtgac cgatgggccc tgccgcagcg ccaagccggt caccgagctg 240  
 gtgtgctccg gccagtgcgg cccggcgcg cgtgtgccca acgcatcgg ccgcggcaag 300  
 tggtagcgac ctagtgggccc cgacttccgc tgcattcccc accgctaccg cgcgcagcgc 360  
 gtgcagctgc tgtgtccccg tggtagggcg ccgcgcgcgc gcaaggtgcg cctggtggcc 420

## 15

```

tcgtgcaagt gcaagcgcct caccgccttc cacaaccagt cggagctcaa ggacttcggg 480
accgaggccg ctcgcccgca gaaggccggc aagccgcggc cccgcgcccg gagcgccaaa 540
gcccaaccagg ccgagctgga gaacgcctac                                     570

```

```

<210> 56
<211> 570
<212> DNA
<213> Rattus norvegicus

```

```

<400> 56
caggggtggc aagccttcaa gaatgatgcc acagaaatca tcccgggact cagagagtac 60
ccagagcctc ctccaggaact agagaacaac cagaccatga accgggcccga gaacggaggc 120
agaccccccc accatcctta tgacacccaa gacgtgtccg agtacagctg ccgcgagctg 180
cactacaccc gcttcgtgac cgacggcccc tgccgcagtg ccaagccggg caccgagttg 240
gtgtgctcgg gccagtgcgg ccccgcgccg ctgctgcccc acgccatcgg gcgcgtgaag 300
tggtggcgcc cgaacggacc cgacttcggc tgcattcccg atcgctaccg cgcgagcgcg 360
gtgcagctgc tgtgccccgg cggcgcgccg ccgcgctcgc gcaagggtgcg tctggtaggc 420
tcgtgcaagt gcaagcgcct caccgccttc cacaaccagt cggagctcaa ggacttcgga 480
cctgagaccg cgcggccgca gaagggtcgc aagccgcggc cccgcgcccg gggagccaaa 540
gcccaaccagg ccgagctgga gaacgcctac                                     570

```

```

<210> 57
<211> 2301
<212> DNA
<213> Homo sapiens

```

```

<400> 57
agagcctgtg ctactggaag gtggcgtgcc ctccctctggc tggtagcatg cagctcccac 60
tggccctgtg tctcgtctgc ctgctggtag acacagcctt ccgtgtagtg gagggccagg 120
ggtggcaggc gttcaagaat gatgccacgg aaatcatccc cgagctcgga gactaccccg 180
agcctccacc ggagctggag aacaacaaga ccatgaaccg ggcggagaac ggagggcgcc 240
ctccccacca cccctttgag accaaagacg tgtccgagta cagctgccgc gagctgcact 300
tcacccgcta cgtgaccgat gggcctgtgc gcagcgccaa gccggtcacc gagctggtgt 360
gtcccgccca gtgcggcccc gcgcgcctgc tgcccaacgc catcgccgcg ggcaagtggg 420
ggcgacactg tgggcccagc ttccgctgca tccccgaccg ctaccgcgcg cagcgcgctg 480
agctgctgtg tcccgggtggg gaggcgcccc gcgcgcgcaa ggtgcgcctg gtggcctcgt 540
gcaagtgcga ggcctcacc cgcttcacca accagtcgga gctcaaggac ttcgggaccg 600
aggcgctcgc gcgcgagaag ggccggaagc cgcggccccc cgcccggagc gccaaagcca 660
accaggccga gctggagaac gcctactaga gcccgccccg gcccctcccc accggcgggc 720
gccccggccc tgaacccgcg cccacattt ctgtcctctg cgcgtgggtt gattgtttat 780
atttcattgt aaatgcctgc aaccagggc agggggctga gacctccag gccctgagga 840
atcccgggcg ccggcaaggc cccctcagc ccgccaagct aggggtccca cggggcaggg 900
gagggaaatt agagtacag aactgagcc acgcagcccc gcctctgggg ccgcctacct 960
ttgctggctc cacttcagag gaggcagaaa tggaaacatt ttcaccgccc tggggtttta 1020
agggagcggg gtgggagtg gaaagtccag ggactggtta agaaagttgg ataagattcc 1080
cccttgccac tcgctgcccc tcagaaagcc tgaggcgtgc ccagagcaca agactggggg 1140
caactgtaga tgtggtttct agtcctggct ctgccactaa cttgctgtgt aacctgaac 1200
tacacaattc tccttcggga cctcaatttc cactttgtaa aatgaggggt gaggtgggaa 1260
taggatctcg aggagactat tggcatatga ttccaaggac tccagtgcc tttgaatggg 1320
cagaggtgag agagagagag agaaagagag agaataatg cagttgcatt gattcagtgc 1380
caaggtcact tccagaattc agagttgtga tgcctctctc tgacagccaa agatgaaaaa 1440
caaacagaaa aaaaaaagta aagagtctat ttatggctga catatttacg gctgacaaac 1500
tcctggaaga agctatgctg cttcccagcc tggcttcccc ggatgttttg ctacctccac 1560
ccctccatct caaagaaata acatcatcca ttggggtaga aaaggagagg gtccgagggt 1620
gggtgggagg atagaaatca catccgcccc aacttcccaa agagcagcat ccctcccccg 1680
acccatagcc atgttttaaa gtcaccttcc gaagagaagt gaaagggtca aggacactgg 1740
ccttgaggc ccgagggagc agccatcaca aactcacaga ccagcacatc ccttttgaga 1800
caccgccttc tgcccaccac tcacggacac atttctgcct agaaaacagc ttcttactgc 1860

```

16

```

tcttacatgt gatggcatat cttacactaa aagaatatta ttgggggaaa aactacaagt 1920
gctgtacata tgctgagaaa ctgcagagca taatagctgc cacccaaaaa tctttttgaa 1980
aatcatttcc agacaacctc ttactttctg tgtagttttt aattgttaaa aaaaaaaagt 2040
tttaaacaga agcacatgac atatgaaagc ctgcaggact ggtagttttt ttggcaattc 2100
ttccacgtgg gacttgtcca caagaatgaa agtagtggtt tttaaagagt taagttacat 2160
atatttttcc tcaacttaagt tatttatgca aaagtttttc ttgtagagaa tgacaatgtt 2220
aatattgctt tatgaattaa cagtctgttc ttccagagtc cagagacatt gttaataaag 2280
acaatgaatc atgaccgaaa g                                     2301

```

&lt;210&gt; 58

&lt;211&gt; 213

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 58

```

Met Gln Leu Pro Leu Ala Leu Cys Leu Val Cys Leu Leu Val His Thr
  1           5           10           15
Ala Phe Arg Val Val Glu Gly Gln Gly Trp Gln Ala Phe Lys Asn Asp
      20           25           30
Ala Thr Glu Ile Ile Pro Glu Leu Gly Glu Tyr Pro Glu Pro Pro Pro
      35           40           45
Glu Leu Glu Asn Asn Lys Thr Met Asn Arg Ala Glu Asn Gly Gly Arg
      50           55           60
Pro Pro His His Pro Phe Glu Thr Lys Asp Val Ser Glu Tyr Ser Cys
      65           70           75           80
Arg Glu Leu His Phe Thr Arg Tyr Val Thr Asp Gly Pro Cys Arg Ser
      85           90           95
Ala Lys Pro Val Thr Glu Leu Val Cys Ser Gly Gln Cys Gly Pro Ala
      100          105          110
Arg Leu Leu Pro Asn Ala Ile Gly Arg Gly Lys Trp Trp Arg Pro Ser
      115          120          125
Gly Pro Asp Phe Arg Cys Ile Pro Asp Arg Tyr Arg Ala Gln Arg Val
      130          135          140
Gln Leu Leu Cys Pro Gly Gly Glu Ala Pro Arg Ala Arg Lys Val Arg
      145          150          155          160
Leu Val Ala Ser Cys Lys Cys Lys Arg Leu Thr Arg Phe His Asn Gln
      165          170          175
Ser Glu Leu Lys Asp Phe Gly Thr Glu Ala Ala Arg Pro Gln Lys Gly
      180          185          190
Arg Lys Pro Arg Pro Arg Ala Arg Ser Ala Lys Ala Asn Gln Ala Glu
      195          200          205
Leu Glu Asn Ala Tyr
      210

```

&lt;210&gt; 59

&lt;211&gt; 2301

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 59

```

agagcctgtg ctactggaag gtggcgtgcc ctccctctggc tggtagcatg cagctccac 60
tgccctgtg tctcatctgc ctgctggtac acacagcctt ccgtgtagtg gagggccagg 120
ggtggcaggc gttcaagaat gatgccacgg aaatcatccg cgagctcgga gtagtccccg 180
agcctccacc ggagctggag aacaacaaga ccatgaaccg ggcggagaac ggagggcggc 240
ctccccacca cccctttgag accaaagacg tgtccgagta cagctgccgc gagctgcact 300
tcaccgcgta cgtgaccgat gggccgtgcc gcagcgccaa gccggtcacc gagctggtgt 360
gctccggcca gtgcggcccg gcgcgcctgc tgcccaacgc catcggccgc ggcaagtggg 420

```



```

ggcgacctag tgggccccgac ttccgctgca tccccgaccg ctaccgcgcg cagcgcgtgc 480
agctgctgtg tcccgggtggt gaggcgcgcg gcgcgcgcaa ggtgcgcctg gtggcctcgt 540
gcaagtgcaa gcgcctcacc cgcttcacaca accagtgcga gctcaaggac ttcgggaccg 600
aggcgcgtcg gccgcagaag ggccggaagc cgcggccccc cgcgggagc gccaaagcca 660
accaggccga gctggagaac gcctactaga gcccgcgcgc gccctcccc accggcgggc 720
gccccggccc tgaacccgcg cccacattt ctgtcctctg cgcgtggtt gattgtttat 780
atttcattgt aaatgcctgc aacccagggc agggggctga gaccttcag gccctgagga 840
atcccgggcg cgggcaaggc cccctcagc ccgcagctg aggggtccca cggggcaggg 900
gaggggaattg agagtacag acactgagcc acgcagccc gcctctgggg ccgcctacct 960
ttgctggtcc cacttcagag gaggcagaaa tggaaagcatt ttcaccgccc tggggtttta 1020
agggagcggg gtgggagtgg gaaagtccag ggactgggta agaaagtgg ataagattcc 1080
cccttgacc tcgctgccc tcagaaagcc tgaggcgtgc ccagagcaca agactggggg 1140
caactgtaga tgtggtttct agtcctggct ctgccactaa cttgctgtgt aacctgaac 1200
tacacaattc tccttcggga cctcaatttc cactttgtaa aatgagggtg gaggtgggaa 1260
taggatctcg aggagactat tggcatatga ttccaaggac tccagtgcct tttgaatggg 1320
cagaggtgag agagagagag agaaagagag agaataatg cagttgcatt gattcagtgc 1380
caaggctcact tccagaattc agagtgtgta tgctctcttc tgacagccaa agatgaaaaa 1440
caaacagaaa aaaaaaagta aagagtctat ttatggctga catatttacg gctgacaaac 1500
tcctggaaga agctatgctg cttcccagcc tggcttcccc ggatgtttgg ctacctccac 1560
ccctccatct caaagaaata acatcatcca ttggggtaga aaaggagagg gtccgagggt 1620
ggtgggaggg atagaaatca catccgcccc aacttcccaa agagcagcat ccctcccccg 1680
acccatagcc atgttttaaa gtcaccttcc gaagagaagt gaaagggtca aggacactgg 1740
ccttgaggc cggaggagc agccatcaca aactcacaga ccagcacatc ccttttgaga 1800
caccgccttc tgcccaccac tcacggacac atttctgcct agaaaacagc ttcttactgc 1860
tcttacatgt gatggcatat cttacactaa aagaatatta ttgggggaaa aactacaagt 1920
gctgtacata tgctgagaaa ctgcagagca taatagctgc caccacaaaa tctttttgaa 1980
aatcatttcc agacaacctc ttactttctg tgtagttttt aattgttaaa aaaaaaagt 2040
tttaaacaga agcacatgac atatgaaagc ctgcaggact ggtcggtttt ttggcaattc 2100
ttccacgtgg gacttgcca caagaatgaa agtagtggt tttaaagagt taagttacat 2160
atttatttcc ccacttaagt tatttatgca aaagttttcc ttgtagagaa tgacaatggt 2220
aatattgctt tatgaattaa cagtctgttc ttccagagtc cagagacatt gttaataaag 2280
acaatgaatc atgaccgaaa g

```

&lt;210&gt; 60

&lt;211&gt; 213

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 60

```

Met Gln Leu Pro Leu Ala Leu Cys Leu Ile Cys Leu Leu Val His Thr
1           5           10           15
Ala Phe Arg Val Val Glu Gly Gln Gly Trp Gln Ala Phe Lys Asn Asp
20           25           30
Ala Thr Glu Ile Ile Arg Glu Leu Gly Glu Tyr Pro Glu Pro Pro Pro
35           40           45
Glu Leu Glu Asn Asn Lys Thr Met Asn Arg Ala Glu Asn Gly Gly Arg
50           55           60
Pro Pro His His Pro Phe Glu Thr Lys Asp Val Ser Glu Tyr Ser Cys
65           70           75           80
Arg Glu Leu His Phe Thr Arg Tyr Val Thr Asp Gly Pro Cys Arg Ser
85           90           95
Ala Lys Pro Val Thr Glu Leu Val Cys Ser Gly Gln Cys Gly Pro Ala
100          105          110
Arg Leu Leu Pro Asn Ala Ile Gly Arg Gly Lys Trp Trp Arg Pro Ser
115          120          125
Gly Pro Asp Phe Arg Cys Ile Pro Asp Arg Tyr Arg Ala Gln Arg Val
130          135          140
Gln Leu Leu Cys Pro Gly Gly Glu Ala Pro Arg Ala Arg Lys Val Arg

```

145                                      150                                      155                                      160  
 Leu Val Ala Ser Cys Lys Cys Lys Arg Leu Thr Arg Phe His Asn Gln  
    165                                      170                                      175  
 Ser Glu Leu Lys Asp Phe Gly Thr Glu Ala Ala Arg Pro Gln Lys Gly  
    180                                      185                                      190  
 Arg Lys Pro Arg Pro Arg Ala Arg Ser Ala Lys Ala Asn Gln Ala Glu  
    195                                      200                                      205  
 Leu Glu Asn Ala Tyr  
    210

&lt;210&gt; 61

&lt;211&gt; 2301

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 61

```

agagcctgtg ctactggaag gtggcgtgcc ctctctggc tgggtaccatg cagctcccac 60
tggccctgtg tctcgtctgc ctgctggtac acacagcctt ccgtgtagtg gagggccagg 120
ggtggcaggc gttcaagaat gatgccacgg aaatcatccg cgagctcggg gagtaccacc 180
agcctccacc ggagctggag aacaacaaga ccatgaaccg ggcggagaac ggagggcggc 240
ctccccacca cccctttgag accaaagacg tgctcgagta cagctgccgc gagctgcact 300
tcacccgcta cgtgaccgat gggccgtgcc gcagcgccaa gccggtcacc gagctggtgt 360
gctccggcca gtgcggcccg gcgcgcctgc tgcccaacgc catcggccgc ggcaagtggg 420
ggcgacctag tgggcccagc ttccgctgca tccccgaccg ctaccgcgcg cagcgcgtgc 480
agctgctgtg tcccgggtgt gaggcgcgcg gcgcgcgcaa ggtgcgcctg gtggcctcgt 540
gcaagtgcga gcgcctcacc cgcttccaca accagtcgga gctcaaggac ttcgggaccg 600
aggccgctcg gccgcagaag ggccggaagc cgccggcccc cgcccgaggc gccaaagcca 660
accaggccga gctggagaac gcctactaga gcccgccccg gccctcccc accggcgggc 720
gccccggccc tgaacccgcg ccccacattt ctgtcctctg cgcgtggttt gattgtttat 780
atctcattgt aaatgcctgc aaccagggc agggggctga gaccttccag gccctgagga 840
atcccgggcg ccggcaaggc cccctcagc ccgccagctg aggggtccca cggggcaggg 900
gaggggaattg agagtacag acactgagcc acgcagcccc gcctctgggg ccgcctacct 960
ttgctggtcc cacttcagag gaggcagaaa tggaaagcatt ttcaccgccc tgggggtttta 1020
agggagcggg gtgggagtg gaaagtccag ggactgggta agaaagtgtg ataagattcc 1080
ccctgcacc tcgctgcccc tcagaaagcc tgaggcgtgc ccagagcaca agactggggg 1140
caactgtaga tgtggtttct agtctggct ctgccactaa ctgtctgtgt aaccttgaac 1200
tacacaattc tccttcggga cctcaatttc cactttgtaa aatgagggtg gaggtgggaa 1260
taggatctcg aggagactat tggcatatga ttccaaggac tccagtgcct tttgaatggg 1320
cagaggtgag agagagagag agaaagagag agaatgaatg cagttgcatt gattcagtgc 1380
caaggtcact tccagaattc agagttgtga tgctctcttc tgacagccaa agatgaaaaa 1440
caaacagaaa aaaaaaagta aagagtctat ttatggctga catatttacg gctgacaaac 1500
tcttggaaga agctatgctg cttcccagcc tggcttcccc ggatgtttgg ctacctccac 1560
ccctccatct caaagaaaata acatcatcca ttggggtaga aaaggagagg gtccgagggg 1620
ggtgggaggg atagaaatca catccgcccc aacttcccaa agagcagcat ccctcccccg 1680
acctatagcc atgtttttaa gtcaccttcc gaagagaagt gaaaggttca aggacactgg 1740
ccttgaggc ccgagggagc agccatcaca aactcacaga ccagcacatc ccttttgaga 1800
caccgccttc tgcccaccac tcacggacac atttctgcct agaaaacagc ttcttactgc 1860
tcttacatgt gatggcatat cttacactaa aagaatatta ttgggggaaa aactacaagt 1920
gctgtacata tgctgagaaa ctgcagagca taatagctgc cacccaaaaa tctttttgaa 1980
aatctttcc agacaacctc ttactttctg tgtagtttt aattgtttaa aaaaaaagt 2040
tttaaacaga agcacatgac atatgaaagc ctgcaggact ggtcgttttt ttggcaattc 2100
ttccacgtgg gacttgtcca caagaatgaa agtagtggtt tttaaagagt taagttacat 2160
atatttttc tactttaagt tatttatgca aaagtttttc ttgtagagaa tgacaatgtt 2220
aatattgctt tatgaattaa cagtctgttc ttccagagtc cagagacatt gttaataaag 2280
acaatgaatc atgaccgaaa g

```

&lt;210&gt; 62

<211> 213  
 <212> PRT  
 <213> Homo sapiens

<400> 62  
 Met Gln Leu Pro Leu Ala Leu Cys Leu Val Cys Leu Leu Val His Thr  
 1 5 10 15  
 Ala Phe Arg Val Val Glu Gly Gln Gly Trp Gln Ala Phe Lys Asn Asp  
 20 25 30  
 Ala Thr Glu Ile Ile Arg Glu Leu Gly Glu Tyr Pro Glu Pro Pro Pro  
 35 40 45  
 Glu Leu Glu Asn Asn Lys Thr Met Asn Arg Ala Glu Asn Gly Gly Arg  
 50 55 60  
 Pro Pro His His Pro Phe Glu Thr Lys Asp Val Ser Glu Tyr Ser Cys  
 65 70 75 80  
 Arg Glu Leu His Phe Thr Arg Tyr Val Thr Asp Gly Pro Cys Arg Ser  
 85 90 95  
 Ala Lys Pro Val Thr Glu Leu Val Cys Ser Gly Gln Cys Gly Pro Ala  
 100 105 110  
 Arg Leu Leu Pro Asn Ala Ile Gly Arg Gly Lys Trp Trp Arg Pro Ser  
 115 120 125  
 Gly Pro Asp Phe Arg Cys Ile Pro Asp Arg Tyr Arg Ala Gln Arg Val  
 130 135 140  
 Gln Leu Leu Cys Pro Gly Gly Glu Ala Pro Arg Ala Arg Lys Val Arg  
 145 150 155 160  
 Leu Val Ala Ser Cys Lys Cys Lys Arg Leu Thr Arg Phe His Asn Gln  
 165 170 175  
 Ser Glu Leu Lys Asp Phe Gly Thr Glu Ala Ala Arg Pro Gln Lys Gly  
 180 185 190  
 Arg Lys Pro Arg Pro Arg Ala Arg Ser Ala Lys Ala Asn Gln Ala Glu  
 195 200 205  
 Leu Glu Asn Ala Tyr  
 210

<210> 63  
 <211> 642  
 <212> DNA  
 <213> Cercopithecus pygerythrus

<400> 63  
 atgcagctcc cactggccct gtgtcttgct tgccctgctgg tacacgcagc cttccgtgta 60  
 gtggagggcc aggggtggca ggccttcaag aatgatgcca cggaaatcat ccccgagctc 120  
 ggagagtacc ccgagcctcc accggagctg gagaacaaca agaccatgaa ccgggcggag 180  
 aatggagggc ggcctcccca ccaccccttt gagaccaaag acgtgtccga gtacagctgc 240  
 cgagagctgc acttcacccg ctacgtgacc gatgggcccgt gccgcagcgc caagccagtc 300  
 accgagttgg tgtgctccgg ccagtgcggc ccggcacgcc tgctgcccga cgccatcggc 360  
 cgcggaagt ggtggcgccc gagtgggccc gacttccgct gcatccccga ccgctaccgc 420  
 gcgcagcgtg tgcagctgct gtgtcccggg ggtgccgcgc cgcgcgcgcg caaggtgcgc 480  
 ctggtggcct cgtgcaagtg caagcgccct acccgcttcc acaaccagtc ggagctcaag 540  
 gacttcggtc ccgaggccgc tcggccgcag aagggccgga agccgcggcc ccgcgcccgg 600  
 ggggccaag ccaatcaggc cgagctggag aacgcctact ag 642

<210> 64  
 <211> 213  
 <212> PRT  
 <213> Cercopithecus pygerythrus

20

&lt;400&gt; 64

```

Met Gln Leu Pro Leu Ala Leu Cys Leu Val Cys Leu Leu Val His Ala
 1          5          10          15
Ala Phe Arg Val Val Glu Gly Gln Gly Trp Gln Ala Phe Lys Asn Asp
 20          25          30
Ala Thr Glu Ile Ile Pro Glu Leu Gly Glu Tyr Pro Glu Pro Pro Pro
 35          40          45
Glu Leu Glu Asn Asn Lys Thr Met Asn Arg Ala Glu Asn Gly Gly Arg
 50          55          60
Pro Pro His His Pro Phe Glu Thr Lys Asp Val Ser Glu Tyr Ser Cys
 65          70          75          80
Arg Glu Leu His Phe Thr Arg Tyr Val Thr Asp Gly Pro Cys Arg Ser
 85          90          95
Ala Lys Pro Val Thr Glu Leu Val Cys Ser Gly Gln Cys Gly Pro Ala
100          105          110
Arg Leu Leu Pro Asn Ala Ile Gly Arg Gly Lys Trp Trp Arg Pro Ser
115          120          125
Gly Pro Asp Phe Arg Cys Ile Pro Asp Arg Tyr Arg Ala Gln Arg Val
130          135          140
Gln Leu Leu Cys Pro Gly Gly Ala Ala Pro Arg Ala Arg Lys Val Arg
145          150          155          160
Leu Val Ala Ser Cys Lys Cys Lys Arg Leu Thr Arg Phe His Asn Gln
165          170          175
Ser Glu Leu Lys Asp Phe Gly Pro Glu Ala Ala Arg Pro Gln Lys Gly
180          185          190
Arg Lys Pro Arg Pro Arg Ala Arg Gly Ala Lys Ala Asn Gln Ala Glu
195          200          205
Leu Glu Asn Ala Tyr
210

```

&lt;210&gt; 65

&lt;211&gt; 638

&lt;212&gt; DNA

&lt;213&gt; Mus musculus

&lt;400&gt; 65

```

atgcagccct cactagcccc gtgcctcatc tgcctacttg tgcacgctgc cttctgtgct 60
gtggagggcc aggggtggca agccttcagg aatgatgcc aagaggtcat cccagggctt 120
ggagagtacc ccgagcctcc tcctgagaac aaccagacca tgaaccgggc ggagaatgga 180
ggcagacctc cccaccatcc ctatgacgcc aaaggtgtgt ccgagtacag ctgccgcgag 240
ctgcactaca cccgcttcct gacagacggc ccattgccga gcgccaagcc ggtcaccgag 300
ttggtgtgct ccggccagtg cggccccgcg cggtgtgtgc ccaacgccat cgggcgcgtg 360
aagtgggtggc gcccgaacgg accggatttc cgctgcatcc cggatcgcta ccgcgcgcag 420
cgggtgcagc tgctgtgccc cgggggcgcg gcgcgcgcgt cgcgcaaggt gcgtctggtg 480
gcctcgtgca agtgcaagcg cctcaccgcg ttccacaacc agtcggagct caaggacttc 540
gggcgcgaga ccgcgcggcc gcagaagggt cgcaagccgc ggcccggcgc ccggggagcc 600
aaagccaacc aggcggagct ggagaacgcc tactagag 638

```

&lt;210&gt; 66

&lt;211&gt; 211

&lt;212&gt; PRT

&lt;213&gt; Mus musculus

&lt;400&gt; 66

```

Met Gln Pro Ser Leu Ala Pro Cys Leu Ile Cys Leu Leu Val His Ala
 1          5          10          15
Ala Phe Cys Ala Val Glu Gly Gln Gly Trp Gln Ala Phe Arg Asn Asp

```

```
<210> 67
<211> 674
<212> DNA
<213> Rattus norvegicus
```

<400> 67						
gaggaccgag	tgbcccttct	ccttcttgga	ccatgcagct	ctcactagcc	ccttgccttg	60
cctgcctgct	tgtacatgca	gccttcggtg	ctgtggagag	ccaggggtgg	caagccttca	120
agaatgatgc	cacagaaatc	atcccgggac	tcagagagta	cccagagcct	cctcaggaac	180
tagaagaaca	ccagaccatg	aaccggcgct	agaacggagg	cgagaccccc	caccattcct	240
atgacaccaa	agacgtgtcc	gagtagcagct	gccgcgagct	gcactacacc	cgcttcgtga	300
ccgacggccc	gtgccgcagt	gccaaagccgg	tcaccgagtt	gggtgtgctcg	ggccagtgcg	360
gccccgcgcg	gctgctgccc	aacgccatcg	ggcgcgtgaa	gtggtggcgc	ccgaacggac	420
ccgacttccg	ctgcatcccg	gatcgctacc	gcgcgcagcg	gggtgcagctg	ctgtgccccg	480
gcggcgcggc	gccgcgctcg	cgcaaggtgc	gtctggtggc	ctcgtgcaag	tgcaagcgcc	540
tcacccgcct	ccacaaccag	tcggagctca	aggacttcgg	acctgagacc	gcgcggccgc	600
agaagggtcg	caagccgcgg	ccccgcgccc	ggggagccaa	agccaaccag	gcggagctgg	660
agaacgccta	ctag					674

```
<210> 68
<211> 213
<212> PRT
<213> Rattus norvegicus
```

<400> 68  
Met Gln Leu Ser Leu Ala Pro Cys Leu Ala Cys Leu Leu Val His Ala  
1 5 10 15  
Ala Phe Val Ala Val Glu Ser Gln Gly Trp Gln Ala Phe Lys Asn Asp  
20 25 30  
Ala Thr Glu Ile Ile Pro Gly Leu Arg Glu Tyr Pro Glu Pro Pro Gln  
35 40 45

22

Glu Leu Glu Asn Asn Gln Thr Met Asn Arg Ala Glu Asn Gly Gly Arg  
 50 55 60  
 Pro Pro His His Pro Tyr Asp Thr Lys Asp Val Ser Glu Tyr Ser Cys  
 65 70 75 80  
 Arg Glu Leu His Tyr Thr Arg Phe Val Thr Asp Gly Pro Cys Arg Ser  
 85 90 95  
 Ala Lys Pro Val Thr Glu Leu Val Cys Ser Gly Gln Cys Gly Pro Ala  
 100 105 110  
 Arg Leu Leu Pro Asn Ala Ile Gly Arg Val Lys Trp Trp Arg Pro Asn  
 115 120 125  
 Gly Pro Asp Phe Arg Cys Ile Pro Asp Arg Tyr Arg Ala Gln Arg Val  
 130 135 140  
 Gln Leu Leu Cys Pro Gly Gly Ala Ala Pro Arg Ser Arg Lys Val Arg  
 145 150 155 160  
 Leu Val Ala Ser Cys Lys Cys Lys Arg Leu Thr Arg Phe His Asn Gln  
 165 170 175  
 Ser Glu Leu Lys Asp Phe Gly Pro Glu Thr Ala Arg Pro Gln Lys Gly  
 180 185 190  
 Arg Lys Pro Arg Pro Arg Ala Arg Gly Ala Lys Ala Asn Gln Ala Glu  
 195 200 205  
 Leu Glu Asn Ala Tyr  
 210

<210> 69  
 <211> 532  
 <212> DNA  
 <213> Bos torus

<400> 69  
 agaatgatgc cacagaaatc atccccgagc tgggcgagta ccccgagcct ctgccagagc 60  
 tgaacaacaa gac'catgaac cggg'cgagaga acggagggag acctccccac cacc'cctttg 120  
 agaccaaaga cgcctccgag tacagctgcc gggagctgca cttc'acccgc tacgtgaccg 180  
 atgggcccgtg ccgcagcgcc aagccgggtca ccgagctggt gtgctcgggc cagtgcggcc 240  
 cggcgcgccct gctgccc'aaac gccatcggcc gcggcaagt'g gtggcgccca agcgggcccc 300  
 acttccgctg catccccgac cgctaccgcg cgcagcgggt gcagctgttg t'gtcctggcg 360  
 gcgcggcgccc gcgcgcgcgc aaggtgcgcg tgg'tggcctc gtgcaagtgc aagcgcctca 420  
 ctgcgttcca caaccagtcc gagctcaagg acttcggggc cgaggccgcg cggccgcaaa 480  
 cgggcccggaa gctgcggccc cgcgcccggg gcaccaaagc cagccggg'cc ga 532

<210> 70  
 <211> 176  
 <212> PRT  
 <213> Bos torus

<400> 70  
 Asn Asp Ala Thr Glu Ile Ile Pro Glu Leu Gly Glu Tyr Pro Glu Pro  
 1 5 10 15  
 Leu Pro Glu Leu Asn Asn Lys Thr Met Asn Arg Ala Glu Asn Gly Gly  
 20 25 30  
 Arg Pro Pro His His Pro Phe Glu Thr Lys Asp Ala Ser Glu Tyr Ser  
 35 40 45  
 Cys Arg Glu Leu His Phe Thr Arg Tyr Val Thr Asp Gly Pro Cys Arg  
 50 55 60  
 Ser Ala Lys Pro Val Thr Glu Leu Val Cys Ser Gly Gln Cys Gly Pro  
 65 70 75 80  
 Ala Arg Leu Leu Pro Asn Ala Ile Gly Arg Gly Lys Trp Trp Arg Pro  
 85 90 95

23

```

Ser Gly Pro Asp Phe Arg Cys Ile Pro Asp Arg Tyr Arg Ala Gln Arg
      100      105      110
Val Gln Leu Leu Cys Pro Gly Gly Ala Ala Pro Arg Ala Arg Lys Val
      115      120      125
Arg Leu Val Ala Ser Cys Lys Cys Lys Arg Leu Thr Arg Phe His Asn
      130      135      140
Gln Ser Glu Leu Lys Asp Phe Gly Pro Glu Ala Ala Arg Pro Gln Thr
145      150      155      160
Gly Arg Lys Leu Arg Pro Arg Ala Arg Gly Thr Lys Ala Ser Arg Ala
      165      170      175

```

&lt;210&gt; 71

&lt;211&gt; 532

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 71

```

Met Thr Gln Leu Tyr Ile Tyr Ile Arg Leu Leu Gly Ala Tyr Leu Phe
 1      5      10      15
Ile Ile Ser Arg Val Gln Gly Gln Asn Leu Asp Ser Met Leu His Gly
      20      25      30
Thr Gly Met Lys Ser Asp Ser Asp Gln Lys Lys Ser Glu Asn Gly Val
      35      40      45
Thr Leu Ala Pro Glu Asp Thr Leu Pro Phe Leu Lys Cys Tyr Cys Ser
      50      55      60
Gly His Cys Pro Asp Asp Ala Ile Asn Asn Thr Cys Ile Thr Asn Gly
65      70      75      80
His Cys Phe Ala Ile Glu Glu Asp Asp Gln Gly Glu Thr Thr Leu
      85      90      95
Ala Ser Gly Cys Met Lys Tyr Glu Gly Ser Asp Phe Gln Cys Lys Asp
      100      105      110
Ser Pro Lys Ala Gln Leu Arg Arg Thr Ile Glu Cys Cys Arg Thr Asn
      115      120      125
Leu Cys Asn Gln Tyr Leu Gln Pro Thr Leu Pro Pro Val Val Ile Gly
      130      135      140
Pro Phe Phe Asp Gly Ser Ile Arg Trp Leu Val Leu Leu Ile Ser Met
145      150      155      160
Ala Val Cys Ile Ile Ala Met Ile Ile Phe Ser Ser Cys Phe Cys Tyr
      165      170      175
Lys His Tyr Cys Lys Ser Ile Ser Ser Arg Arg Arg Tyr Asn Arg Asp
      180      185      190
Leu Glu Gln Asp Glu Ala Phe Ile Pro Val Gly Glu Ser Leu Lys Asp
      195      200      205
Leu Ile Asp Gln Ser Gln Ser Ser Gly Ser Gly Ser Gly Leu Pro Leu
      210      215      220
Leu Val Gln Arg Thr Ile Ala Lys Gln Ile Gln Met Val Arg Gln Val
225      230      235      240
Gly Lys Gly Arg Tyr Gly Glu Val Trp Met Gly Lys Trp Arg Gly Glu
      245      250      255
Lys Val Ala Val Lys Val Phe Phe Thr Thr Glu Glu Ala Ser Trp Phe
      260      265      270
Arg Glu Thr Glu Ile Tyr Gln Thr Val Leu Met Arg His Glu Asn Ile
      275      280      285
Leu Gly Phe Ile Ala Ala Asp Ile Lys Gly Thr Gly Ser Trp Thr Gln
290      295      300
Leu Tyr Leu Ile Thr Asp Tyr His Glu Asn Gly Ser Leu Tyr Asp Phe
305      310      315      320

```

```
<210> 72
<211> 502
<212> PRT
<213> Homo sapiens
```

<400>	72															
Met	Leu	Leu	Arg	Ser	Ala	Gly	Lys	Leu	Asn	Val	Gly	Thr	Lys	Lys	Glu	
1				5					10					15		
Asp	Gly	Glu	Ser	Thr	Ala	Pro	Thr	Pro	Arg	Pro	Lys	Val	Leu	Arg	Cys	
			20					25					30			
Lys	Cys	His	His	His	Cys	Pro	Glu	Asp	Ser	Val	Asn	Asn	Ile	Cys	Ser	
		35					40					45				
Thr	Asp	Gly	Tyr	Cys	Phe	Thr	Met	Ile	Glu	Glu	Asp	Asp	Ser	Gly	Leu	
	50					55					60					
Pro	Val	Val	Thr	Ser	Gly	Cys	Leu	Gly	Leu	Glu	Gly	Ser	Asp	Phe	Gln	
65					70					75					80	
Cys	Arg	Asp	Thr	Pro	Ile	Pro	His	Gln	Arg	Arg	Ser	Ile	Glu	Cys	Cys	
				85					90					95		
Thr	Glu	Arg	Asn	Glu	Cys	Asn	Lys	Asp	Leu	His	Pro	Thr	Leu	Pro	Pro	
			100					105					110			
Leu	Lys	Asn	Arg	Asp	Phe	Val	Asp	Gly	Pro	Ile	His	His	Arg	Ala	Leu	
		115					120					125				
Leu	Ile	Ser	Val	Thr	Val	Cys	Ser	Leu	Leu	Leu	Val	Leu	Ile	Ile	Leu	
	130					135					140					
Phe	Cys	Tyr	Phe	Arg	Tyr	Lys	Arg	Gln	Glu	Thr	Arg	Pro	Arg	Tyr	Ser	
145					150					155					160	
Ile	Gly	Leu	Glu	Gln	Asp	Glu	Thr	Tyr	Ile	Pro	Pro	Gly	Glu	Ser	Leu	
				165					170					175		



25

Arg Asp Leu Ile Glu Gln Ser Gln Ser Ser Gly Ser Gly Ser Gly Leu  
 180 185 190  
 Pro Leu Leu Val Gln Arg Thr Ile Ala Lys Gln Ile Gln Met Val Lys  
 195 200 205  
 Gln Ile Gly Lys Gly Arg Tyr Gly Glu Val Trp Met Gly Lys Trp Arg  
 210 215 220  
 Gly Glu Lys Val Ala Val Lys Val Phe Phe Thr Thr Glu Glu Ala Ser  
 225 230 235 240  
 Trp Phe Arg Glu Thr Glu Ile Tyr Gln Thr Val Leu Met Arg His Glu  
 245 250 255  
 Asn Ile Leu Gly Phe Ile Ala Ala Asp Ile Lys Gly Thr Gly Ser Trp  
 260 265 270  
 Thr Gln Leu Tyr Leu Ile Thr Asp Tyr His Glu Asn Gly Ser Leu Tyr  
 275 280 285  
 Asp Tyr Leu Lys Ser Thr Thr Leu Asp Ala Lys Ser Met Leu Lys Leu  
 290 295 300  
 Ala Tyr Ser Ser Val Ser Gly Leu Cys His Leu His Thr Glu Ile Phe  
 305 310 315 320  
 Ser Thr Gln Gly Lys Pro Ala Ile Ala His Arg Asp Leu Lys Ser Lys  
 325 330 335  
 Asn Ile Leu Val Lys Lys Asn Gly Thr Cys Cys Ile Ala Asp Leu Gly  
 340 345 350  
 Leu Ala Val Lys Phe Ile Ser Asp Thr Asn Glu Val Asp Ile Pro Pro  
 355 360 365  
 Asn Thr Arg Val Gly Thr Lys Arg Tyr Met Pro Pro Glu Val Leu Asp  
 370 375 380  
 Glu Ser Leu Asn Arg Asn His Phe Gln Ser Tyr Ile Met Ala Asp Met  
 385 390 395 400  
 Tyr Ser Phe Gly Leu Ile Leu Trp Glu Val Ala Arg Arg Cys Val Ser  
 405 410 415  
 Gly Gly Ile Val Glu Glu Tyr Gln Leu Pro Tyr His Asp Leu Val Pro  
 420 425 430  
 Ser Asp Pro Ser Tyr Glu Asp Met Arg Glu Ile Val Cys Ile Lys Lys  
 435 440 445  
 Leu Arg Pro Ser Phe Pro Asn Arg Trp Ser Ser Asp Glu Cys Leu Arg  
 450 455 460  
 Gln Met Gly Lys Leu Met Thr Glu Cys Trp Ala His Asn Pro Ala Ser  
 465 470 475 480  
 Arg Leu Thr Ala Leu Arg Val Lys Lys Thr Leu Ala Lys Met Ser Glu  
 485 490 495  
 Ser Gln Asp Ile Lys Leu  
 500

&lt;210&gt; 73

&lt;211&gt; 502

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 73

Met Leu Leu Arg Ser Ala Gly Lys Leu Asn Val Gly Thr Lys Lys Glu  
 1 5 10 15  
 Asp Gly Glu Ser Thr Ala Pro Thr Pro Arg Pro Lys Val Leu Arg Cys  
 20 25 30  
 Lys Cys His His His Cys Pro Glu Asp Ser Val Asn Asn Ile Cys Ser  
 35 40 45  
 Thr Asp Gly Tyr Cys Phe Thr Met Ile Glu Glu Asp Asp Ser Gly Leu  
 50 55 60

Pro	Val	Val	Thr	Ser	Gly	Cys	Leu	Gly	Leu	Glu	Gly	Ser	Asp	Phe	Gln	65	70	75	80
Cys	Arg	Asp	Thr	Pro	Ile	Pro	His	Gln	Arg	Arg	Ser	Ile	Glu	Cys	Cys	85	90	95	
Thr	Glu	Arg	Asn	Glu	Cys	Asn	Lys	Asp	Leu	His	Pro	Thr	Leu	Pro	Pro	100	105	110	
Leu	Lys	Asn	Arg	Asp	Phe	Val	Asp	Gly	Pro	Ile	His	His	Arg	Ala	Leu	115	120	125	
Leu	Ile	Ser	Val	Thr	Val	Cys	Ser	Leu	Leu	Leu	Val	Leu	Ile	Ile	Leu	130	135	140	
Phe	Cys	Tyr	Phe	Arg	Tyr	Lys	Arg	Gln	Glu	Thr	Arg	Pro	Arg	Tyr	Ser	145	150	155	160
Ile	Gly	Leu	Glu	Gln	Asp	Glu	Thr	Tyr	Ile	Pro	Pro	Gly	Glu	Ser	Leu	165	170	175	
Arg	Asp	Leu	Ile	Glu	Gln	Ser	Gln	Ser	Ser	Gly	Ser	Gly	Ser	Gly	Leu	180	185	190	
Pro	Leu	Leu	Val	Gln	Arg	Thr	Ile	Ala	Lys	Gln	Ile	Gln	Met	Val	Lys	195	200	205	
Gln	Ile	Gly	Lys	Gly	Arg	Tyr	Gly	Glu	Val	Trp	Met	Gly	Lys	Trp	Arg	210	215	220	
Gly	Glu	Lys	Val	Ala	Val	Lys	Val	Phe	Phe	Thr	Thr	Glu	Glu	Ala	Ser	225	230	235	240
Trp	Phe	Arg	Glu	Thr	Glu	Ile	Tyr	Gln	Thr	Val	Leu	Met	Arg	His	Glu	245	250	255	
Asn	Ile	Leu	Gly	Phe	Ile	Ala	Ala	Asp	Ile	Lys	Gly	Thr	Gly	Ser	Trp	260	265	270	
Thr	Gln	Leu	Tyr	Leu	Ile	Thr	Asp	Tyr	His	Glu	Asn	Gly	Ser	Leu	Tyr	275	280	285	
Asp	Tyr	Leu	Lys	Ser	Thr	Thr	Leu	Asp	Ala	Lys	Ser	Met	Leu	Lys	Leu	290	295	300	
Ala	Tyr	Ser	Ser	Val	Ser	Gly	Leu	Cys	His	Leu	His	Thr	Glu	Ile	Phe	305	310	315	320
Ser	Thr	Gln	Gly	Lys	Pro	Ala	Ile	Ala	His	Arg	Asp	Leu	Lys	Ser	Lys	325	330	335	
Asn	Ile	Leu	Val	Lys	Lys	Asn	Gly	Thr	Cys	Cys	Ile	Ala	Asp	Leu	Gly	340	345	350	
Leu	Ala	Val	Lys	Phe	Ile	Ser	Asp	Thr	Asn	Glu	Val	Asp	Ile	Pro	Pro	355	360	365	
Asn	Thr	Arg	Val	Gly	Thr	Lys	Arg	Tyr	Met	Pro	Pro	Glu	Val	Leu	Asp	370	375	380	
Glu	Ser	Leu	Asn	Arg	Asn	His	Phe	Gln	Ser	Tyr	Ile	Met	Ala	Asp	Met	385	390	395	400
Tyr	Ser	Phe	Gly	Leu	Ile	Leu	Trp	Glu	Val	Ala	Arg	Arg	Cys	Val	Ser	405	410	415	
Gly	Gly	Ile	Val	Glu	Glu	Tyr	Gln	Leu	Pro	Tyr	His	Asp	Leu	Val	Pro	420	425	430	
Ser	Asp	Pro	Ser	Tyr	Glu	Asp	Met	Arg	Glu	Ile	Val	Cys	Ile	Lys	Lys	435	440	445	
Leu	Arg	Pro	Ser	Phe	Pro	Asn	Arg	Trp	Ser	Ser	Asp	Glu	Cys	Leu	Arg	450	455	460	
Gln	Met	Gly	Lys	Leu	Met	Thr	Glu	Cys	Trp	Ala	His	Asn	Pro	Ala	Ser	465	470	475	480
Arg	Leu	Thr	Ala	Leu	Arg	Val	Lys	Lys	Thr	Leu	Ala	Lys	Met	Ser	Glu	485	490	495	
Ser	Gln	Asp	Ile	Lys	Leu											500			

<210> 74  
 <211> 532  
 <212> PRT  
 <213> Rattus sp.

<400> 74

Met	Thr	Gln	Leu	Tyr	Thr	Tyr	Ile	Arg	Leu	Leu	Gly	Ala	Cys	Leu	Phe	1	5	10	15
Ile	Ile	Ser	His	Val	Gln	Gly	Gln	Asn	Leu	Asp	Ser	Met	Leu	His	Gly	20	25	30	
Thr	Gly	Met	Lys	Ser	Asp	Val	Asp	Gln	Lys	Lys	Pro	Glu	Asn	Gly	Val	35	40	45	
Thr	Leu	Ala	Pro	Glu	Asp	Thr	Leu	Pro	Phe	Leu	Lys	Cys	Tyr	Cys	Ser	50	55	60	
Gly	His	Cys	Pro	Asp	Asp	Ala	Ile	Asn	Asn	Thr	Cys	Ile	Thr	Asn	Gly	65	70	75	80
His	Cys	Phe	Ala	Ile	Ile	Glu	Glu	Asp	Asp	Gln	Gly	Glu	Thr	Thr	Leu	85	90	95	
Thr	Ser	Gly	Cys	Met	Lys	Tyr	Glu	Gly	Ser	Asp	Phe	Gln	Cys	Lys	Asp	100	105	110	
Ser	Pro	Lys	Ala	Gln	Leu	Arg	Arg	Thr	Ile	Glu	Cys	Cys	Arg	Thr	Asn	115	120	125	
Leu	Cys	Asn	Gln	Tyr	Leu	Gln	Pro	Thr	Leu	Pro	Pro	Val	Val	Ile	Gly	130	135	140	
Pro	Phe	Phe	Asp	Gly	Ser	Val	Arg	Trp	Leu	Ala	Val	Leu	Ile	Ser	Met	145	150	155	160
Ala	Val	Cys	Ile	Val	Ala	Met	Ile	Val	Phe	Ser	Ser	Cys	Phe	Cys	Tyr	165	170	175	
Lys	His	Tyr	Cys	Lys	Ser	Ile	Ser	Ser	Arg	Gly	Arg	Tyr	Asn	Arg	Asp	180	185	190	
Leu	Glu	Gln	Asp	Glu	Ala	Phe	Ile	Pro	Val	Gly	Glu	Ser	Leu	Lys	Asp	195	200	205	
Leu	Ile	Asp	Gln	Ser	Gln	Ser	Ser	Gly	Ser	Gly	Ser	Gly	Leu	Pro	Leu	210	215	220	
Leu	Val	Gln	Arg	Thr	Ile	Ala	Lys	Gln	Ile	Gln	Met	Val	Arg	Gln	Val	225	230	235	240
Gly	Lys	Gly	Arg	Tyr	Gly	Glu	Val	Trp	Met	Gly	Lys	Trp	Arg	Gly	Glu	245	250	255	
Lys	Val	Ala	Val	Lys	Val	Phe	Phe	Thr	Thr	Glu	Glu	Ala	Ser	Trp	Phe	260	265	270	
Arg	Glu	Thr	Glu	Ile	Tyr	Gln	Thr	Val	Leu	Met	Arg	His	Glu	Asn	Ile	275	280	285	
Leu	Gly	Phe	Ile	Ala	Ala	Asp	Ile	Lys	Gly	Thr	Gly	Ser	Trp	Thr	Gln	290	295	300	
Leu	Tyr	Leu	Ile	Thr	Asp	Tyr	His	Glu	Asn	Gly	Ser	Leu	Tyr	Asp	Phe	305	310	315	320
Leu	Lys	Cys	Ala	Thr	Leu	Asp	Thr	Arg	Ala	Leu	Leu	Lys	Leu	Ala	Tyr	325	330	335	
Ser	Ala	Ala	Cys	Gly	Leu	Cys	His	Leu	His	Thr	Glu	Ile	Tyr	Gly	Thr	340	345	350	
Gln	Gly	Lys	Pro	Ala	Ile	Ala	His	Arg	Asp	Leu	Lys	Ser	Lys	Asn	Ile	355	360	365	
Leu	Ile	Lys	Lys	Asn	Gly	Ser	Cys	Cys	Ile	Ala	Asp	Leu	Gly	Leu	Ala	370	375	380	
Val	Lys	Phe	Asn	Ser	Asp	Thr	Asn	Glu	Val	Asp	Ile	Pro	Leu	Asn	Thr	385	390	395	400
Arg	Val	Gly	Thr	Arg	Arg	Tyr	Met	Ala	Pro	Glu	Val	Leu	Asp	Glu	Ser	405	410	415	

28

Leu Ser Lys Asn His Phe Gln Pro Tyr Ile Met Ala Asp Ile Tyr Ser  
                   420                  425                  430  
 Phe Gly Leu Ile Ile Trp Glu Met Ala Arg Arg Cys Ile Thr Gly Gly  
                   435                  440                  445  
 Ile Val Glu Glu Tyr Gln Leu Pro Tyr Tyr Asn Met Val Pro Ser Asp  
                   450                  455                  460  
 Pro Ser Tyr Glu Asp Met Arg Glu Val Val Cys Val Lys Arg Leu Arg  
                   465                  470                  475                  480  
 Pro Ile Val Ser Asn Arg Trp Asn Ser Asp Glu Cys Leu Arg Ala Val  
                   485                  490                  495  
 Leu Lys Leu Met Ser Glu Cys Trp Ala His Asn Pro Ala Ser Arg Leu  
                   500                  505                  510  
 Thr Ala Leu Arg Ile Lys Lys Thr Leu Ala Lys Met Val Glu Ser Gln  
                   515                  520                  525  
 Asp Val Lys Ile  
                   530

<210> 75  
 <211> 532  
 <212> PRT  
 <213> Rattus norvegicus

<400> 75  
 Met Thr Gln Leu Tyr Thr Tyr Ile Arg Leu Leu Gly Ala Cys Leu Phe  
   1                  5                  10                  15  
 Ile Ile Ser His Val Gln Gly Gln Asn Leu Asp Ser Met Leu His Gly  
                   20                  25                  30  
 Thr Gly Met Lys Ser Asp Val Asp Gln Lys Lys Pro Glu Asn Gly Val  
                   35                  40                  45  
 Thr Leu Ala Pro Glu Asp Thr Leu Pro Phe Leu Lys Cys Tyr Cys Ser  
                   50                  55                  60  
 Gly His Cys Pro Asp Asp Ala Ile Asn Asn Thr Cys Ile Thr Asn Gly  
   65                  70                  75                  80  
 His Cys Phe Ala Ile Ile Glu Glu Asp Asp Gln Gly Glu Thr Thr Leu  
                   85                  90                  95  
 Thr Ser Gly Cys Met Lys Tyr Glu Gly Ser Asp Phe Gln Cys Lys Asp  
                   100                  105                  110  
 Ser Pro Lys Ala Gln Leu Arg Arg Thr Ile Glu Cys Cys Arg Thr Asn  
                   115                  120                  125  
 Leu Cys Asn Gln Tyr Leu Gln Pro Thr Leu Pro Pro Val Val Ile Gly  
                   130                  135                  140  
 Pro Phe Phe Asp Gly Ser Val Arg Trp Leu Ala Val Leu Ile Ser Met  
   145                  150                  155                  160  
 Ala Val Cys Ile Val Ala Met Ile Val Phe Ser Ser Cys Phe Cys Tyr  
                   165                  170                  175  
 Lys His Tyr Cys Lys Ser Ile Ser Ser Arg Gly Arg Tyr Asn Arg Asp  
                   180                  185                  190  
 Leu Glu Gln Asp Glu Ala Phe Ile Pro Val Gly Glu Ser Leu Lys Asp  
                   195                  200                  205  
 Leu Ile Asp Gln Ser Gln Ser Ser Gly Ser Gly Ser Gly Leu Pro Leu  
                   210                  215                  220  
 Leu Val Gln Arg Thr Ile Ala Lys Gln Ile Gln Met Val Arg Gln Val  
   225                  230                  235                  240  
 Gly Lys Gly Arg Tyr Gly Glu Val Trp Met Gly Lys Trp Arg Gly Glu  
                   245                  250                  255  
 Lys Val Ala Val Lys Val Phe Phe Thr Thr Glu Glu Ala Ser Trp Phe  
                   260                  265                  270

```
<210> 76
<211> 532
<212> PRT
<213> Rattus norvegicus
```

<400> 76															
Met	Thr	Gln	Leu	Tyr	Thr	Tyr	Ile	Arg	Leu	Leu	Gly	Ala	Cys	Leu	Phe
1				5					10					15	
Ile	Ile	Ser	His	Val	Gln	Gly	Gln	Asn	Leu	Asp	Ser	Met	Leu	His	Gly
			20					25					30		
Thr	Gly	Met	Lys	Ser	Asp	Val	Asp	Gln	Lys	Lys	Pro	Glu	Asn	Gly	Val
		35					40					45			
Thr	Leu	Ala	Pro	Glu	Asp	Thr	Leu	Pro	Phe	Leu	Lys	Cys	Tyr	Cys	Ser
	50					55					60				
Gly	His	Cys	Pro	Asp	Asp	Ala	Ile	Asn	Asn	Thr	Cys	Ile	Thr	Asn	Gly
65				70						75				80	
His	Cys	Phe	Ala	Ile	Ile	Glu	Glu	Asp	Asp	Gln	Gly	Glu	Thr	Thr	Leu
			85						90					95	
Thr	Ser	Gly	Cys	Met	Lys	Tyr	Glu	Gly	Ser	Asp	Phe	Gln	Cys	Lys	Asp
			100					105					110		
Ser	Pro	Lys	Ala	Gln	Leu	Arg	Arg	Thr	Ile	Glu	Cys	Cys	Arg	Thr	Asn
		115					120					125			

Leu Cys Asn Gln Tyr Leu Gln Pro Thr Leu Pro Pro Val Val Ile Gly  
 130 135 140  
 Pro Phe Phe Asp Gly Ser Val Arg Trp Leu Ala Val Leu Ile Ser Met  
 145 150 155 160  
 Ala Val Cys Ile Val Ala Met Ile Val Phe Ser Ser Cys Phe Cys Tyr  
 165 170 175  
 Lys His Tyr Cys Lys Ser Ile Ser Ser Arg Gly Arg Tyr Asn Arg Asp  
 180 185 190  
 Leu Glu Gln Asp Glu Ala Phe Ile Pro Val Gly Glu Ser Leu Lys Asp  
 195 200 205  
 Leu Ile Asp Gln Ser Gln Ser Ser Gly Ser Gly Ser Gly Leu Pro Leu  
 210 215 220  
 Leu Val Gln Arg Thr Ile Ala Lys Gln Ile Gln Met Val Arg Gln Val  
 225 230 235 240  
 Gly Lys Gly Arg Tyr Gly Glu Val Trp Met Gly Lys Trp Arg Gly Glu  
 245 250 255  
 Lys Val Ala Val Lys Val Phe Phe Thr Thr Glu Glu Ala Ser Trp Phe  
 260 265 270  
 Arg Glu Thr Glu Ile Tyr Gln Thr Val Leu Met Arg His Glu Asn Ile  
 275 280 285  
 Leu Gly Phe Ile Ala Ala Asp Ile Lys Gly Thr Gly Ser Trp Thr Gln  
 290 295 300  
 Leu Tyr Leu Ile Thr Asp Tyr His Glu Asn Gly Ser Leu Tyr Asp Phe  
 305 310 315 320  
 Leu Lys Cys Ala Thr Leu Asp Thr Arg Ala Leu Leu Lys Leu Ala Tyr  
 325 330 335  
 Ser Ala Ala Cys Gly Leu Cys His Leu His Thr Glu Ile Tyr Gly Thr  
 340 345 350  
 Gln Gly Lys Pro Ala Ile Ala His Arg Asp Leu Lys Ser Lys Asn Ile  
 355 360 365  
 Leu Ile Lys Lys Asn Gly Ser Cys Cys Ile Ala Asp Leu Gly Leu Ala  
 370 375 380  
 Val Lys Phe Asn Ser Asp Thr Asn Glu Val Asp Ile Pro Leu Asn Thr  
 385 390 395 400  
 Arg Val Gly Thr Arg Arg Tyr Met Ala Pro Glu Val Leu Asp Glu Ser  
 405 410 415  
 Leu Ser Lys Asn His Phe Gln Pro Tyr Ile Met Ala Asp Ile Tyr Ser  
 420 425 430  
 Phe Gly Leu Ile Ile Trp Glu Met Ala Arg Arg Cys Ile Thr Gly Gly  
 435 440 445  
 Ile Val Glu Glu Tyr Gln Leu Pro Tyr Tyr Asn Met Val Pro Ser Asp  
 450 455 460  
 Pro Ser Tyr Glu Asp Met Arg Glu Val Val Cys Val Lys Arg Leu Arg  
 465 470 475 480  
 Pro Ile Val Ser Asn Arg Trp Asn Ser Asp Glu Cys Leu Arg Ala Val  
 485 490 495  
 Leu Lys Leu Met Ser Glu Cys Trp Ala His Asn Pro Ala Ser Arg Leu  
 500 505 510  
 Thr Ala Leu Arg Ile Lys Lys Thr Leu Ala Lys Met Val Glu Ser Gln  
 515 520 525  
 Asp Val Lys Ile  
 530

&lt;210&gt; 77

&lt;211&gt; 502

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 77

```

Met Leu Leu Arg Ser Ala Gly Lys Leu Asn Val Gly Thr Lys Lys Glu
 1      5      10      15
Asp Gly Glu Ser Thr Ala Pro Thr Pro Arg Pro Lys Val Leu Arg Cys
      20      25      30
Lys Cys His His Cys Pro Glu Asp Ser Val Asn Asn Ile Cys Ser
      35      40      45
Thr Asp Gly Tyr Cys Phe Thr Met Ile Glu Glu Asp Asp Ser Gly Leu
      50      55      60
Pro Val Val Thr Ser Gly Cys Leu Gly Leu Glu Gly Ser Asp Phe Gln
      65      70      75      80
Cys Arg Asp Thr Pro Ile Pro His Gln Arg Arg Ser Ile Glu Cys Cys
      85      90      95
Thr Glu Arg Asn Glu Cys Asn Lys Asp Leu His Pro Thr Leu Pro Pro
      100      105      110
Leu Lys Asn Arg Asp Phe Val Asp Gly Pro Ile His His Arg Ala Leu
      115      120      125
Leu Ile Ser Val Thr Val Cys Ser Leu Leu Leu Val Leu Ile Ile Leu
      130      135      140
Phe Cys Tyr Phe Arg Tyr Lys Arg Gln Glu Thr Arg Pro Arg Tyr Ser
      145      150      155      160
Ile Gly Leu Glu Gln Asp Glu Thr Tyr Ile Pro Pro Gly Glu Ser Leu
      165      170      175
Arg Asp Leu Ile Glu Gln Ser Gln Ser Ser Gly Ser Gly Ser Gly Leu
      180      185      190
Pro Leu Leu Val Gln Arg Thr Ile Ala Lys Gln Ile Gln Met Val Lys
      195      200      205
Gln Ile Gly Lys Gly Arg Tyr Gly Glu Val Trp Met Gly Lys Trp Arg
      210      215      220
Gly Glu Lys Val Ala Val Lys Val Phe Phe Thr Thr Glu Glu Ala Ser
      225      230      235      240
Trp Phe Arg Glu Thr Glu Ile Tyr Gln Thr Val Leu Met Arg His Glu
      245      250      255
Asn Ile Leu Gly Phe Ile Ala Ala Asp Ile Lys Gly Thr Gly Ser Trp
      260      265      270
Thr Gln Leu Tyr Leu Ile Thr Asp Tyr His Glu Asn Gly Ser Leu Tyr
      275      280      285
Asp Tyr Leu Lys Ser Thr Thr Leu Asp Ala Lys Ser Met Leu Lys Leu
      290      295      300
Ala Tyr Ser Ser Val Ser Gly Leu Cys His Leu His Thr Glu Ile Phe
      305      310      315      320
Ser Thr Gln Gly Lys Pro Ala Ile Ala His Arg Asp Leu Lys Ser Lys
      325      330      335
Asn Ile Leu Val Lys Lys Asn Gly Thr Cys Cys Ile Ala Asp Leu Gly
      340      345      350
Leu Ala Val Lys Phe Ile Ser Asp Thr Asn Glu Val Asp Ile Pro Pro
      355      360      365
Asn Thr Arg Val Gly Thr Lys Arg Tyr Met Pro Pro Glu Val Leu Asp
      370      375      380
Glu Ser Leu Asn Arg Asn His Phe Gln Ser Tyr Ile Met Ala Asp Met
      385      390      395      400
Tyr Ser Phe Gly Leu Ile Leu Trp Glu Val Ala Arg Arg Cys Val Ser
      405      410      415
Gly Gly Ile Val Glu Glu Tyr Gln Leu Pro Tyr His Asp Leu Val Pro
      420      425      430
Ser Asp Pro Ser Tyr Glu Asp Met Arg Glu Ile Val Cys Ile Lys Lys
      435      440      445

```

32

Leu Arg Pro Ser Phe Pro Asn Arg Trp Ser Ser Asp Glu Cys Leu Arg  
 450 455 460  
 Gln Met Gly Lys Leu Met Thr Glu Cys Trp Ala His Asn Pro Ala Ser  
 465 470 475 480  
 Arg Leu Thr Ala Leu Arg Val Lys Lys Thr Leu Ala Lys Met Ser Glu  
 485 490 495  
 Ser Gln Asp Ile Lys Leu  
 500

<210> 78  
 <211> 502  
 <212> PRT  
 <213> Homo sapiens

<400> 78  
 Met Leu Leu Arg Ser Ala Gly Lys Leu Asn Val Gly Thr Lys Lys Glu  
 1 5 10 15  
 Asp Gly Glu Ser Thr Ala Pro Thr Pro Arg Pro Lys Val Leu Arg Cys  
 20 25 30  
 Lys Cys His His Cys Pro Glu Asp Ser Val Asn Asn Ile Cys Ser  
 35 40 45  
 Thr Asp Gly Tyr Cys Phe Thr Met Ile Glu Glu Asp Asp Ser Gly Leu  
 50 55 60  
 Pro Val Val Thr Ser Gly Cys Leu Gly Leu Glu Gly Ser Asp Phe Gln  
 65 70 75 80  
 Cys Arg Asp Thr Pro Ile Pro His Gln Arg Arg Ser Ile Glu Cys Cys  
 85 90 95  
 Thr Glu Arg Asn Glu Cys Asn Lys Asp Leu His Pro Thr Leu Pro Pro  
 100 105 110  
 Leu Lys Asn Arg Asp Phe Val Asp Gly Pro Ile His His Arg Ala Leu  
 115 120 125  
 Leu Ile Ser Val Thr Val Cys Ser Leu Leu Leu Val Leu Ile Ile Leu  
 130 135 140  
 Phe Cys Tyr Phe Arg Tyr Lys Arg Gln Glu Thr Arg Pro Arg Tyr Ser  
 145 150 155 160  
 Ile Gly Leu Glu Gln Asp Glu Thr Tyr Ile Pro Pro Gly Glu Ser Leu  
 165 170 175  
 Arg Asp Leu Ile Glu Gln Ser Gln Ser Ser Gly Ser Gly Ser Gly Leu  
 180 185 190  
 Pro Leu Leu Val Gln Arg Thr Ile Ala Lys Gln Ile Gln Met Val Lys  
 195 200 205  
 Gln Ile Gly Lys Gly Arg Tyr Gly Glu Val Trp Met Gly Lys Trp Arg  
 210 215 220  
 Gly Glu Lys Val Ala Val Lys Val Phe Phe Thr Thr Glu Glu Ala Ser  
 225 230 235 240  
 Trp Phe Arg Glu Thr Glu Ile Tyr Gln Thr Val Leu Met Arg His Glu  
 245 250 255  
 Asn Ile Leu Gly Phe Ile Ala Ala Asp Ile Lys Gly Thr Gly Ser Trp  
 260 265 270  
 Thr Gln Leu Tyr Leu Ile Thr Asp Tyr His Glu Asn Gly Ser Leu Tyr  
 275 280 285  
 Asp Tyr Leu Lys Ser Thr Thr Leu Asp Ala Lys Ser Met Leu Lys Leu  
 290 295 300  
 Ala Tyr Ser Ser Val Ser Gly Leu Cys His Leu His Thr Glu Ile Phe  
 305 310 315 320  
 Ser Thr Gln Gly Lys Pro Ala Ile Ala His Arg Asp Leu Lys Ser Lys  
 325 330 335



Asn Ile Leu Val Lys Lys Asn Gly Thr Cys Cys Ile Ala Asp Leu Gly  
                   340                  345                  350  
 Leu Ala Val Lys Phe Ile Ser Asp Thr Asn Glu Val Asp Ile Pro Pro  
                   355                  360                  365  
 Asn Thr Arg Val Gly Thr Lys Arg Tyr Met Pro Pro Glu Val Leu Asp  
                   370                  375                  380  
 Glu Ser Leu Asn Arg Asn His Phe Gln Ser Tyr Ile Met Ala Asp Met  
 385                  390                  395                  400  
 Tyr Ser Phe Gly Leu Ile Leu Trp Glu Val Ala Arg Arg Cys Val Ser  
                   405                  410                  415  
 Gly Gly Ile Val Glu Glu Tyr Gln Leu Pro Tyr His Asp Leu Val Pro  
                   420                  425                  430  
 Ser Asp Pro Ser Tyr Glu Asp Met Arg Glu Ile Val Cys Ile Lys Lys  
                   435                  440                  445  
 Leu Arg Pro Ser Phe Pro Asn Arg Trp Ser Ser Asp Glu Cys Leu Arg  
 450                  455                  460  
 Gln Met Gly Lys Leu Met Thr Glu Cys Trp Ala His Asn Pro Ala Ser  
 465                  470                  475                  480  
 Arg Leu Thr Ala Leu Arg Val Lys Lys Thr Leu Ala Lys Met Ser Glu  
                   485                  490                  495  
 Ser Gln Asp Ile Lys Leu  
                   500

<210> 79  
 <211> 532  
 <212> PRT  
 <213> Rattus sp.

<400> 79  
 Met Thr Gln Leu Tyr Thr Tyr Ile Arg Leu Leu Gly Ala Cys Leu Phe  
 1                  5                  10                  15  
 Ile Ile Ser His Val Gln Gly Gln Asn Leu Asp Ser Met Leu His Gly  
                   20                  25                  30  
 Thr Gly Met Lys Ser Asp Val Asp Gln Lys Lys Pro Glu Asn Gly Val  
                   35                  40                  45  
 Thr Leu Ala Pro Glu Asp Thr Leu Pro Phe Leu Lys Cys Tyr Cys Ser  
 50                  55                  60  
 Gly His Cys Pro Asp Asp Ala Ile Asn Asn Thr Cys Ile Thr Asn Gly  
 65                  70                  75                  80  
 His Cys Phe Ala Ile Ile Glu Glu Asp Asp Gln Gly Glu Thr Thr Leu  
                   85                  90                  95  
 Thr Ser Gly Cys Met Lys Tyr Glu Gly Ser Asp Phe Gln Cys Lys Asp  
                   100                  105                  110  
 Ser Pro Lys Ala Gln Leu Arg Arg Thr Ile Glu Cys Cys Arg Thr Asn  
                   115                  120                  125  
 Leu Cys Asn Gln Tyr Leu Gln Pro Thr Leu Pro Pro Val Val Ile Gly  
 130                  135                  140  
 Pro Phe Phe Asp Gly Ser Val Arg Trp Leu Ala Val Leu Ile Ser Met  
 145                  150                  155                  160  
 Ala Val Cys Ile Val Ala Met Ile Val Phe Ser Ser Cys Phe Cys Tyr  
                   165                  170                  175  
 Lys His Tyr Cys Lys Ser Ile Ser Ser Arg Gly Arg Tyr Asn Arg Asp  
                   180                  185                  190  
 Leu Glu Gln Asp Glu Ala Phe Ile Pro Val Gly Glu Ser Leu Lys Asp  
                   195                  200                  205  
 Leu Ile Asp Gln Ser Gln Ser Ser Gly Ser Gly Ser Gly Leu Pro Leu  
 210                  215                  220

Leu Val Gln Arg Thr Ile Ala Lys Gln Ile Gln Met Val Arg Gln Val  
 225 230 235 240  
 Gly Lys Gly Arg Tyr Gly Glu Val Trp Met Gly Lys Trp Arg Gly Glu  
 245 250 255  
 Lys Val Ala Val Lys Val Phe Phe Thr Thr Glu Glu Ala Ser Trp Phe  
 260 265 270  
 Arg Glu Thr Glu Ile Tyr Gln Thr Val Leu Met Arg His Glu Asn Ile  
 275 280 285  
 Leu Gly Phe Ile Ala Ala Asp Ile Lys Gly Thr Gly Ser Trp Thr Gln  
 290 295 300  
 Leu Tyr Leu Ile Thr Asp Tyr His Glu Asn Gly Ser Leu Tyr Asp Phe  
 305 310 315 320  
 Leu Lys Cys Ala Thr Leu Asp Thr Arg Ala Leu Leu Lys Leu Ala Tyr  
 325 330 335  
 Ser Ala Ala Cys Gly Leu Cys His Leu His Thr Glu Ile Tyr Gly Thr  
 340 345 350  
 Gln Gly Lys Pro Ala Ile Ala His Arg Asp Leu Lys Ser Lys Asn Ile  
 355 360 365  
 Leu Ile Lys Lys Asn Gly Ser Cys Cys Ile Ala Asp Leu Gly Leu Ala  
 370 375 380  
 Val Lys Phe Asn Ser Asp Thr Asn Glu Val Asp Ile Pro Leu Asn Thr  
 385 390 395 400  
 Arg Val Gly Thr Arg Arg Tyr Met Ala Pro Glu Val Leu Asp Glu Ser  
 405 410 415  
 Leu Ser Lys Asn His Phe Gln Pro Tyr Ile Met Ala Asp Ile Tyr Ser  
 420 425 430  
 Phe Gly Leu Ile Ile Trp Glu Met Ala Arg Arg Cys Ile Thr Gly Gly  
 435 440 445  
 Ile Val Glu Glu Tyr Gln Leu Pro Tyr Tyr Asn Met Val Pro Ser Asp  
 450 455 460  
 Pro Ser Tyr Glu Asp Met Arg Glu Val Val Cys Val Lys Arg Leu Arg  
 465 470 475 480  
 Pro Ile Val Ser Asn Arg Trp Asn Ser Asp Glu Cys Leu Arg Ala Val  
 485 490 495  
 Leu Lys Leu Met Ser Glu Cys Trp Ala His Asn Pro Ala Ser Arg Leu  
 500 505 510  
 Thr Ala Leu Arg Ile Lys Lys Thr Leu Ala Lys Met Val Glu Ser Gln  
 515 520 525  
 Asp Val Lys Ile  
 530

&lt;210&gt; 80

&lt;211&gt; 530

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 80

Met Thr Ser Ser Leu Gln Arg Pro Trp Arg Val Pro Trp Leu Pro Trp  
 1 5 10 15  
 Thr Ile Leu Leu Val Ser Thr Ala Ala Ser Gln Asn Gln Glu Arg  
 20 25 30  
 Leu Cys Ala Phe Lys Asp Pro Tyr Gln Gln Asp Leu Gly Ile Gly Glu  
 35 40 45  
 Ser Arg Ile Ser His Glu Asn Gly Thr Ile Leu Cys Ser Lys Gly Ser  
 50 55 60  
 Thr Cys Tyr Gly Leu Trp Glu Lys Ser Lys Gly Asp Ile Asn Leu Val  
 65 70 75 80

Lys Gln Gly Cys Trp Ser His Ile Gly Asp Pro Gln Glu Cys His Tyr  
 85 90 95  
 Glu Glu Cys Val Val Thr Thr Thr Pro Pro Ser Ile Gln Asn Gly Thr  
 100 105 110  
 Tyr Arg Phe Cys Cys Cys Ser Thr Asp Leu Cys Asn Val Asn Phe Thr  
 115 120 125  
 Glu Asn Phe Pro Pro Pro Asp Thr Thr Pro Leu Ser Pro Pro His Ser  
 130 135 140  
 Phe Asn Arg Asp Glu Thr Ile Ile Ile Ala Leu Ala Ser Val Ser Val  
 145 150 155 160  
 Leu Ala Val Leu Ile Val Ala Leu Cys Phe Gly Tyr Arg Met Leu Thr  
 165 170 175  
 Gly Asp Arg Lys Gln Gly Leu His Ser Met Asn Met Met Glu Ala Ala  
 180 185 190  
 Ala Ser Glu Pro Ser Leu Asp Leu Asp Asn Leu Lys Leu Leu Glu Leu  
 195 200 205  
 Ile Gly Arg Gly Arg Tyr Gly Ala Val Tyr Lys Gly Ser Leu Asp Glu  
 210 215 220  
 Arg Pro Val Ala Val Lys Val Phe Ser Phe Ala Asn Arg Gln Asn Phe  
 225 230 235 240  
 Ile Asn Glu Lys Asn Ile Tyr Arg Val Pro Leu Met Glu His Asp Asn  
 245 250 255  
 Ile Ala Arg Phe Ile Val Gly Asp Glu Arg Val Thr Ala Asp Gly Arg  
 260 265 270  
 Met Glu Tyr Leu Leu Val Met Glu Tyr Tyr Pro Asn Gly Ser Leu Cys  
 275 280 285  
 Lys Tyr Leu Ser Leu His Thr Ser Asp Trp Val Ser Ser Cys Arg Leu  
 290 295 300  
 Ala His Ser Val Thr Arg Gly Leu Ala Tyr Leu His Thr Glu Leu Pro  
 305 310 315 320  
 Arg Gly Asp His Tyr Lys Pro Ala Ile Ser His Arg Asp Leu Asn Ser  
 325 330 335  
 Arg Asn Val Leu Val Lys Asn Asp Gly Thr Cys Val Ile Ser Asp Phe  
 340 345 350  
 Gly Leu Ser Met Arg Leu Thr Gly Asn Arg Leu Val Arg Pro Gly Glu  
 355 360 365  
 Glu Asp Asn Ala Ala Ile Ser Glu Val Gly Thr Ile Arg Tyr Met Ala  
 370 375 380  
 Pro Glu Val Leu Glu Gly Ala Val Asn Leu Arg Asp Cys Glu Ser Ala  
 385 390 395 400  
 Leu Lys Gln Val Asp Met Tyr Ala Leu Gly Leu Ile Tyr Trp Glu Ile  
 405 410 415  
 Phe Met Arg Cys Thr Asp Leu Phe Pro Gly Glu Ser Val Pro Glu Tyr  
 420 425 430  
 Gln Met Ala Phe Gln Thr Glu Val Gly Asn His Pro Thr Phe Glu Asp  
 435 440 445  
 Met Gln Val Leu Val Ser Arg Glu Lys Gln Arg Pro Lys Phe Pro Glu  
 450 455 460  
 Ala Trp Lys Glu Asn Ser Leu Ala Val Arg Ser Leu Lys Glu Thr Ile  
 465 470 475 480  
 Glu Asp Cys Trp Asp Gln Asp Ala Glu Ala Arg Leu Thr Ala Gln Cys  
 485 490 495  
 Ala Glu Glu Arg Met Ala Glu Leu Met Met Ile Trp Glu Arg Asn Lys  
 500 505 510  
 Ser Val Ser Pro Thr Val Asn Pro Met Ser Thr Ala Met Gln Asn Glu  
 515 520 525  
 Arg Arg  
 530

<210> 81  
 <211> 530  
 <212> PRT  
 <213> Homo sapiens

<400> 81

Met	Thr	Ser	Ser	Leu	Gln	Arg	Pro	Trp	Arg	Val	Pro	Trp	Leu	Pro	Trp
1				5					10				15		
Thr	Ile	Leu	Leu	Val	Ser	Thr	Ala	Ala	Ala	Ser	Gln	Asn	Gln	Glu	Arg
			20					25					30		
Leu	Cys	Ala	Phe	Lys	Asp	Pro	Tyr	Gln	Gln	Asp	Leu	Gly	Ile	Gly	Glu
		35					40					45			
Ser	Arg	Ile	Ser	His	Glu	Asn	Gly	Thr	Ile	Leu	Cys	Ser	Lys	Gly	Ser
	50					55					60				
Thr	Cys	Tyr	Gly	Leu	Trp	Glu	Lys	Ser	Lys	Gly	Asp	Ile	Asn	Leu	Val
65					70					75					80
Lys	Gln	Gly	Cys	Trp	Ser	His	Ile	Gly	Asp	Pro	Gln	Glu	Cys	His	Tyr
			85						90					95	
Glu	Glu	Cys	Val	Val	Thr	Thr	Thr	Pro	Pro	Ser	Ile	Gln	Asn	Gly	Thr
			100					105					110		
Tyr	Arg	Phe	Cys	Cys	Cys	Ser	Thr	Asp	Leu	Cys	Asn	Val	Asn	Phe	Thr
		115					120					125			
Glu	Asn	Phe	Pro	Pro	Pro	Asp	Thr	Thr	Pro	Leu	Ser	Pro	Pro	His	Ser
	130					135					140				
Phe	Asn	Arg	Asp	Glu	Thr	Ile	Ile	Ile	Ala	Leu	Ala	Ser	Val	Ser	Val
145					150					155					160
Leu	Ala	Val	Leu	Ile	Val	Ala	Leu	Cys	Phe	Gly	Tyr	Arg	Met	Leu	Thr
				165					170					175	
Gly	Asp	Arg	Lys	Gln	Gly	Leu	His	Ser	Met	Asn	Met	Met	Glu	Ala	Ala
			180					185					190		
Ala	Ser	Glu	Pro	Ser	Leu	Asp	Leu	Asp	Asn	Leu	Lys	Leu	Leu	Glu	Leu
		195					200					205			
Ile	Gly	Arg	Gly	Arg	Tyr	Gly	Ala	Val	Tyr	Lys	Gly	Ser	Leu	Asp	Glu
	210					215					220				
Arg	Pro	Val	Ala	Val	Lys	Val	Phe	Ser	Phe	Ala	Asn	Arg	Gln	Asn	Phe
225					230					235					240
Ile	Asn	Glu	Lys	Asn	Ile	Tyr	Arg	Val	Pro	Leu	Met	Glu	His	Asp	Asn
				245					250					255	
Ile	Ala	Arg	Phe	Ile	Val	Gly	Asp	Glu	Arg	Val	Thr	Ala	Asp	Gly	Arg
		260					265						270		
Met	Glu	Tyr	Leu	Leu	Val	Met	Glu	Tyr	Tyr	Pro	Asn	Gly	Ser	Leu	Cys
		275					280					285			
Lys	Tyr	Leu	Ser	Leu	His	Thr	Ser	Asp	Trp	Val	Ser	Ser	Cys	Arg	Leu
	290					295					300				
Ala	His	Ser	Val	Thr	Arg	Gly	Leu	Ala	Tyr	Leu	His	Thr	Glu	Leu	Pro
305					310					315					320
Arg	Gly	Asp	His	Tyr	Lys	Pro	Ala	Ile	Ser	His	Arg	Asp	Leu	Asn	Ser
				325					330					335	
Arg	Asn	Val	Leu	Val	Lys	Asn	Asp	Gly	Thr	Cys	Val	Ile	Ser	Asp	Phe
			340					345					350		
Gly	Leu	Ser	Met	Arg	Leu	Thr	Gly	Asn	Arg	Leu	Val	Arg	Pro	Gly	Glu
		355					360					365			
Glu	Asp	Asn	Ala	Ala	Ile	Ser	Glu	Val	Gly	Thr	Ile	Arg	Tyr	Met	Ala
	370					375					380				
Pro	Glu	Val	Leu	Glu	Gly	Ala	Val	Asn	Leu	Arg	Asp	Cys	Glu	Ser	Ala
385					390					395					400

[illegible]

<210> 82

<211> 1038

<212> PRT

<213> Homo sapiens

<400> 82

Met	Thr	Ser	Ser	Leu	Gln	Arg	Pro	Trp	Arg	Val	Pro	Trp	Leu	Pro	Trp
1				5					10					15	
Thr	Ile	Leu	Leu	Val	Ser	Thr	Ala	Ala	Ala	Ser	Gln	Asn	Gln	Glu	Arg
		20					25						30		
Leu	Cys	Ala	Phe	Lys	Asp	Pro	Tyr	Gln	Gln	Asp	Leu	Gly	Ile	Gly	Glu
		35					40					45			
Ser	Arg	Ile	Ser	His	Glu	Asn	Gly	Thr	Ile	Leu	Cys	Ser	Lys	Gly	Ser
	50					55					60				
Thr	Cys	Tyr	Gly	Leu	Trp	Glu	Lys	Ser	Lys	Gly	Asp	Ile	Asn	Leu	Val
65				70						75					80
Lys	Gln	Gly	Cys	Trp	Ser	His	Ile	Gly	Asp	Pro	Gln	Glu	Cys	His	Tyr
				85					90					95	
Glu	Glu	Cys	Val	Val	Thr	Thr	Thr	Pro	Pro	Ser	Ile	Gln	Asn	Gly	Thr
			100					105					110		
Tyr	Arg	Phe	Cys	Cys	Cys	Ser	Thr	Asp	Leu	Cys	Asn	Val	Asn	Phe	Thr
		115					120					125			
Glu	Asn	Phe	Pro	Pro	Pro	Asp	Thr	Thr	Pro	Leu	Ser	Pro	Pro	His	Ser
	130					135					140				
Phe	Asn	Arg	Asp	Glu	Thr	Ile	Ile	Ile	Ala	Leu	Ala	Ser	Val	Ser	Val
145				150						155					160
Leu	Ala	Val	Leu	Ile	Val	Ala	Leu	Cys	Phe	Gly	Tyr	Arg	Met	Leu	Thr
				165					170					175	
Gly	Asp	Arg	Lys	Gln	Gly	Leu	His	Ser	Met	Asn	Met	Met	Glu	Ala	Ala
			180					185					190		
Ala	Ser	Glu	Pro	Ser	Leu	Asp	Leu	Asp	Asn	Leu	Lys	Leu	Leu	Glu	Leu
		195					200					205			
Ile	Gly	Arg	Gly	Arg	Tyr	Gly	Ala	Val	Tyr	Lys	Gly	Ser	Leu	Asp	Glu
	210					215					220				
Arg	Pro	Val	Ala	Val	Lys	Val	Phe	Ser	Phe	Ala	Asn	Arg	Gln	Asn	Phe
225				230						235					240
Ile	Asn	Glu	Lys	Asn	Ile	Tyr	Arg	Val	Pro	Leu	Met	Glu	His	Asp	Asn
				245					250					255	

Ile	Ala	Arg	Phe	Ile	Val	Gly	Asp	Glu	Arg	Val	Thr	Ala	Asp	Gly	Arg	260	265	270
Met	Glu	Tyr	Leu	Leu	Val	Met	Glu	Tyr	Tyr	Pro	Asn	Gly	Ser	Leu	Cys	275	280	285
Lys	Tyr	Leu	Ser	Leu	His	Thr	Ser	Asp	Trp	Val	Ser	Ser	Cys	Arg	Leu	290	295	300
Ala	His	Ser	Val	Thr	Arg	Gly	Leu	Ala	Tyr	Leu	His	Thr	Glu	Leu	Pro	305	310	315
Arg	Gly	Asp	His	Tyr	Lys	Pro	Ala	Ile	Ser	His	Arg	Asp	Leu	Asn	Ser	325	330	335
Arg	Asn	Val	Leu	Val	Lys	Asn	Asp	Gly	Thr	Cys	Val	Ile	Ser	Asp	Phe	340	345	350
Gly	Leu	Ser	Met	Arg	Leu	Thr	Gly	Asn	Arg	Leu	Val	Arg	Pro	Gly	Glu	355	360	365
Glu	Asp	Asn	Ala	Ala	Ile	Ser	Glu	Val	Gly	Thr	Ile	Arg	Tyr	Met	Ala	370	375	380
Pro	Glu	Val	Leu	Glu	Gly	Ala	Val	Asn	Leu	Arg	Asp	Cys	Glu	Ser	Ala	385	390	395
Leu	Lys	Gln	Val	Asp	Met	Tyr	Ala	Leu	Gly	Leu	Ile	Tyr	Trp	Glu	Ile	405	410	415
Phe	Met	Arg	Cys	Thr	Asp	Leu	Phe	Pro	Gly	Glu	Ser	Val	Pro	Glu	Tyr	420	425	430
Gln	Met	Ala	Phe	Gln	Thr	Glu	Val	Gly	Asn	His	Pro	Thr	Phe	Glu	Asp	435	440	445
Met	Gln	Val	Leu	Val	Ser	Arg	Glu	Lys	Gln	Arg	Pro	Lys	Phe	Pro	Glu	450	455	460
Ala	Trp	Lys	Glu	Asn	Ser	Leu	Ala	Val	Arg	Ser	Leu	Lys	Glu	Thr	Ile	465	470	475
Glu	Asp	Cys	Trp	Asp	Gln	Asp	Ala	Glu	Ala	Arg	Leu	Thr	Ala	Gln	Cys	485	490	495
Ala	Glu	Glu	Arg	Met	Ala	Glu	Leu	Met	Met	Ile	Trp	Glu	Arg	Asn	Lys	500	505	510
Ser	Val	Ser	Pro	Thr	Val	Asn	Pro	Met	Ser	Thr	Ala	Met	Gln	Asn	Glu	515	520	525
Arg	Asn	Leu	Ser	His	Asn	Arg	Arg	Val	Pro	Lys	Ile	Gly	Pro	Tyr	Pro	530	535	540
Asp	Tyr	Ser	Ser	Ser	Ser	Tyr	Ile	Glu	Asp	Ser	Ile	His	His	Thr	Asp	545	550	555
Ser	Ile	Val	Lys	Asn	Ile	Ser	Ser	Glu	His	Ser	Met	Ser	Ser	Thr	Pro	565	570	575
Leu	Thr	Ile	Gly	Glu	Lys	Asn	Arg	Asn	Ser	Ile	Asn	Tyr	Glu	Arg	Gln	580	585	590
Gln	Ala	Gln	Ala	Arg	Ile	Pro	Ser	Pro	Glu	Thr	Ser	Val	Thr	Ser	Leu	595	600	605
Ser	Thr	Asn	Thr	Thr	Thr	Thr	Asn	Thr	Thr	Gly	Leu	Thr	Pro	Ser	Thr	610	615	620
Gly	Met	Thr	Thr	Ile	Ser	Glu	Met	Pro	Tyr	Pro	Asp	Glu	Thr	Asn	Leu	625	630	635
His	Thr	Thr	Asn	Val	Ala	Gln	Ser	Ile	Gly	Pro	Thr	Pro	Val	Cys	Leu	645	650	655
Gln	Leu	Thr	Glu	Glu	Asp	Leu	Glu	Thr	Asn	Lys	Leu	Asp	Pro	Lys	Glu	660	665	670
Val	Asp	Lys	Asn	Leu	Lys	Glu	Ser	Ser	Asp	Glu	Asn	Leu	Met	Glu	His	675	680	685
Ser	Leu	Lys	Gln	Phe	Ser	Gly	Pro	Asp	Pro	Leu	Ser	Ser	Thr	Ser	Ser	690	695	700
Ser	Leu	Leu	Tyr	Pro	Leu	Ile	Lys	Leu	Ala	Val	Glu	Ala	Thr	Gly	Gln	705	710	715
																		720

```
<210> 83
<211> 1038
<212> PRT
<213> Homo sapiens
```

<400> 83															
Met	Thr	Ser	Ser	Leu	Gln	Arg	Pro	Trp	Arg	Val	Pro	Trp	Leu	Pro	Trp
1				5					10					15	
Thr	Ile	Leu	Leu	Val	Ser	Thr	Ala	Ala	Ala	Ser	Gln	Asn	Gln	Glu	Arg
			20					25					30		
Leu	Cys	Ala	Phe	Lys	Asp	Pro	Tyr	Gln	Gln	Asp	Leu	Gly	Ile	Gly	Glu
		35					40					45			
Ser	Arg	Ile	Ser	His	Glu	Asn	Gly	Thr	Ile	Leu	Cys	Ser	Lys	Gly	Ser
	50					55					60				
Thr	Cys	Tyr	Gly	Leu	Trp	Glu	Lys	Ser	Lys	Gly	Asp	Ile	Asn	Leu	Val
65					70					75					80

Lys	Gln	Gly	Cys	Trp	Ser	His	Ile	Gly	Asp	Pro	Gln	Glu	Cys	His	Tyr		
				85					90					95			
Glu	Glu	Cys	Val	Val	Thr	Thr	Thr	Pro	Pro	Ser	Ile	Gln	Asn	Gly	Thr		
			100					105					110				
Tyr	Arg	Phe	Cys	Cys	Cys	Ser	Thr	Asp	Leu	Cys	Asn	Val	Asn	Phe	Thr		
		115					120					125					
Glu	Asn	Phe	Pro	Pro	Pro	Asp	Thr	Thr	Pro	Leu	Ser	Pro	Pro	His	Ser		
		130				135					140						
Phe	Asn	Arg	Asp	Glu	Thr	Ile	Ile	Ile	Ala	Leu	Ala	Ser	Val	Ser	Val		
145				150						155					160		
Leu	Ala	Val	Leu	Ile	Val	Ala	Leu	Cys	Phe	Gly	Tyr	Arg	Met	Leu	Thr		
				165					170					175			
Gly	Asp	Arg	Lys	Gln	Gly	Leu	His	Ser	Met	Asn	Met	Met	Glu	Ala	Ala		
			180					185					190				
Ala	Ser	Glu	Pro	Ser	Leu	Asp	Leu	Asp	Asn	Leu	Lys	Leu	Leu	Glu	Leu		
		195				200						205					
Ile	Gly	Arg	Gly	Arg	Tyr	Gly	Ala	Val	Tyr	Lys	Gly	Ser	Leu	Asp	Glu		
		210				215					220						
Arg	Pro	Val	Ala	Val	Lys	Val	Phe	Ser	Phe	Ala	Asn	Arg	Gln	Asn	Phe		
225					230					235					240		
Ile	Asn	Glu	Lys	Asn	Ile	Tyr	Arg	Val	Pro	Leu	Met	Glu	His	Asp	Asn		
			245						250					255			
Ile	Ala	Arg	Phe	Ile	Val	Gly	Asp	Glu	Arg	Val	Thr	Ala	Asp	Gly	Arg		
		260						265					270				
Met	Glu	Tyr	Leu	Leu	Val	Met	Glu	Tyr	Tyr	Pro	Asn	Gly	Ser	Leu	Cys		
		275					280					285					
Lys	Tyr	Leu	Ser	Leu	His	Thr	Ser	Asp	Trp	Val	Ser	Ser	Cys	Arg	Leu		
	290					295					300						
Ala	His	Ser	Val	Thr	Arg	Gly	Leu	Ala	Tyr	Leu	His	Thr	Glu	Leu	Pro		
305					310					315					320		
Arg	Gly	Asp	His	Tyr	Lys	Pro	Ala	Ile	Ser	His	Arg	Asp	Leu	Asn	Ser		
			325						330					335			
Arg	Asn	Val	Leu	Val	Lys	Asn	Asp	Gly	Thr	Cys	Val	Ile	Ser	Asp	Phe		
		340						345					350				
Gly	Leu	Ser	Met	Arg	Leu	Thr	Gly	Asn	Arg	Leu	Val	Arg	Pro	Gly	Glu		
		355					360					365					
Glu	Asp	Asn	Ala	Ala	Ile	Ser	Glu	Val	Gly	Thr	Ile	Arg	Tyr	Met	Ala		
		370				375					380						
Pro	Glu	Val	Leu	Glu	Gly	Ala	Val	Asn	Leu	Arg	Asp	Cys	Glu	Ser	Ala		
385					390					395					400		
Leu	Lys	Gln	Val	Asp	Met	Tyr	Ala	Leu	Gly	Leu	Ile	Tyr	Trp	Glu	Ile		
			405						410					415			
Phe	Met	Arg	Cys	Thr	Asp	Leu	Phe	Pro	Gly	Glu	Ser	Val	Pro	Glu	Tyr		
		420						425					430				
Gln	Met	Ala	Phe	Gln	Thr	Glu	Val	Gly	Asn	His	Pro	Thr	Phe	Glu	Asp		
		435					440					445					
Met	Gln	Val	Leu	Val	Ser	Arg	Glu	Lys	Gln	Arg	Pro	Lys	Phe	Pro	Glu		
		450				455					460						
Ala	Trp	Lys	Glu	Asn	Ser	Leu	Ala	Val	Arg	Ser	Leu	Lys	Glu	Thr	Ile		
465					470					475					480		
Glu	Asp	Cys	Trp	Asp	Gln	Asp	Ala	Glu	Ala	Arg	Leu	Thr	Ala	Gln	Cys		
			485						490					495			
Ala	Glu	Glu	Arg	Met	Ala	Glu	Leu	Met	Met	Ile	Trp	Glu	Arg	Asn	Lys		
			500					505					510				
Ser	Val	Ser	Pro	Thr	Val	Asn	Pro	Met	Ser	Thr	Ala	Met	Gln	Asn	Glu		
		515				520						525					
Arg	Asn	Leu	Ser	His	Asn	Arg	Arg	Val	Pro	Lys	Ile	Gly	Pro	Tyr	Pro		
		530				535						540					



Asp	Tyr	Ser	Ser	Ser	Ser	Tyr	Ile	Glu	Asp	Ser	Ile	His	His	Thr	Asp
545					550					555					560
Ser	Ile	Val	Lys	Asn	Ile	Ser	Ser	Glu	His	Ser	Met	Ser	Ser	Thr	Pro
				565					570						575
Leu	Thr	Ile	Gly	Glu	Lys	Asn	Arg	Asn	Ser	Ile	Asn	Tyr	Glu	Arg	Gln
			580					585					590		
Gln	Ala	Gln	Ala	Arg	Ile	Pro	Ser	Pro	Glu	Thr	Ser	Val	Thr	Ser	Leu
		595					600					605			
Ser	Thr	Asn	Thr	Thr	Thr	Thr	Asn	Thr	Thr	Gly	Leu	Thr	Pro	Ser	Thr
	610					615					620				
Gly	Met	Thr	Thr	Ile	Ser	Glu	Met	Pro	Tyr	Pro	Asp	Glu	Thr	Asn	Leu
625					630					635					640
His	Thr	Thr	Asn	Val	Ala	Gln	Ser	Ile	Gly	Pro	Thr	Pro	Val	Cys	Leu
			645						650					655	
Gln	Leu	Thr	Glu	Glu	Asp	Leu	Glu	Thr	Asn	Lys	Leu	Asp	Pro	Lys	Glu
			660					665					670		
Val	Asp	Lys	Asn	Leu	Lys	Glu	Ser	Ser	Asp	Glu	Asn	Leu	Met	Glu	His
		675					680					685			
Ser	Leu	Lys	Gln	Phe	Ser	Gly	Pro	Asp	Pro	Leu	Ser	Ser	Thr	Ser	Ser
	690					695					700				
Ser	Leu	Leu	Tyr	Pro	Leu	Ile	Lys	Leu	Ala	Val	Glu	Ala	Thr	Gly	Gln
705					710					715					720
Gln	Asp	Phe	Thr	Gln	Thr	Ala	Asn	Gly	Gln	Ala	Cys	Leu	Ile	Pro	Asp
				725					730					735	
Val	Leu	Pro	Thr	Gln	Ile	Tyr	Pro	Leu	Pro	Lys	Gln	Gln	Asn	Leu	Pro
			740					745					750		
Lys	Arg	Pro	Thr	Ser	Leu	Pro	Leu	Asn	Thr	Lys	Asn	Ser	Thr	Lys	Glu
		755					760					765			
Pro	Arg	Leu	Lys	Phe	Gly	Ser	Lys	His	Lys	Ser	Asn	Leu	Lys	Gln	Val
	770					775					780				
Glu	Thr	Gly	Val	Ala	Lys	Met	Asn	Thr	Ile	Asn	Ala	Ala	Glu	Pro	His
785					790					795					800
Val	Val	Thr	Val	Thr	Met	Asn	Gly	Val	Ala	Gly	Arg	Asn	His	Ser	Val
				805					810					815	
Asn	Ser	His	Ala	Ala	Thr	Thr	Gln	Tyr	Ala	Asn	Arg	Thr	Val	Leu	Ser
			820					825					830		
Gly	Gln	Thr	Thr	Asn	Ile	Val	Thr	His	Arg	Ala	Gln	Glu	Met	Leu	Gln
		835					840					845			
Asn	Gln	Phe	Ile	Gly	Glu	Asp	Thr	Arg	Leu	Asn	Ile	Asn	Ser	Ser	Pro
	850					855					860				
Asp	Glu	His	Glu	Pro	Leu	Leu	Arg	Arg	Glu	Gln	Gln	Ala	Gly	His	Asp
865					870					875					880
Glu	Gly	Val	Leu	Asp	Arg	Leu	Val	Asp	Arg	Glu	Arg	Pro	Leu	Glu	
				885					890					895	
Gly	Gly	Arg	Thr	Asn	Ser	Asn	Asn	Asn	Ser	Asn	Pro	Cys	Ser	Glu	
			900					905				910			
Gln	Asp	Val	Leu	Ala	Gln	Gly	Val	Pro	Ser	Thr	Ala	Ala	Asp	Pro	Gly
		915						920				925			
Pro	Ser	Lys	Pro	Arg	Arg	Ala	Gln	Arg	Pro	Asn	Ser	Leu	Asp	Leu	Ser
	930					935					940				
Ala	Thr	Asn	Val	Leu	Asp	Gly	Ser	Ser	Ile	Gln	Ile	Gly	Glu	Ser	Thr
945					950					955					960
Gln	Asp	Gly	Lys	Ser	Gly	Ser	Gly	Glu	Lys	Ile	Lys	Lys	Arg	Val	Lys
				965					970					975	
Thr	Pro	Tyr	Ser	Leu	Lys	Arg	Trp	Arg	Pro	Ser	Thr	Trp	Val	Ile	Ser
			980					985					990		
Thr	Glu	Ser	Leu	Asp	Cys	Glu	Val	Asn	Asn	Asn	Gly	Ser	Asn	Arg	Ala
		995					1000						1005		

42

Val His Ser Lys Ser Ser Thr Ala Val Tyr Leu Ala Glu Gly Gly Thr  
 1010 1015 1020  
 Ala Thr Thr Met Val Ser Lys Asp Ile Gly Met Asn Cys Leu  
 1025 1030 1035

<210> 84  
 <211> 1038  
 <212> PRT  
 <213> Homo sapiens

<400> 84  
 Met Thr Ser Ser Leu Gln Arg Pro Trp Arg Val Pro Trp Leu Pro Trp  
 1 5 10 15  
 Thr Ile Leu Leu Val Ser Thr Ala Ala Ser Gln Asn Gln Glu Arg  
 20 25 30  
 Leu Cys Ala Phe Lys Asp Pro Tyr Gln Gln Asp Leu Gly Ile Gly Glu  
 35 40 45  
 Ser Arg Ile Ser His Glu Asn Gly Thr Ile Leu Cys Ser Lys Gly Ser  
 50 55 60  
 Thr Cys Tyr Gly Leu Trp Glu Lys Ser Lys Gly Asp Ile Asn Leu Val  
 65 70 75 80  
 Lys Gln Gly Cys Trp Ser His Ile Gly Asp Pro Gln Glu Cys His Tyr  
 85 90 95  
 Glu Glu Cys Val Val Thr Thr Thr Pro Pro Ser Ile Gln Asn Gly Thr  
 100 105 110  
 Tyr Arg Phe Cys Cys Cys Ser Thr Asp Leu Cys Asn Val Asn Phe Thr  
 115 120 125  
 Glu Asn Phe Pro Pro Pro Asp Thr Thr Pro Leu Ser Pro Pro His Ser  
 130 135 140  
 Phe Asn Arg Asp Glu Thr Ile Ile Ile Ala Leu Ala Ser Val Ser Val  
 145 150 155 160  
 Leu Ala Val Leu Ile Val Ala Leu Cys Phe Gly Tyr Arg Met Leu Thr  
 165 170 175  
 Gly Asp Arg Lys Gln Gly Leu His Ser Met Asn Met Met Glu Ala Ala  
 180 185 190  
 Ala Ser Glu Pro Ser Leu Asp Leu Asp Asn Leu Lys Leu Leu Glu Leu  
 195 200 205  
 Ile Gly Arg Gly Arg Tyr Gly Ala Val Tyr Lys Gly Ser Leu Asp Glu  
 210 215 220  
 Arg Pro Val Ala Val Lys Val Phe Ser Phe Ala Asn Arg Gln Asn Phe  
 225 230 235 240  
 Ile Asn Glu Lys Asn Ile Tyr Arg Val Pro Leu Met Glu His Asp Asn  
 245 250 255  
 Ile Ala Arg Phe Ile Val Gly Asp Glu Arg Val Thr Ala Asp Gly Arg  
 260 265 270  
 Met Glu Tyr Leu Leu Val Met Glu Tyr Tyr Pro Asn Gly Ser Leu Cys  
 275 280 285  
 Lys Tyr Leu Ser Leu His Thr Ser Asp Trp Val Ser Ser Cys Arg Leu  
 290 295 300  
 Ala His Ser Val Thr Arg Gly Leu Ala Tyr Leu His Thr Glu Leu Pro  
 305 310 315 320  
 Arg Gly Asp His Tyr Lys Pro Ala Ile Ser His Arg Asp Leu Asn Ser  
 325 330 335  
 Arg Asn Val Leu Val Lys Asn Asp Gly Thr Cys Val Ile Ser Asp Phe  
 340 345 350  
 Gly Leu Ser Met Arg Leu Thr Gly Asn Arg Leu Val Arg Pro Gly Glu  
 355 360 365

Glu	Asp	Asn	Ala	Ala	Ile	Ser	Glu	Val	Gly	Thr	Ile	Arg	Tyr	Met	Ala	370	375	380
Pro	Glu	Val	Leu	Glu	Gly	Ala	Val	Asn	Leu	Arg	Asp	Cys	Glu	Ser	Ala	385	390	395
Leu	Lys	Gln	Val	Asp	Met	Tyr	Ala	Leu	Gly	Leu	Ile	Tyr	Trp	Glu	Ile	405	410	415
Phe	Met	Arg	Cys	Thr	Asp	Leu	Phe	Pro	Gly	Glu	Ser	Val	Pro	Glu	Tyr	420	425	430
Gln	Met	Ala	Phe	Gln	Thr	Glu	Val	Gly	Asn	His	Pro	Thr	Phe	Glu	Asp	435	440	445
Met	Gln	Val	Leu	Val	Ser	Arg	Glu	Lys	Gln	Arg	Pro	Lys	Phe	Pro	Glu	450	455	460
Ala	Trp	Lys	Glu	Asn	Ser	Leu	Ala	Val	Arg	Ser	Leu	Lys	Glu	Thr	Ile	465	470	475
Glu	Asp	Cys	Trp	Asp	Gln	Asp	Ala	Glu	Ala	Arg	Leu	Thr	Ala	Gln	Cys	485	490	495
Ala	Glu	Glu	Arg	Met	Ala	Glu	Leu	Met	Met	Ile	Trp	Glu	Arg	Asn	Lys	500	505	510
Ser	Val	Ser	Pro	Thr	Val	Asn	Pro	Met	Ser	Thr	Ala	Met	Gln	Asn	Glu	515	520	525
Arg	Asn	Leu	Ser	His	Asn	Arg	Arg	Val	Pro	Lys	Ile	Gly	Pro	Tyr	Pro	530	535	540
Asp	Tyr	Ser	Ser	Ser	Ser	Tyr	Ile	Glu	Asp	Ser	Ile	His	His	Thr	Asp	545	550	555
Ser	Ile	Val	Lys	Asn	Ile	Ser	Ser	Glu	His	Ser	Met	Ser	Ser	Thr	Pro	565	570	575
Leu	Thr	Ile	Gly	Glu	Lys	Asn	Arg	Asn	Ser	Ile	Asn	Tyr	Glu	Arg	Gln	580	585	590
Gln	Ala	Gln	Ala	Arg	Ile	Pro	Ser	Pro	Glu	Thr	Ser	Val	Thr	Ser	Leu	595	600	605
Ser	Thr	Asn	Thr	Thr	Thr	Thr	Asn	Thr	Thr	Gly	Leu	Thr	Pro	Ser	Thr	610	615	620
Gly	Met	Thr	Thr	Ile	Ser	Glu	Met	Pro	Tyr	Pro	Asp	Glu	Thr	Asn	Leu	625	630	635
His	Thr	Thr	Asn	Val	Ala	Gln	Ser	Ile	Gly	Pro	Thr	Pro	Val	Cys	Leu	645	650	655
Gln	Leu	Thr	Glu	Glu	Asp	Leu	Glu	Thr	Asn	Lys	Leu	Asp	Pro	Lys	Glu	660	665	670
Val	Asp	Lys	Asn	Leu	Lys	Glu	Ser	Ser	Asp	Glu	Asn	Leu	Met	Glu	His	675	680	685
Ser	Leu	Lys	Gln	Phe	Ser	Gly	Pro	Asp	Pro	Leu	Ser	Ser	Thr	Ser	Ser	690	695	700
Ser	Leu	Leu	Tyr	Pro	Leu	Ile	Lys	Leu	Ala	Val	Glu	Ala	Thr	Gly	Gln	705	710	715
Gln	Asp	Phe	Thr	Gln	Thr	Ala	Asn	Gly	Gln	Ala	Cys	Leu	Ile	Pro	Asp	725	730	735
Val	Leu	Pro	Thr	Gln	Ile	Tyr	Pro	Leu	Pro	Lys	Gln	Gln	Asn	Leu	Pro	740	745	750
Lys	Arg	Pro	Thr	Ser	Leu	Pro	Leu	Asn	Thr	Lys	Asn	Ser	Thr	Lys	Glu	755	760	765
Pro	Arg	Leu	Lys	Phe	Gly	Ser	Lys	His	Lys	Ser	Asn	Leu	Lys	Gln	Val	770	775	780
Glu	Thr	Gly	Val	Ala	Lys	Met	Asn	Thr	Ile	Asn	Ala	Ala	Glu	Pro	His	785	790	795
Val	Val	Thr	Val	Thr	Met	Asn	Gly	Val	Ala	Gly	Arg	Asn	His	Ser	Val	805	810	815
Asn	Ser	His	Ala	Ala	Thr	Thr	Gln	Tyr	Ala	Asn	Arg	Thr	Val	Leu	Ser	820	825	830

Gly Gln Thr Thr Asn Ile Val Thr His Arg Ala Gln Glu Met Leu Gln  
                   835                  840                  845  
 Asn Gln Phe Ile Gly Glu Asp Thr Arg Leu Asn Ile Asn Ser Ser Pro  
           850                  855                  860  
 Asp Glu His Glu Pro Leu Leu Arg Arg Glu Gln Ala Gly His Asp  
 865                  870                  875                  880  
 Glu Gly Val Leu Asp Arg Leu Val Asp Arg Arg Glu Arg Pro Leu Glu  
                   885                  890                  895  
 Gly Gly Arg Thr Asn Ser Asn Asn Asn Asn Ser Asn Pro Cys Ser Glu  
           900                  905                  910  
 Gln Asp Val Leu Ala Gln Gly Val Pro Ser Thr Ala Ala Asp Pro Gly  
           915                  920                  925  
 Pro Ser Lys Pro Arg Arg Ala Gln Arg Pro Asn Ser Leu Asp Leu Ser  
           930                  935                  940  
 Ala Thr Asn Val Leu Asp Gly Ser Ser Ile Gln Ile Gly Glu Ser Thr  
 945                  950                  955                  960  
 Gln Asp Gly Lys Ser Gly Ser Gly Glu Lys Ile Lys Lys Arg Val Lys  
                   965                  970                  975  
 Thr Pro Tyr Ser Leu Lys Arg Trp Arg Pro Ser Thr Trp Val Ile Ser  
           980                  985                  990  
 Thr Glu Ser Leu Asp Cys Glu Val Asn Asn Asn Gly Ser Asn Arg Ala  
           995                  1000                  1005  
 Val His Ser Lys Ser Ser Thr Ala Val Tyr Leu Ala Glu Gly Gly Thr  
           1010                  1015                  1020  
 Ala Thr Thr Met Val Ser Lys Asp Ile Gly Met Asn Cys Leu  
 1025                  1030                  1035

&lt;210&gt; 85

&lt;211&gt; 2932

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 85

gctccgcgcc gagggctgga ggatgcgttc cctgggggtcc ggacttatga aaatatgcat 60  
 cagtttaata ctgtcttgga attcatgaga tggagcata ggtcaaagct gtttgagaa 120  
 aatcagaagt acagttttat ctaggcacat cttggaggag tcgtaagaaa gcagtgggag 180  
 ttgaagtcac tgtcaagtgc ttgcgatctt ttacaagaaa atctcactga atgatagtc 240  
 tttaaattgg tgaagtagca agaccaatta ttaaagggtga cagtacacag gaaacattac 300  
 aattgaacaa tgactcagct atacatttac atcagattat tgggagccta tttgttcac 360  
 atttctcgtg ttcaaggaca gaatctggat agtatgcttc atggcactgg gatgaaatca 420  
 gactccgacc agaaaaagtc agaaaaatgga gtaaccttag caccagagga taccttgct 480  
 tttttaaagt gctattgctc agggcactgt ccagatgatg ctattaataa cacatgcata 540  
 actaatggac attgctttgc catcatagaa gaagatgacc agggagaaac cacattagct 600  
 tcagggtgta tgaaatatga aggatctgat tttcagtgc aagattctcc aaaagcccag 660  
 ctacgccgga caatagaatg ttgtcggacc aatttatgta accagtattt gcaaccaca 720  
 ctgccccctg ttgtcatagg tccgtttttt gatggcagca ttcgatggct gggtttgctc 780  
 atttctatgg ctgtctgcat aattgctatg atcatcttct ccagctgctt ttgttacaaa 840  
 cattattgca agagcatctc aagcagacgt cgttacaatc gtgatttgga acaggatgaa 900  
 gcatttattc cagttggaga atcactaaaa gaccttattg accagtcaca aagttctgg 960  
 agtgggtctg gactaccttt attggttcag cgaactattg ccaaacagat tcagatggtc 1020  
 cggcaagttg gtaaaaggccg atatggagaa gtatggatgg gcaaattggc tggcgaaaaa 1080  
 gtggcgggtg aagtattctt taccactgaa gaagccagct gggttcgaga aacagaaatc 1140  
 taccaaactg tgctaatgcg ccatgaaaaa atacttggtt tcatagcggc agacattaaa 1200  
 ggtacaggtt cctggactca gctctatttg atactgatt accatgaaaa tggatctctc 1260  
 tatgacttcc tgaaatgtgc tacactggac accagacccc tgcttaaat ggcttattca 1320  
 gctgcctgtg gtctgtgcca cctgcacaca gaaatttatg gcacccaagg aaagcccgc 1380  
 attgctcatc gagacctaaa gagcaaaaac atctctatca agaaaaatgg gagttgctgc 1440

```

attgctgacc tgggccttgc tgttaaattc aacagtgaca caaatgaagt tgatgtgccc 1500
ttgaatacca ggggtgggcac caaacgctac atggctcccg aagtgtctgga cgaaagcctg 1560
aacaaaaacc acttccagcc ctacatcatg gctgacatct acagcttcgg cctaatactt 1620
tgggagatgg ctcgctcgttg tatcacagga gggatcgtgg aagaatacca attgccatat 1680
tacaacatgg taccgagtga tccgtcatat gaagatatgc gtgaggttgt gtgtgtcaaa 1740
cgtttgccgg caatttgtgc taatcgggtg aacagtgatg aatgtctacg agcagttttg 1800
aagctaattg cagaatgctg ggcccacaat ccagcctcca gactcacagc attgagaatt 1860
aagaagacgc ttgccaagat ggttgaatcc caagatgtaa aaatctgatg gttaaaccat 1920
cggaggagaa actctagact gcaagaactg tttttaccca tggcatgggt ggaattagag 1980
tggaaataagg atgttaactt ggttctcaga ctctttcttc actacgtgtt cacaggctgc 2040
taatattaaa cttttcagta ctcttattag gatacaagct gggaaacttct aaacacttca 2100
ttctttatat atggacagct ttattttaaa tgtggttttt gatgcctttt ttttaagtggg 2160
tttttatgaa ctgcatcaag acttcaatcc tgattagtgt ctccagtc aa gctctgggta 2220
ctgaattgcc tgttcataaa acggtgcttt ctgtgaaagc ctttaagaaga taaatgagcg 2280
cagcagagat ggagaaatag actttgcctt ttacctgaga cattcagttc gtttgtattc 2340
tacctttgta aaacagccta tagatgatga tgtgtttggg atactgctta ttttatgata 2400
gtttgtcctg tgccttagt gatgtgtgtg tgtctccatg cacatgcacg ccgggattcc 2460
tctgctgcc a tttgaattag aagaaaataa tttatatgca tgcacaggaa gatattggtg 2520
gccggtgggt ttgtgcttta aaaatgcaat atctgaccaa gattcgccaa tctcatacaa 2580
gccatttact ttgcaagtga gatagcttcc ccaccagctt tattttttta catgaaagct 2640
gatgccaagg ccaaaagaag tttaaagcat ctgtaaat t ggactgtttt ccttcaacca 2700
ccattttttt tgtggttatt attttgtca cggaaagcat cctctccaaa gttggagctt 2760
ctattgccat gaaccatgct taaaaagaaa gcacttctta ttgaagtga ttcctgcatt 2820
tgatagcaat gtaagtgcct ataaccatgt tctatatctt ttattctcag taacttttaa 2880
aagggaagtt atttatattt tgtgtataat gtgctttatt tgcaaatcac cc 2932

```

&lt;210&gt; 86

&lt;211&gt; 1575

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 86

```

gcaaaacttcc ttgataacat gcttttgcca agtgcaggaa aattaaatgt gggcaccaag 60
aaagaggatg gtgagagtac agccccacc ccccgccaa aggtcttgcg ttgtaaatgc 120
caccaccatt gtccagaaga ctcagtcaac aatatttgca gcacagacgg atattgtttc 180
acgatgatag aagaggatga ctctgggttg cctgtggtca cttctgggtg cctaggacta 240
gaaggctcag attttcagtg tcgggacact cccattcctc atcaaagaag atcaattgaa 300
tgetgcacag aaaggaacga atgtaataaa gacctacacc ctacactgcc tccattgaaa 360
aacagagatt ttgttgatgg acctatacac cacagggtct tacttatatc tgtgactgtc 420
tgtagtttgc tcttggtcct tatcatatta ttttgttact tccggtataa aagacaagaa 480
accagacctc gatacagcat tgggttagaa caggatgaaa cttacattcc tccctggagaa 540
tccctgagag acttaattga gcagtctcag agctcaggaa gtggatcagg cctccctctg 600
ctgggtccaa ggactatagc taagcagatt cagatggtga aacagattgg aaaaggtcgc 660
tatggggaag tttggatggg aaagtggcgt ggcgaaaagg tagctgtgaa agtgttcttc 720
accacagagg aagccagctg gttcagagag acagaaatat atcagacagt gttgatgagg 780
catgaaaaca ttttgggttt cattgctgca gatatacaag ggacagggtc ctggaccag 840
ttgtacctaa tcacagacta tcatgaaaat ggttcccttt atgattatct gaagtcacc 900
accctagacg ctaaatcaat gctgaagtta gcctactctt ctgtcagtgg cttatgtcat 960
ttacacacag aaatctttag tactcaaggc aaaccagcaa ttgcccacg agatctgaaa 1020
agtaaaaaca ttctgggtgaa gaaaaatgga acttgctgta ttgctgacct gggcctggct 1080
gttaaattta ttagtgatag aaatgaagtt gacataccac ctaacactcg agttggcacc 1140
aaacgctata tgcctccaga agtggtggac gagagcttga acagaaatca cttccagctc 1200
tacatcatgg ctgacatgta tagttttggc ctcatccttt gggaggttgc taggagatgt 1260
gtatcaggag gtagtgga agaataccag cttccttata atgacctagt gccagtgac 1320
ccctcttatg aggacatgag ggagattgtg tgcataaga agttacgcc ctcattccca 1380
aaccggtgga gcagtgatga gtgtctaagg cagatgggaa aactcatgac agaattgtg 1440
gctcacaatc ctgcatcaag gctgacagcc ctgcggtta agaaaacact tgccaaaatg 1500
tcagagtccc aggacattaa actctgatag gagaggaaaa gtaagcatct ctgcagaaag 1560

```

```

tctgattttc aatgcaagga ttcacaaaaa gccagctac gcaggacaat agaatgttgt 600
cggaccaatt tgtgcaacca atatttgcag cctacactgc cccctgtcgt tataggccca 660
ttctttgatg gcagcgtccg atggctggct gtgctcatct ctatggctgt ctgtattgtc 720
gccatgatcg tcttctccag ctgcttctgt tacaacatt actgtaagag tatctcaagc 780
agaggctcgtt acaaccgtga cttggaacag gatgaagcat ttattccagt aggagaatca 840
ctgaaagacc tgattgacca gtcacaaaagc tctggtagtg gatctggatt acctttattg 900
gttcagcgaa ctattgccaa acagattcag atgggtcggc aggttggtaa aggccggtat 960
ggagaagtat ggatgggtaa atggcgtggt gaaaaagtgg ctgtcaaagt attttttacc 1020
actgaagaag ctagctggtt tagagaaaca gaaatctacc agacgggtgtt aatgcgtcat 1080
gaaaatatac ttggttttat agctgcagac attaaaggca ccggttcctg gactcagctg 1140
tatttgatta ctgattacca tgagaatggg tctctctatg acttcttgaa atgtgccacc 1200
ctggacacca gagccctact caagttagct tattctgctg cctgtggtct gtgccacctc 1260
cacacagaaa tttatggcac gcaaggcaag cctgcaattg ctcatcgaga cctgaagagc 1320
aaaaacatcc ttattaagaa aaatggtagt tgctgtattg ctgacctggg cctagctggt 1380
aaattcaaca gtgacacaaa tgaagtgtac atacccttga acaccagggt gggcaccagg 1440
cggtagcatg tccagaagt gctggacgag agcctgagta aaaaccattt ccagccctac 1500
atcatggctg acatctacag ctttggtttg atcatttggg agatggcccg tcgctgtatt 1560
acaggaggaa tcgtggagga atatcaatta ccatattaca acatgggtgcc tagtgaccca 1620
tcttatgaag acatgcgtga ggtcgtgtgt gtgaaacgct tgcggccaat cgtctctaac 1680
cgctggaaca gtgatgaatg tcttcgagcc gttttgaagc tgatgtcaga atgctgggcc 1740
cataatccag catccagact cacagctttg agaatcaaga agacgctcgc aaagatgggt 1800
gaatcccagg atgtaaagat ttgacaaaaca gttttgagaa agaatttaga ctgcaagaaa 1860
ttcaccgag gaagggtgga gttagcatgg actaggatgt cggcttggtt tccagactct 1920
ctcctctaca tcttcacagg ctgctaacag taaactttca ggactctgca gaatgcaggg 1980
ttggagcttc agacatagga cttcagacat gctgttcttt gcgtatggac agctttgttt 2040
taaagtgtgg cttttgatgc ctttttggtt tttatgaatt gcatcaagac tccaatcctg 2100
ataagaagtc tctggtcaaa ctctggttac tcaactatct gtccataaag tgggtgctttc 2160
tgtgaaagcc ttaaggaaat tagtgagctc agcagagatg gagaaaggca tatttgccct 2220
ctacagagaa aatatctgtc tgtgttctgt ctctgtaaac agcctggact atgatctctt 2280
tgggtagctg cttggttgat gatggtgcat catgcctctg atatgcatac cagacttctt 2340
ctgctgccat gggcttacaa gacaagaatg tgaagggtgc acaggacggt atttgtggcc 2400
agtggtttaa atatgcaata tctaactgac attcgccaat ctcataaaaag ccatctacct 2460
tgtaactgaa gtaacttctc taccaacttt atttttagca taatagttgt aaaggccaaa 2520
ctatgtataa agtggtccata gactcgaact gttttctctc agtcaccatt ttgttttctt 2580
tttggttaatt atttttgtta tataattcct cctatccaga attggcgtct actgtcttga 2640
accatacttt gaaagaaatg cctcttctct gagtctgcct tactgcactc gatcaccatg 2700
tgcatacctc tgatcaaatt ctggagtctt tgttctcggt acctcttaaa aagggaat 2760
gtgtatcatg tgtagtgtgc ttttattttc aaaatcttca tagcctttat tctagccatt 2820
tttacctaca tactcattct gtacaaaaca gctcactcgg tctcacggct gatcctcagt 2880
ggaaatgatt taaagtagag ctgtgtacga atttcagaat tcatgtattt aaaaacttca 2940
cactaacact ttactaagat attgtctcat atcttttatg aggatgtcag ctgattttca 3000
atgactataa atgtatctta gctatctaaa tcttttgaaa tttggtttta taatttctgg 3060
tccctaactt gtgaagacaa agaggcagaa gtacccagtc taccacattt acactgtaca 3120
ttattaaata aaaaaatgta tattttaaaa aaaaaaaaaa aaaaaaa 3167

```

&lt;210&gt; 89

&lt;211&gt; 3167

&lt;212&gt; DNA

&lt;213&gt; Rattus norvegicus

&lt;400&gt; 89

```

gaattcatga gatggaaaca taggtcaaag ctgtttggag aaattggaac tacagtttta 60
tctagccaca tctctgagaa gtctgaagaa agcagcaggt gaaagtcatt gtcaagtgat 120
tttgttcttc tgtaaggaaa cctcgttcag taaggccggt tacttcagtg aaacagcagg 180
accagtaatc aagggtggccc ggacaggaca cgtgcgaatt ggacaatgac tcagctatac 240
acttacatca gattactggg agcctgtctg ttcattcatt ctcatgttca agggcagaat 300
ctagatagta tgctccatgg tactggtatg aaatcagacg tggaccagaa gaagccggaa 360
aatggagtga cgtagcacc agaggacacc ttacctttct taaaatgcta ttgctcagg 420

```

```

cactgcccag atgacgctat taataacaca tgcataacta atggccattg ctttgccatt 480
atagaagaag atgacagagg agaaaccacg ttaacttctg ggtgtatgaa gtatgaaggc 540
tctgattttc aatgcaagga ttcacaaaaa gccagctac gcaggacaat agaattgtgt 600
cggaccaatt tgtgcaacca atatttgcag cctacactgc cccctgtcgt tataggccca 660
ttctttgatg gcagcgctcg atggctggct gtgctcatct ctatggctgt ctgtattgtc 720
gccatgatcg tcttctccag ctgcttctgt taaaaacatt actgtaagag tatctcaagc 780
agaggctcgtt acaaccgtga cttggaacag gatgaagcat ttattccagt aggagaatca 840
ctgaaagacc tgattgacca gtcacaaagc tctggtagtgt gatctggatt acctttattg 900
gttcacgcaa ctattgccaa acagattcag atggttcggc aggttggttaa aggcgggtat 960
ggagaagtat ggatgggtaa atggcgtggt gaaaaagtgg ctgtcaaagt attttttacc 1020
actgaagaag ctagctgggt tagagaaaca gaaatctacc agacggtgtt aatgcgtcat 1080
gaaaatatac ttggttttat agctgcagac attaaaggca ccggttcctg gactcagctg 1140
tatttgatta ctgattacca tgagaatggg tctctctatg acttccctgaa atgtgccacc 1200
ctggacacca gagccctact caagttagct tattctgctg cctgtgggtct gtgccacctc 1260
cacacagaaa tttatggcac gcaaggcaag cctgcaattg ctcatcgaga cctgaagagc 1320
aaaaacatcc ctatccagact cacagctttg agaatcaaga agacgctcgc aaagatggtt 1380
aaattcaaca gtgacacaaa tgaagttgac atacccttga acaccagggg gggcaccagg 1440
cggtagatgg ctccagaagt gctggacgag agcctgagta aaaaccattt ccagccctac 1500
atcatggctg acatctacag ctttggtttg atcatttggg agatggcccg tcgctgtatt 1560
acaggaggaa tcgtggagga atatcaatta ccatattaca acatggtgcc tagtgacca 1620
tcttatgaag acatgcgtga ggtcgtgtgt gtgaaacgct tgcggccaat cgtctctaac 1680
cgctggaaca gtgatgaatg tcttcgagcc gttttgaagc tgatgtcaga atgctgggcc 1740
cataatccag catccagact cacagctttg agaatcaaga agacgctcgc aaagatggtt 1800
gaatcccagg atgtaaagat ttgacaaaaca gttttgagaa agaatttaga ctgcaagaaa 1860
ttcacccgag gaagggtgga gttagcatgg actaggatgt cggcttggtt tccagactct 1920
ctcctctaca tcttcacagg ctgctaacag taaactttca ggactctgca gaatgcaggg 1980
ttggagcttc agacatagga cttcagacat gctgttcttt gcgtatggac agctttgttt 2040
taaatgtggg cttttgatgc ctttttggtt tttatgaatt gcatcaagac tccaatcctg 2100
ataagaagtc tctggtcaaa ctctggttac tcaactatct gtccataaag tgggtgctttc 2160
tgtgaaagcc ttaaggaaat tagtgagctc agcagagatg gagaaaggca tatttgccct 2220
ctacagagaa aatatctgtc tgtgttctgt ctctgtaaac agcctggact atgatctctt 2280
tgggatgctg cctggttgat gatggtgcat catgcctctg atatgcatac cagacttcct 2340
ctgctgccat gggcttacaa gacaagaatg tgaaggttgc acaggacggt atttgtggcc 2400
agtggtttaa atatgcaata tctaactgac attcgccaat ctcataaaaag ccatctacct 2460
tgtaactgaa gtaacttctc taccaacttt atttttagca taatagttgt aaaggccaaa 2520
ctatgtataa agtggtccata gactcgaact gttttcctcc agtcaccatt ttgttttctc 2580
tttggttaatt atttttgtta tataattcct cctatccaga attggcgctc actgtcttga 2640
accatacttt gaaagaaatg cctcttctct gagtctgcct tactgcatct gatcaccatg 2700
tgcatacctc tgatcaaat ctggagtctt tgttctcggt acctcttaa aagggaaatt 2760
gtgtatcatg tgtagtgtgc ttttattttc aaaatcttca tagcctttat tctagccatt 2820
tttacctaca tactcattct gtacaaaaca gctcactcgg tctcacggct gatcctcagt 2880
ggaaatgatt taaagtagag ctgtgtacga atttcagaat tcatgtattt aaaaacttca 2940
cactaacact ttactaagat attgtctcat atcttttatg aggatgtcag ctgattttca 3000
atgactataa atgtatctta gctatctaaa tcttttgaaa tttggtttta taatttctgg 3060
tcctaactt gtgaagacaa agaggcagaa gtaccagtc taccacattt aactgtaca 3120
ttattaaata aaaaaatgta tatttttaaa aaaaaaaaaa aaaaaaa 3167

```

&lt;210&gt; 90

&lt;211&gt; 3003

&lt;212&gt; DNA

&lt;213&gt; Rattus norvegicus

&lt;400&gt; 90

```

cgttcagtaa ggccggtttac ttcagtgaag cagcaggacc agtaatcaag gtggcccggg 60
caggacacgt gcgaattgga caatgactca gctatacact tacatcagat tactgggagc 120
ctgtctgttc atcatctctc atgttcaagg gcagaatcta gatagtatgc tccatggtac 180
tggtatgaaa tcagacgtgg accagaagaa gccggaaaat ggagtacgt tagcaccaga 240
ggacacctta cctttcttaa aatgctattg ctcaggacac tgcccagatg acgctattaa 300

```

ccaacaggta ccctt

1575

&lt;210&gt; 87

&lt;211&gt; 2032

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 87

```
cgcgggggcg cgagtcggcg gggcctcgcg ggacgcgggc agtgcggaga ccgcggcgct 60
gaggacgcgg gagccgggag cgcacgcgcg gggaggagtt cagcctactc tttcttagat 120
gtgaaaggaa aggaagatca tttcatgcct tgttgataaa ggttcagact tctgctgatt 180
cataaccatt tggctctgag ctatgacaag agaggaaaca aaaagttaaa cttacaagcc 240
tgccataagt gagaagcaaa cttccttgat aacatgcttt tgcgaagtgc aggaaaatta 300
aatgtgggca ccaagaaaga ggatggtag agtacagccc ccaccccccg tccaaaggct 360
ttgcgttgta aatgccacca ccattgtcca gaagactcag tcaacaatat ttgcagcaca 420
gacggatatt gtttcacgat gatagaagag gatgactctg ggttgccctgt ggtcacttct 480
ggttgcctag gactagaagg ctacagatctt cagtgtcggg acactcccat tcctcatcaa 540
agaagatcaa ttgaatgctg cacagaaagg aacgaatgta ataaagacct acaccctaca 600
ctgcctccat tgaaaaacag agattttgtt gatggacctt tacaccacag ggctttactt 660
atatctgtga ctgtctgtag tttgctcttg gtccttatca tattattttg ttacttccg 720
tataaaagac aagaaaccag acctcgatac agcattgggt tagaacagga tgaaacttac 780
attcctctctg gagaatccct gagagactta attgagcagt ctacagagctc aggaagtggg 840
tcaggcctcc ctctgctggt ccaaaggact atagctaagc agattcagat ggtgaaacag 900
attggaaaag gtcgctatgg ggaagtttgg atgggaaagt ggcgtggcga aaaggtagct 960
gtgaaagtgt tcttcaccac agaggaagcc agctgggtca gagagacaga aatatatcag 1020
acagtgttga tgaggcatga aaacattttg ggtttcattg ctgcagatat caaagggaca 1080
gggtcctgga cccagttgta cctaatacaca gactatcatg aaaatgggtc cctttatgat 1140
tatctgaagt ccaccaccct agacgctaaa tcaatgctga agttagccta ctcttctgtc 1200
agtggcttat gtcatttaca cacagaaatc tttagtactc aaggcaaac agcaattgcc 1260
catcgagatc tgaaaagtaa aaacattctg gtgaagaaaa atggaacttg ctgtattgct 1320
gacctgggccc tggctgttaa atttattagt gatacaaatg aagttgacat accacctaac 1380
actcgagttg gcaccaaacc ctatatgcct ccagaagtgt tggacgagag cttgaacaga 1440
aatcacttcc agtcttacat catggctgac atgtatagtt ttggcctcat cctttgggag 1500
gttgctagga gctgtgtatc aggaggtata gtggaagaat accagcttcc ttatcatgac 1560
ctagtccccg gtgacccctc ttatgaggac atgagggaga ttgtgtgcat caagaagtta 1620
cgccctcat tcccaaaccg gtggagcagt gatgagtgtc taaggcagat gggaaaactc 1680
atgacagaat gctgggctca caatcctgca tcaaggctga cagccctgcg ggttaagaaa 1740
acacttgcca aaatgtcaga gtcccaggac attaaactct gataggagag gaaaagtaag 1800
catctctgca gaaagccaac aggtactctt ctgtttgtgg gcagagcaaa agacatcaaa 1860
taagcatcca cagtacaagc cttgaacatc gtcctgcttc ccagtgggtt cagacctcac 1920
ctttcagggg cgcagctggg caaagacaga gaagctccca gaaggagaga ttgatccgtg 1980
tctgtttgta ggcggagaaa ccgttgggta acttgttcaa gatatgatgc at 2032
```

&lt;210&gt; 88

&lt;211&gt; 3167

&lt;212&gt; DNA

&lt;213&gt; Rattus sp.

&lt;400&gt; 88

```
gaattcatga gatggaaaca taggtcaaag ctgtttggag aaattggaac tacagtttta 60
tctagccaca tctctgagaa gtctgaagaa agcagcagg gaaagtcatt gtcaagtgat 120
tttgttcttc tgtaaggaaa cctcgttcag taaggccggt tacttcagtg aaacagcagg 180
accagtaatc aagggtggccc ggacaggaca cgtgcgaatt ggacaatgac tcagctatac 240
acttacatca gattactggg agcctgtctg ttcattcttt ctcatgttca agggcagaat 300
ctagatagta tgctccatgg tactggtag aaatcagacg tggaccagaa gaagccggaa 360
aatggagtga cgtagcacc agaggacacc ttacctttct taaaatgcta ttgctcagga 420
cactgcccag atgacgctat taataacaca tgcataacta atggccattg ctttgccatt 480
atagaagaag atgatcaggg agaaaccacg ttaacttctg ggtgtatgaa gtatgaaggc 540
```



```

tctgatttttc aatgcaagga ttcaccaaaa gccagctac gcaggacaat agaatgttgt 600
cggaccaatt tgtgcaacca atatttgag cctacactgc cccctgtcgt tataggccca 660
ttctttgatg gcagcgctcg atggctggct gtgctcatct ctatggctgt ctgtattgtc 720
gccatgatcg tcttctccag ctgcttctgt tacaaacatt actgtaagag tatctcaagc 780
agaggtcgtt acaaccgtga cttggaacag gatgaagcat ttattccagt aggagaatca 840
ctgaaagacc tgattgacca gtcacaaagc tctggtagtg gatctggatt acctttattg 900
gttcagcgaa ctattgccaa acagattcag atgggtcggc aggttggtaa aggccggtat 960
ggagaagtat ggatgggtaa atggcggtgt gaaaaagtgg ctgtcaaagt attttttacc 1020
actgaagaag ctagctgggt tagagaaaca gaaatctacc agacggtgtt aatgcgtcat 1080
gaaaatatac ttgggttttat agctgcagac attaaaggca ccggttcctg gactcagctg 1140
tatttgatta ctgattacca tgagaatggg tctctctatg acttcctgaa atgtgccacc 1200
ctggacacca gagccctact caagttagct tattctgctg cctgtggctc gtgccacctc 1260
cacacagaaa tttatggcac gcaaggcaag cctgcaattg ctcatcgaga cctgaagagc 1320
aaaaacatcc ttattaagaa aaatggtagt tgctgtattg ctgacctggg cctagctgtt 1380
aaattcaaca gtgacacaaa tgaagttgac atacccttga acaccagggt gggcaccagg 1440
cggtagcatg tccagaagt gctggacgag agcctgagta aaaaccattt ccagccctac 1500
atcatggctg acatctacag ctttggtttg atcatttggg agatggcccg tcgctgtatt 1560
acaggaggaa tcgtggagga atatcaatta ccatattaca acatggtgcc tagtgacca 1620
tcttatgaag acatgcgtga ggtcgtgtgt gtgaaacgct tgcggccaat cgtctctaac 1680
cgctggaaca gtgatgaatg tcttcgagcc gttttgaagc tgatgtcaga atgtcgggcc 1740
cataatccag catccagact cacagctttg agaattcaaga agacgctcgc aaagatgggt 1800
gaatcccagg atgtaaagat ttgacaaaaca gttttgagaa agaatttaga ctgcaagaaa 1860
ttcacccgag gaagggtgga gttagcatgg actaggatgt cggcttggtt tccagactct 1920
ctcctctaca tcttcacagg ctgctaacag taaactttca ggactctgca gaatgcaggg 1980
ttggagcttc agacatagga cttcagacat gctgttcttt gcgtatggac agctttgttt 2040
taaagtgtgg cttttgatgc ctttttggtt tttatgaatt gcataagac tccaatcctg 2100
ataagaagtc tctggtcaaa ctctggttac tcactatcct gtccataaag tgggtgctttc 2160
tgtgaaagcc ttaaggaaat tagtgagctc agcagagatg gagaaaggca tatttgccct 2220
ctacagagaa aatatctgtc tgtgttctgt ctctgtaaac agcctggact atgatctctt 2280
tgggatcgct cctggttgat gatggtgcat catgcctctg atatgcatac cagacttcct 2340
ctgctgccat gggcttacia gacaagaatg tgaagggtgc acaggacggt atttgtggcc 2400
agtggtttaa atatgcaata tctaactgac attcgccaat ctcataaaag ccatctacct 2460
tgtaactgaa gtaacttctc taccaacttt attttttagc taatagttgt aaaggccaaa 2520
ctatgtataa agtgccata gactcgaact gttttcctcc agtcaccatt ttgttttctc 2580
tttggttaatt atttttgtta tataattcct cctatccaga attggcgctc actgtcttga 2640
accatacttt gaaagaaatg cctcttctcg gagtctgcct tactgcatct gatcaccatg 2700
tgcatacctc tgatcaaat ctggagtctt tgttctcggt acctcttaaa aagggaaatt 2760
gtgtatcatg tgtagtgtgc ttttattttc aaaatcttca tagcctttat tctagccatt 2820
tttacctaca tactcattct gtacaaaaca gctcactcgg tctcacggct gatcctcagt 2880
ggaaatgatt taaagtagag ctgtgtacga atttcagaat tcatgtattt aaaaacttca 2940
cactaacact ttactaagat attgtctcat atcttttatg aggatgtcag ctgattttca 3000
atgactataa atgtatctta gctatctaaa tcttttgaaa tttggtttta taatttctgg 3060
tccctaactt gtgaagacaa agaggcagaa gtaccagtc taccacattt acactgtaca 3120
ttattaataa aaaaaatgta tattttaaaa aaaaaaaaaa aaaaaaa 3167

```

&lt;210&gt; 89

&lt;211&gt; 3167

&lt;212&gt; DNA

<213> *Rattus norvegicus*

&lt;400&gt; 89

```

gaattcatga gatggaaaca taggtcaaag ctgtttggag aaattggaac tacagtttta 60
tctagccaca tctctgagaa gtctgaagaa agcagcaggt gaaagtcat gtcaagtgat 120
tttgttcttc tgtaaggaaa cctcgttcag taaggccgtt tacttcagt aaacagcagg 180
accagtaatc aagggtggccc ggacaggaca cgtgcgaatt ggacaatgac tcagctatac 240
acttacatca gattactggg agcctgtctg ttcattcatt ctcatgttca agggcagaat 300
ctagatagta tgctccatgg tactggtagt aaatcacagc tggaccagaa gaagccgga 360
aatggagtga cgtttagcacc agaggacacc ttacctttct taaaatgcta ttgctcagga 420

```

```

cactgcccag atgacgctat taataacaca tgcataacta atggccattg ctttgccatt 480
atagaagaag atgatcaggg agaaaccacg ttaacttctg ggtgtatgaa gtatgaaggc 540
tctgattttc aatgcaagga ttcacaaaaa gccagctac gcaggacaat agaagtgtgt 600
cggaccaatt tgtgcaacca atatttgcag cctacactgc cccctgtcgt tataggccca 660
ttctttgatg gcagcgccg atggctggct gtgctcatct ctatggctgt ctgtattgtc 720
gccatgatcg tcttctccag ctgcttctgt tacaacatt actgtaagag tatctcaagc 780
agaggctcgtt acaaccgtga cttggaacag gatgaagcat ttattccagt aggagaatca 840
ctgaaagacc tgattgacca gtcacaaagc tctggtagtg gatctggatt acctttattg 900
gttcagcgaa ctattgccaa acagattcag atggttcggc aggttggtaa aggccgggtat 960
ggagaagtat ggatgggtaa atggcgtggt gaaaaagtgg ctgtcaaagt attttttacc 1020
actgaagaag ctagctgggt tagagaaaca gaaatctacc agacggtgtt aatgcgtcat 1080
gaaaatatac ttggttttat agctgcagac attaaaggca ccggttcctg gactcagctg 1140
tatttgatta ctgattacca tgagaatggg tctctctatg acttcctgaa atgtgccacc 1200
ctggacacca gagccctact caagttagct tattctgctg cctgtggctc gtgccacctc 1260
cacacagaaa tttatggcac gcaaggcaag cctgcaattg ctcatcgaga cctgaagagc 1320
aaaaacatcc ttattaagaa aaatggtagt tgctgtattg ctgacctggg cctagctggt 1380
aaattcaaca gtgacacaaa tgaagttgac atacccttga acaccagggt gggcaccagg 1440
cggtagatgg ctccagaagt gctggacgag agcctgagta aaaaccattt ccagccctac 1500
atcatggctg acatctacag ctttggtttg atcatttggg agatggcccg tcgctgtatt 1560
acaggaggaa tcgtggagga atatcaatta ccatattaca acatggtgcc tagtgacca 1620
tcttatgaag acatgcgtga ggtcgtgtgt gtgaaacgct tgcgccaat cgtctctaac 1680
cgctggaaca gtgatgaatg tcttcgagcc gttttgaagc tgatgtcaga atgctgggcc 1740
cataatccag catccagact cacagctttg agaatcaaga agacgctcgc aaagatgggt 1800
gaatcccagg atgtaaagat ttgacaaaaca gttttgagaa agaatttaga ctgcaagaaa 1860
ttcacccgag gaaggggtgga gttagcatgg actaggatgt cggtctgggt tccagactct 1920
ctcctctaca tcttcacagg ctgctaacag taaactttca ggactctgca gaatgcaggg 1980
ttggagcttc agacatagga cttcagacat gctgttcttt gcgtatggac agctttgttt 2040
taaatgtggg cttttgatgc ctttttggtt tttatgaatt gcatcaagac tccaatcctg 2100
ataagaagtc tctgggtcaa ctctggttac tcactatcct gtccataaag tgggtgctttc 2160
tgtgaaagcc ttaaggaaat tagtgagctc agcagagatg gagaaaggca tatttgccct 2220
ctacagagaa aatatctgtc tgtgttctgt ctctgtaaac agcctggact atgatctctt 2280
tggtgatgct cctgggtgat gatggtgcat catgcctctg atatgcatac cagacttctt 2340
ctgctgccat gggcttacaa gacaagaatg tgaagggttg acaggacggg atttgtggcc 2400
agtggtttaa atatgcaata tctaactgac attcgccaat ctcataaaaag ccatctacct 2460
tgtaactgaa gtaacttctc taccaacttt ttttttagca taatagtgt aaaggccaaa 2520
ctatgtataa agtggtccata gactcgaact gttttctctc agtcaccatt ttgttttctt 2580
tttggttaatt atttttgtta tataattcct cctatccaga attggcgctc actgtcttga 2640
accatacttt gaaagaaatg cctcttctct gagtctgcct tactgcatct gatcaccatg 2700
tgcatacctc tgatcaaatt ctggagtctt tgttctcggt acctcttaaa aagggaaatt 2760
gtgtatcatg tgtagtgtgc ttttattttc aaaatcttca tagcctttat tctagccatt 2820
tttacctaca tactcattct gtacaaaaca gctcactcgg tctcacggct gatcctcagt 2880
ggaaatgatt taaagtagag ctgtgtacga atttcagaat tcatgtattt aaaaacttca 2940
cactaacact ttactaagat attgtctcat atcttttatg aggatgtcag ctgattttca 3000
atgactataa atgtatctta gctatctaaa tcttttgaaa tttggtttta taatttctgg 3060
tccctaactt gtgaagacaa agaggcagaa gtacccagtc taccacattt acactgtaca 3120
ttatataata aaaaaatgta tattttaaaa aaaaaaaaaa aaaaaaa 3167

```

&lt;210&gt; 90

&lt;211&gt; 3003

&lt;212&gt; DNA

&lt;213&gt; Rattus norvegicus

&lt;400&gt; 90

```

cgttcagtaa ggccgtttac ttcagtgaag cagcaggacc agtaatcaag gtggcccgga 60
caggacacgt gcgaattgga caatgactca gctatacact tacatcagat tactgggagc 120
ctgctgtgtc atcatttctc atgttcaagg gcagaatcta gatagtatgc tccatggtac 180
tggtatgaaa tcagacgtgg accagaagaa gccggaaaat ggagtgcgt tagcaccaga 240
ggacacctta cttttcttaa aatgctattg ctcaggacac tgcccagatg acgctattaa 300

```

```

taacacatgc ataactaatg gccattgctt tgccattata gaagaagatg atcagggaga 360
aaccacgtta acttctgggt gtatgaagta tgaaggctct gattttcaat gcaaggattc 420
accaaaagcc cagctacgca ggacaataga atgttgctcg accaatttgt gcaaccaata 480
tttgacgctt acactgcccc ctgtcggtat aggcccattc ttgatggca gcgtccgatg 540
gctggctgtg ctcatctcta tggctgtctg tattgtcgcc atgatcgtct tctccagctg 600
cttctgttac aaacattact gtaagagtat ctcaagcaga ggtcggtaca accgtgactt 660
ggaaacaggat gaagcattta ttccagtagg agaactcactg aaagacctga ttgaccagt 720
acaaagctct ggtagtggat ctggattacc ttatttgggt cagcgaacta ttgccaaca 780
gattcagatg gttcggcagg ttggtaaagg ccggtatgga gaagtatgga tgggtaaatg 840
gcgtgggtgaa aaagtggctg tcaaagtatt ttttaccact gaagaagcta gctgggttag 900
agaaacagaa atctaccaga cgggtgttaat gcgtcatgaa aatatacttg gttttatagc 960
tgcagacatt aaaggcaccg gttcctggac tcagctgtat ttgattactg attaccatga 1020
gaatgggtct ctctatgact tcttgaagtg tgccacctg gacaccagag ccctactcaa 1080
gttagcttat tctgtgcct gtggctctgt ccacctccac acagaaattt atggcacgca 1140
aggcaagcct gcaattgtct atcgagacct gaagagcaaa aacatcctta ttaagaaaaa 1200
tggtagttgc tgtattgtct acctgggctt agctgttaaa ttcaacagtg acacaaatga 1260
agttgacata cccttgaaca ccagggtggg caccaggcgg tacatggctc cagaagtgtc 1320
ggacgagagc ctgagtaaaa accatttcca gccctacatc atggctgaca tctacagctt 1380
tggtttgatc atttgggaga tggcccgctg ctgtattaca ggaggaatcg tggaggaata 1440
tcaattacca tattacaaca tgggtgcctag tgacctatct tatgaagaca tgcgtgaggt 1500
cgtgtgtgtg aaacgcttgc ggccaatcgt ctctaaccgc tggaaacagt atgaatgtct 1560
tcgagccgtt ttgaagctga tgtcagaatg ctgggcccat aatccagcat ccagactcac 1620
agctttgaga atcaagaaga cgctcgcaaa gatggttgaa tcccaggatg taaagatttg 1680
acaaacagtt ttgagaaaga atttagactg caagaaattc acccgaggaa ggggtggagt 1740
agcatggact aggatgtcgg cttgggttcc agactctctc ctctacatct tcacaggctg 1800
ctaacagtaa actttcagga ctctgcagaa tgcaggggtg gagcttcaga cataggactt 1860
cagacatgct gttctttgct tatggacagc ttgttttaa atgtgggctt ttgatgcctt 1920
tttgggtttt atgaattgca tcaagactcc aatcctgata agaagctctc ggtcaaactc 1980
tggttactca ctatcctgtc cataaagtgg tgctttctgt gaaagcctta aggaattag 2040
tgagctcagc agagatggag aaaggcatat ttgccctcta cagagaaaat atctgtctgt 2100
gttctgtctc tgtaaacagc ctggactatg atctctttgg gatgctgcct ggttgatgat 2160
ggtgcatcat gcctctgata tgcataccag acttctctct ctgccatggg cttacaagac 2220
aagaatgtga aggttgaca ggacgggtatt tgtggccagt ggtttaaata tgcaatatct 2280
aatcgacatt cgccaatctc ataaaagcca tctacctgtt aactgaagta acttctctac 2340
caactttatt tttagcataa tagttgtaaa ggccaaacta tgtataaagt gtccatagac 2400
tcgaactgtt ttctccagt caccattttg ttttctttt ggtaattatt tttgttatat 2460
aattctctct atccagaatt ggcgctcact gtcttgaacc atactttgaa agaaatgcct 2520
cttctggag tctgccttac tgcactctgat caccatgtgc atacctctga tcaaattctg 2580
gagttttgt tctcggtaac tcttaaaaag ggaattgtg tatcatgtgt agtgtgtctt 2640
tattttcaaa atcttcatag cctttattct agccattttt acctacatac tcatctgtga 2700
caaaacagct cactcggctc cacggctgat cctcagtgga aatgatttaa agtagagctg 2760
tgtacgaatt tcagaattca tgtattttaa aacttcacac taacacttta ctaagatatt 2820
gtctcatatc ttttatgagg atgtcagctg attttcaatg actataaatg tatcttagct 2880
atctaaatct tttgaaattt ggttttataa tttctggctc ctaacttgtg aagacaaaga 2940
ggcagaagta ccagctctac cacatttaca ctgtacatta ttaaataaaa aaatgtatat 3000
ttt 3003

```

&lt;210&gt; 91

&lt;211&gt; 2063

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 91

```

gaattccggg gatgatgatg gtgatggtga tgatgggtgat gaggatgatg gtgatgatga 60
tgatgggtgtt ggtgatggtt tttgcatctt ccattcatga actaagtact cttattagt 120
aatttctttt ctttgccttc ctgattcttt gctggcccag ggatgacttc ctgcgtgcag 180
cggccctggc ggggtgcctg gctaccatgg accatcctgc tggtcagcac tgcgggtgct 240
tcgcagaatc aagaacggct atgtgcgttt aaagatccgt atcagcaaga ccttgggata 300

```

```

ggtgagagta gaatctctca tgaaaatggg acaatattat gctcgaaagg tagcacctgc 360
tatggccttt gggagaaatc aaaaggggac ataaatcttg taaaacaagg atgttggtct 420
cacattggag atccccaaga gtgtcactat gaagaatgtg tagtaactac cactcctccc 480
tcaattcaga atggaacata ccgtttctgc tgtgtagca cagatttatg taatgtcaac 540
tttactgaga attttccacc tcctgacaca acaccactca gtccacctca ttcatTTAAC 600
cgagatgaga caataatcat tgctttggca tcagtctctg tattagctgt tttgatagtt 660
gccttatgct ttggatacag aatgttgaca ggagaccgta aacaaggctc tcacagtatg 720
aacatgatgg aggcagcagc atccgaaccc tctcttgatc tagataatct gaaactgttg 780
gagctgattg gccgaggtcg atatggagca gtatataaag gctccttgga tgagcgcca 840
gttgctgtaa aagtgttttc ctttgcaaac cgtcagaatt ttatcaacga aaagaacatt 900
tacagagtgc ctttgatgga acatgacaac attgcccgtc ttatagttgg agatgagaga 960
gtcactgcag atggacgcat ggaatatttg cttgtgatgg agtactatcc caatggatct 1020
ttatgcaagt atttaagtct ccacacaagt gactgggtaa gctcttgccg tcttgctcat 1080
tctgttacta gaggactggc ttatcttcac acagaattac cacgaggaga tcattataaa 1140
cctgcaattt cccatcgaga tttaaacagc agaaatgtcc tagtgaaaaa tgatggaacc 1200
tgtgttatta gtgactttgg actgtccatg aggtgactg gaaatagact ggtgcgcca 1260
ggggaggaag ataatgcagc cataagcgag gttggcacta tcagatatat ggcaccagaa 1320
gtgctagaag gagctgtgaa cttgaggggac tgtgaatcag ctttgaaaca agtagacatg 1380
tatgctcttg gactaatcta ttgggagata tttatgagat gtacagacct cttcccaggg 1440
gaatccgtac cagagtacca gatggctttt cagacagagg ttggaaacca tcccactttt 1500
gaggatatgc aggttctcgt gtctagggaa aaacagagac ccaagttccc agaagcctgg 1560
aaagaaaata gcctggcagt gaggtcactc aaggagacaa tcgaagactg ttgggaccag 1620
gatgcagagg ctcggtttac tgcacagtgt gctgaggaaa ggatggctga acttatgatg 1680
atttgggaaa gaaacaaatc tgtgagccca acagtcaatc caatgtctac tgctatgcag 1740
aatgaacgta ggtgagtcaa cacaagatgg caaatcagga tcaggtgaaa agatcaagaa 1800
acgtgtgaaa actccctatt ctcttaagcg gtggcgcccc tccacctggg tcatctccac 1860
tgaatcgctg gactgtgaag tcaacaataa tggcagttaac agggcagttc attccaaatc 1920
cagcactgct gtttaccttg cagaaggagg cactgctaca accatgggtg ctaaagatat 1980
aggaatgaac tgtctgtgaa atgttttcaa gcctatggag tgaaattatt ttttgcata 2040
tttaaacatg cagaagatgt tta
2063

```

&lt;210&gt; 92

&lt;211&gt; 1964

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 92

```

atttcttttc tttgccctcc tgattcttgg ctggcccagg gatgacttcc tcgctgcagc 60
ggccctggcg ggtgccctgg ctaccatgga ccatcctgct ggtcagcact gcggtgctt 120
cgcagaatca agaacggcta tgtgcgttta aagatccgta tcagcaagac cttgggatag 180
gtgagagtag aatctctcat gaaaatggga caatattatg ctcgaaaggc agcacctgct 240
atggcctttg ggagaaatca aaaggggaca taaatcttgt aaaacaagga tgttggtctc 300
acattggaga tccccaagag tgtcactatg aagaatgtgt agtaactacc actcctccct 360
caattcagaa tggaacatac cgtttctgct gttgtagcac agatttatgt aatgtcaact 420
ttactgagaa ttttccacct cctgacacaa caccactcag tccacctcat tcatttaacc 480
gagatgagac aataatcatt gctttggcat cagtctctgt attagctgtt ttgatagttg 540
ccttatgctt tggatacaga atgttgacag gagaccgtaa acaaggctctt cacagtatga 600
acatgatgga ggcagcagca tccgaacctt ctcttgatct agataatctg aaactggttg 660
agctgattgg ccgaggtcga tatggagcag tatataaagg ctcttggtat gagcgctccag 720
ttgctgtaaa agtgttttcc tttgcaaac gtcagaattt tatcaacgaa aagaacattt 780
acagagtgcc ttgatggaa catgacaaca ttgccgctt tatagttgga gatgagagag 840
tactgcaga tggagcgtat gaatatattgc ttgtgatgga gtactatccc aatggatctt 900
tatgcaagta ttttaagtctc cacacaagtg actgggttaag ctcttgccgt cttgctcatt 960
ctgttactag aggactggct tatcttcaca cagaattacc acgaggagat cattataaac 1020
ctgcaatttc ccatcgagat ttaaacagca gaaatgtcct agtgaaaaat gatggaacct 1080
gtgttattag tgactttgga ctgtccatga ggctgactgg aaatagactg gtgcgcccag 1140
gggaggaaga taatgcagcc ataagcgagg ttggcactat cagatatatg gcaccagaag 1200
tgctagaagg agctgtgaac ttgagggact gtgaatcagc tttgaaacaa gtagacatgt 1260

```

atgctcttgg	actaatctat	tgggagatat	ttatgagatg	tacagacctc	ttcccagggg	1320
aatccgtacc	agagtaccag	atggcttttc	agacagaggt	tggaaaccat	cccacttttg	1380
aggatatgca	ggttctcgtg	tctagggaaa	aacagagacc	caagttccca	gaagcctgga	1440
aagaaaatag	cctggcagtg	aggctactca	aggagacaat	cgaagactgt	tgggaccagg	1500
atgcagaggc	tgggcttact	gcacagtgtg	ctgaggaaag	gatggctgaa	cttatgatga	1560
tttgggaaaag	aaacaaatct	gtgagcccaa	cagtcaatcc	aatgtctact	gctatgcaga	1620
atgaacgtag	gtgagtcaac	acaagatggc	aaatcaggat	caggtgaaaa	gatcaagaaa	1680
cgtgtgaaaa	ctccctattc	tcttaagcgg	tggcgccctc	ccacctgggt	catctccact	1740
gaatcgctgg	actgtgaagt	caacaataat	ggcagtaaca	gggcagttca	ttccaaatcc	1800
agcactgctg	tttaccttgc	agaaggaggc	actgctacaa	ccatggtgtc	taaagatata	1860
ggaatgaact	gtctgtgaaa	tgttttcaag	cctatggagt	gaaattattt	tttgcacat	1920
ttaaacatgc	agaagatggt	taaaaataaa	aaaaaaactg	cttt		1964

&lt;210&gt; 93

&lt;211&gt; 3611

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 93

cgcccccgga	ccccggatcg	aatccccgcc	ctccgcaccc	tggatatggt	ttctcccaga	60
cctggatatt	tttttgatat	cgtgaaacta	cgagggaat	aatttggggg	atttcttctt	120
ggctccctgc	ttttcccaca	gacatgcctt	ccgtttggag	ggccgcggca	ccccgtccga	180
ggcgaaggaa	ccccccagc	cgcgaggag	agaaatgaag	ggaatttctg	cagcggcatg	240
aaagctctgc	agctagggtc	tctcatcagc	catttgtcct	ttcaaactgt	attgtgatac	300
gggcaggatc	agtccacggg	agagaagacg	agcctcccgg	ctgtttctcc	gccggtctac	360
ttcccatatt	tcttttcttt	gccctcctga	ttcttggctg	gccagggat	gacttctctg	420
ctgcagcggc	cctggcgggg	gccctggcta	ccatggacca	tcctgctggt	cagcactgcg	480
gctgcttcgc	agaatcaaga	acggctatgt	gcgtttaaag	atccgtatca	gcaagacctt	540
gggataggtg	agagtagaat	ctctcatgaa	aatgggacaa	tattatgctc	gaaaggtagc	600
acctgctatg	gcctttggga	gaaatcaaaa	ggggacataa	atccttgtaa	acaaggtagc	660
tggctctaca	ttggagatcc	ccaagagtgt	cactatgaag	aatgtgtagt	aactaccact	720
cctccctcaa	ttcagaatgg	aacataccgt	ttctgctgtt	gtagcacaga	tttatgtaat	780
gtcaacttta	ctgagaattt	tccacctcct	gacacaacac	cactcagtcc	acctcattca	840
tttaaccgag	atgagacaat	aatcattgct	ttggcatcag	tctctgtatt	agctgttttg	900
atagttgcct	tatgcttttg	atacagaatg	ttgacaggag	accgtaaa	aggtcttcac	960
agtatgaaca	tgatggaggc	agcagcatcc	gaacctctc	ttgatctaga	taatctgaaa	1020
ctgttggagc	tgattggccg	aggctgatat	ggagcagtat	ataaaggctc	cttggatgag	1080
cgtccagtgt	ctgtaaaaag	gttttccctt	gcaaaccgtc	agaattttat	caacgaaaag	1140
aacatttaca	gagtgccttt	gatggaacat	gacaacattg	cccgccttat	agttggagat	1200
gagagagtca	ctgcagatgg	acgcatggaa	tatttgcttg	tgatggagta	ctatcccaat	1260
ggatctttat	gcaagtattt	aagtctccac	acaagtgact	gggtaagctc	ttgccgtctt	1320
gctcattctg	ttactagagg	actggcttat	cttcacacag	aattaccacg	aggagatcat	1380
tataaacctg	caatttccca	tcgagattta	aacagcagaa	atgtcctagt	gaaaaatgat	1440
ggaacctgtg	ttattagtga	ctttggactg	tccatgaggc	tgactggaaa	tagactgggtg	1500
cgcccagggg	aggaagataa	tgcagccata	agcgagggtg	gcactatcag	atatatggca	1560
ccagaagtgc	tagaaggagc	tgtgaacttg	agggactgtg	aatcagcttt	gaaacaagta	1620
gacatgtatg	ctcttggact	aatctatttg	gagatattta	tgagatgtac	agacctcttc	1680
ccaggggaat	ccgtaccaga	gtaccagatg	gcttttcaga	cagagggttg	aaaccatccc	1740
acttttgagg	atatgcagggt	tctcgtgtct	agggaaaaac	agagacccaa	gttcccagaa	1800
gacctgaaaag	aaaatagcct	ggcagtgagg	tcactcaagg	agacaatcga	agactgtttg	1860
gaccaggatg	cagaggctcg	gcttactgca	cagtgtgctg	aggaaaggat	ggctgaactt	1920
atgatgatgt	gggaaagaaa	caaactctgt	agcccaacag	tcaatccaat	gtctactgct	1980
atgcagaatg	aacgcaacct	gtcacataat	aggcgtgtgc	caaaaatttg	tccttatcca	2040
gattattctt	cctcctcata	cattgaagac	tctatccatc	atactgacag	catcgtgaag	2100
aatatttcct	ctgagcattc	tatgtccagc	acacctttga	ctatagggga	aaaaaaccga	2160
aattcaatta	actatgaacg	acagcaagca	caagctcgaa	tccccagccc	tgaacaaggt	2220
gtcaccagcc	tctccaccaa	cacaacaacc	acaaacacca	caggactcac	gccaaagtact	2280
ggcatgacta	ctatatctga	gatgccatac	ccagatgaaa	caaactctga	taccacaaat	2340

```

gttgacacagt caattggggc aaccctgtgc tgcttacagc tgacagaaga agacttggaa 2400
accaacaagc tagacccaaa agaagtgtgat aagaacctca aggaaagctc tgatgagaat 2460
ctcatggagc actctcttaa acagttcagt ggcccagacc cactgagcag tactagtctc 2520
agcttgcttt acccactcat aaaacttgca gtagaagcaa ctggacagca ggacttcaca 2580
cagactgcaa atggccaagc atgtttgatt cctgatgttc tgcctactca gatctatcct 2640
ctccccaaagc agcagaacct tcccaagaga cctactagtt tgcctttgaa caccaaaaat 2700
tcaacaaaag agccccggct aaaatttggc agcaagcaca aatcaaactt gaaacaagtc 2760
gaaactggag ttgccaagat gaatacaatc aatgcagcag aacctcatgt ggtgacagtc 2820
accatgaatg gtgtggcagg tagaaaccac agtgttaact cccatgctgc cacaacccaa 2880
tatgccaatg ggacagtact atctggccaa acaaccaaca tagtgacaca tagggcccaa 2940
gaaatgttgc agaatcagtt tattggtgag gacaccggc tgaatattaa ttccagtcct 3000
gatgagcatg agcctttact gagacgagag caacaagctg gccatgatga aggtgttctg 3060
gatcgtcttg tggacaggag ggaacggcca ctagaagggtg gccgaactaa ttccaataac 3120
aacaacagca atccatgttc agaacaagat gttcttgcac aggtgttcc aagcacagca 3180
gcagatcctg ggccatcaaa gccagaaga gcacagaggc ctaattctct ggatctttca 3240
gccacaaatg tcctggatgg cagcagtata cagtaggtg agtcaacaca agatggcaaa 3300
tcaggatcag gtgaaaagat caagaaacgt gtgaaaactc cctattctct taagcgggtg 3360
cgcccccca cctgggtcat ctccactgaa tcgctggact gtgaagtcaa caataatggc 3420
agtaacaggg cagttcatcc caaatccagc actgctgttt accttgca gaaggagcact 3480
gctacaacca tgggtgtctaa agatatagga atgaactgtc tgtgaaatgt tttcaagcct 3540
atggagtga attatTTTTT gcattcttta aacatgcaga agatgtttta aaataaaaaa 3600
aaaactgctt t 3611

```

&lt;210&gt; 94

&lt;211&gt; 3871

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 94

```

ggcctccgca ccctggatat gttttctccc agacctggat atttttttga tatcgtgaaa 60
ctacgaggga aataatttgg gggatttctt ctggctccc tgctttcccc acagacatac 120
cttccgtttg gagggccgcg gcaccccgct cgaggcgaag gaaccccccc atccgcgagg 180
gagagaaatg aagggaaattt ctgcagcggc atgaaagctc tgcagctagg tctctctatc 240
agccatttgt cctttcaaac tgtattgtga tacgggcagg atcagtccac gggagagaag 300
acgagcctcc cggtgtttt tccgcccggc tacttcccat atttcttttc tttgcccctc 360
tgattcttgg ctggcccagg gatgacttcc tcgctgcagc ggccctggcg ggtgcccctg 420
taccatgga ccatcctgct ggtcagcact cggctgctt cgcagaatca agaacggcta 480
tgtgcgttta aagatccgta tcagcaagac cttgggtag gtgagagtag aatctctcat 540
gaaaatggga caatattatg ctcgaaaggt agcacctgct atggcctttg ggagaaatca 600
aaaggggaca taaatcttgt aaaacaagga tgttggcttc acattggaga tccccagag 660
tgtcactatg aagaatgtgt agtaactacc actcctccct caattcagaa tggacatac 720
cgtttctgct gttgtagcac agatttatgt aatgtcaact ttactgagaa ttttccacct 780
cctgacacaa caccactcag tccacctcat tcatttaacc gagatgagac aataatcatt 840
gctttggcat cagtctctgt attagctgtt ttgatagttg ccttatgctt tggatacaga 900
atgttgacag gagaccgtaa acaaggtctt cacagtatga acatgatgga ggcagcagca 960
tccgaaccct ctcttgatct agataatctg aaactgttgg agctgattgg ccgaggtcga 1020
tatggagcag tatataaagg ctcccttggat gagcgtccag ttgctgtaaa agtgttttcc 1080
tttgcaaacc gtcagaattt tatcaacgaa gatgagagag tcactgcaga tggacgcatg 1200
gaatatttgc ttgtgatgga gtactatccc aatggatctt tatgcaagta ttttaagtctc 1260
cacacaagtg actgggtaag ctcttgccgt cttgtctatt ctgttactag aggactggct 1320
tatcttcaca cagaattacc acgaggagat cattataaac ctgcaatttc ccatcgagat 1380
ttaaacagca gaaatgtcct agtgaaaaat gatggaacct gtgttattag tgactttgga 1440
ctgtccatga ggctgactgg aaatagactg gtgcgcccag gggaggaaga taatgcagcc 1500
ataagcgagg ttggcactat cagatatatg gcaccagaag tgctagaagg agctgtgaac 1560
ttgagggact gtgaatcagc tttgaaacaa gtagacatgt atgctcttgg actaatctat 1620
tgggagatat ttatgagatg tacagacctc ttcccagggg aatccgtacc agagtaccag 1680
atggcctttc agacagagggt tggaaacccat cccacttttg aggatatgca ggttctcgtg 1740

```

```

tctagggaaa aacagagacc caagttccca gaagcctgga aagaaaatag cctggcagtg 1800
aggtcactca aggagacaat cgaagactgt tgggaccagg atgcagaggc tgggcttact 1860
gcacagtgtg ctgaggaaag gatggctgaa cttatgatga tttgggaaag aaacaaatct 1920
gtgagcccaa cagtcaatcc aatgtctact gctatgcaga atgaacgcaa cctgtcacat 1980
aataggcgtg tgccaaaaat tggtccttat ccagattatt cttcctcctc atacattgaa 2040
gactctatcc atcatactga cagcatcgtg aagaatatTT cctctgagca ttctatgtcc 2100
agcacacctt tgactatagg ggaaaaaaac cgaaattcaa ttaactatga acgacagcaa 2160
gcacaagctc gaatccccag ccctgaaaca agtgtcacca gcctctccac caacacaaca 2220
accacaaaca ccacaggact cacgccaagt actggcatga ctactatata tgagatgcca 2280
taccagatg aaacaaatct gcataccaca aatgttgcac agtcaattgg gccaacccct 2340
gtctgcttac agctgacaga agaagacttg gaaaccaaca agctagaccc aaaagaagtt 2400
gataagaacc tcaaggaaag ctctgatgag aatctcatgg agcactctct taaacagttc 2460
agtggcccgag acccactgag cagtactagt tctagcttgc tttacccact cataaaactt 2520
gcagtagaag caactggaca gcaggacttc acacagactg caaatggcca agcatgtttg 2580
attcctgatg ttctgcctac tcagatctat cctctcccca agcagcagaa ccttcccaag 2640
agacctacta gtttgccttt gaacacaaa aattcaacaa aagagccccg gctaaaaatt 2700
ggcagcaagc acaaaatctt cttgaaacaa gtgcgaaactg gagttgcca gatgaataca 2760
atcaatgcag cagaacctca tgtggtgaca gtcaccatga atgggtgtggc aggtagaaac 2820
cacagtgtta actcccatgc tgccacaacc caatatgcca ataggacagt actatctggc 2880
caaacaacca acatagtgac acataggggc caagaaatgt tgcagaatca gtttattggg 2940
gaggacaccc ggtgtaatat taattccagt cctgatgagc atgagccttt actgagacga 3000
gagcaacaag ctggccatga tgaagggtgt ctggatcgtc ttgtggacag gagggaacgg 3060
ccactagaag gtggccgaac taattccaat aacaacaaca gcaatccatg ttcagaacaa 3120
gatgttcttg cacagggtgt tccaagcaca gcagcagatc ctgggccatc aaagcccaga 3180
agagcacaga ggccaatctc tctggatctt tcagccacaa atgtcctgga tggcagcagt 3240
atacagatag gtgagtcaac acaagatggc aaatcaggat caggtgaaaa gatcaagaaa 3300
cgtgtgaaaa ctccctatct tcttaagcgg tggcgcccct ccacctgggt catctccact 3360
gaatcgctgg actgtgaagt caacaataat ggcagtaaca gggcagttca ttccaaatcc 3420
agcactgctg tttaccttgc agaaggaggc actgctacaa ccatgggtgtc taaagatata 3480
ggaatgaact gtctgtgaaa tgttttcaag cctatggagt gaaattattt ttgcatcat 3540
ttaaacatgc agaagatgtt tacggggcgg ggtgacagga gagagcgtca gcggcaagct 3600
gtggaggatg gggctcagaa tgcagacctg ggctggccgc atggcctctc cctgagccct 3660
gatttgtggg agggaagcag tatgggtgca gtccctcctc aggcctccct ctgggggtccc 3720
ccgatcctat ccacactctt cagggtgagc cagcctcacc tcttcttagt cctgaggggtg 3780
agggcaggct gaggcaacga gtgggaggtt caacaagag tgggctggag ccaagggaaa 3840
atagagatga tgtaatttct ttccggaatt c 3871

```

&lt;210&gt; 95

&lt;211&gt; 88

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 95

```

Cys Arg Glu Leu His Phe Thr Arg Tyr Val Thr Asp Gly Pro Cys Arg
 1           5           10          15
Ser Ala Lys Pro Val Thr Glu Leu Val Cys Ser Gly Gln Cys Gly Pro
          20          25          30
Ala Arg Leu Leu Pro Asn Ala Ile Gly Arg Gly Lys Trp Trp Arg Pro
          35          40          45
Ser Gly Pro Asp Phe Arg Cys Ile Pro Asp Arg Tyr Arg Ala Gln Arg
          50          55          60
Val Gln Leu Leu Cys Pro Gly Gly Glu Ala Pro Arg Ala Arg Lys Val
          65          70          75          80
Arg Leu Val Ala Ser Cys Lys Cys
          85

```

&lt;210&gt; 96

54

<211> 82  
 <212> PRT  
 <213> Homo sapiens

<400> 96  
 Cys Arg Pro Ile Asn Ala Thr Leu Ala Val Glu Lys Glu Gly Cys Pro  
 1 5 10 15  
 Val Cys Ile Thr Val Asn Thr Thr Ile Cys Ala Gly Tyr Cys Pro Thr  
 20 25 30  
 Met Thr Arg Val Leu Gln Gly Val Leu Pro Ala Leu Pro Gln Val Val  
 35 40 45  
 Cys Asn Tyr Arg Asp Val Arg Phe Glu Ser Ile Arg Leu Pro Gly Cys  
 50 55 60  
 Pro Arg Gly Val Asn Pro Val Val Ser Tyr Ala Val Ala Leu Ser Cys  
 65 70 75 80  
 Gln Cys

<210> 97  
 <211> 82  
 <212> PRT  
 <213> Homo sapiens

<400> 97  
 Cys Glu Leu Thr Asn Ile Thr Ile Ala Ile Glu Lys Glu Glu Cys Arg  
 1 5 10 15  
 Phe Cys Ile Ser Ile Asn Thr Thr Trp Cys Ala Gly Tyr Cys Tyr Thr  
 20 25 30  
 Arg Asp Leu Val Tyr Lys Asp Pro Ala Arg Pro Lys Ile Gln Lys Thr  
 35 40 45  
 Cys Thr Phe Lys Glu Leu Val Tyr Glu Thr Val Arg Val Pro Gly Cys  
 50 55 60  
 Ala His His Ala Asp Ser Leu Tyr Thr Tyr Pro Val Ala Thr Gln Cys  
 65 70 75 80  
 His Cys

<210> 98  
 <211> 84  
 <212> PRT  
 <213> Homo sapiens

<400> 98  
 Cys Ile Pro Thr Glu Tyr Thr Met His Ile Glu Arg Arg Glu Cys Ala  
 1 5 10 15  
 Tyr Cys Leu Thr Ile Asn Thr Thr Ile Cys Ala Gly Tyr Cys Met Thr  
 20 25 30  
 Arg Asp Ile Asn Gly Lys Leu Phe Leu Pro Lys Tyr Ala Leu Ser Gln  
 35 40 45  
 Asp Val Cys Thr Tyr Arg Asp Phe Ile Tyr Arg Thr Val Glu Ile Pro  
 50 55 60  
 Gly Cys Pro Leu His Val Ala Pro Tyr Phe Ser Tyr Pro Val Ala Leu  
 65 70 75 80  
 Ser Cys Lys Cys



55

<210> 99  
 <211> 83  
 <212> PRT  
 <213> Homo sapiens

<400> 99  
 Cys Asn Asp Ile Thr Ala Arg Leu Gln Tyr Val Lys Val Gly Ser Cys  
 1 5 10 15  
 Lys Ser Glu Val Glu Val Asp Ile His Tyr Cys Gln Gly Lys Cys Ala  
 20 25 30  
 Ser Lys Ala Met Tyr Ser Ile Asp Ile Asn Asp Val Gln Asp Gln Cys  
 35 40 45  
 Ser Cys Cys Ser Pro Thr Arg Thr Glu Pro Met Gln Val Ala Leu His  
 50 55 60  
 Cys Thr Asn Gly Ser Val Val Tyr His Glu Val Leu Asn Ala Met Glu  
 65 70 75 80  
 Cys Lys Cys

<210> 100  
 <211> 80  
 <212> PRT  
 <213> Homo sapiens

<400> 100  
 Cys Ser Thr Val Pro Val Thr Thr Glu Val Ser Tyr Ala Gly Cys Thr  
 1 5 10 15  
 Lys Thr Val Leu Met Asn His Cys Ser Gly Ser Cys Gly Thr Phe Val  
 20 25 30  
 Met Tyr Ser Ala Lys Ala Gln Ala Leu Asp His Ser Cys Ser Cys Cys  
 35 40 45  
 Lys Glu Glu Lys Thr Ser Gln Arg Glu Val Val Leu Ser Cys Pro Asn  
 50 55 60  
 Gly Gly Ser Leu Thr His Thr Tyr Thr His Ile Glu Ser Cys Gln Cys  
 65 70 75 80

<210> 101  
 <211> 80  
 <212> PRT  
 <213> Homo sapiens

<400> 101  
 Cys Arg Thr Val Pro Phe Ser Gln Thr Ile Thr His Glu Gly Cys Glu  
 1 5 10 15  
 Lys Val Val Val Gln Asn Asn Leu Cys Phe Gly Lys Cys Gly Ser Val  
 20 25 30  
 His Phe Pro Gly Ala Ala Gln His Ser His Thr Ser Cys Ser His Cys  
 35 40 45  
 Leu Pro Ala Lys Phe Thr Thr Met His Leu Pro Leu Asn Cys Thr Glu  
 50 55 60  
 Leu Ser Ser Val Ile Lys Val Val Met Leu Val Glu Glu Cys Gln Cys  
 65 70 75 80

<210> 102

56

<211> 85  
 <212> PRT  
 <213> Homo sapiens

<400> 102

```

Cys Lys Thr Gln Pro Leu Lys Gln Thr Ile His Glu Glu Gly Cys Asn
 1           5           10           15
Ser Arg Thr Ile Ile Asn Arg Phe Cys Tyr Gly Gln Cys Asn Ser Phe
           20           25           30
Tyr Ile Pro Arg His Ile Arg Lys Glu Glu Gly Ser Phe Gln Ser Cys
           35           40           45
Ser Phe Cys Lys Pro Lys Lys Phe Thr Thr Met Met Val Thr Leu Asn
 50           55           60
Cys Pro Glu Leu Gln Pro Pro Thr Lys Lys Lys Arg Val Thr Arg Val
65           70           75           80
Lys Gln Cys Arg Cys
           85

```

<210> 103  
 <211> 86  
 <212> PRT  
 <213> Homo sapiens

<400> 103

```

Cys Glu Ala Lys Asn Ile Thr Gln Ile Val Gly His Ser Gly Cys Glu
 1           5           10           15
Ala Lys Ser Ile Gln Asn Arg Ala Cys Leu Gly Gln Cys Phe Ser Tyr
           20           25           30
Ser Val Pro Asn Thr Phe Pro Gln Ser Thr Glu Ser Leu Val His Cys
           35           40           45
Asp Ser Cys Met Pro Ala Gln Ser Met Trp Glu Ile Val Thr Leu Glu
 50           55           60
Cys Pro Gly His Glu Glu Val Pro Arg Val Asp Lys Leu Val Glu Lys
65           70           75           80
Ile Leu His Cys Ser Cys
           85

```

<210> 104  
 <211> 70  
 <212> PRT  
 <213> Homo sapiens

<400> 104

```

Cys Ile Arg Thr Pro Lys Ile Ser Lys Pro Ile Lys Phe Glu Leu Ser
 1           5           10           15
Gly Cys Thr Ser Met Lys Thr Tyr Arg Ala Lys Phe Cys Gly Val Cys
           20           25           30
Thr Asp Gly Arg Cys Cys Thr Pro His Arg Thr Thr Thr Leu Pro Val
           35           40           45
Glu Phe Lys Cys Pro Asp Gly Glu Val Met Lys Lys Asn Met Met Phe
 50           55           60
Ile Lys Thr Cys Ala Cys
65           70

```

<210> 105

57

<211> 70  
 <212> PRT  
 <213> Homo sapiens

<400> 105  
 Cys Leu Arg Thr Lys Lys Ser Leu Lys Ala Ile His Leu Gln Phe Lys  
 1 5 10 15  
 Asn Cys Thr Ser Leu His Thr Tyr Lys Pro Arg Phe Cys Gly Val Cys  
 20 25 30  
 Ser Asp Gly Arg Cys Cys Thr Pro His Asn Thr Lys Thr Ile Gln Ala  
 35 40 45  
 Glu Phe Gln Cys Ser Pro Gly Gln Ile Val Lys Lys Pro Val Met Val  
 50 55 60  
 Ile Gly Thr Cys Thr Cys  
 65 70

<210> 106  
 <211> 70  
 <212> PRT  
 <213> Homo sapiens

<400> 106  
 Cys Ser Lys Thr Lys Lys Ser Pro Glu Pro Val Arg Phe Thr Tyr Ala  
 1 5 10 15  
 Gly Cys Leu Ser Val Lys Lys Tyr Arg Pro Lys Tyr Cys Gly Ser Cys  
 20 25 30  
 Val Asp Gly Arg Cys Cys Thr Pro Gln Leu Thr Arg Thr Val Lys Met  
 35 40 45  
 Arg Phe Arg Cys Glu Asp Gly Glu Thr Phe Ser Lys Asn Val Met Met  
 50 55 60  
 Ile Gln Ser Cys Lys Cys  
 65 70

<210> 107  
 <211> 205  
 <212> PRT  
 <213> Homo sapiens

<400> 107  
 Gln His Tyr Leu His Ile Arg Pro Ala Pro Ser Asp Asn Leu Pro Leu  
 1 5 10 15  
 Val Asp Leu Ile Glu His Pro Asp Pro Ile Phe Asp Pro Lys Glu Lys  
 20 25 30  
 Asp Leu Asn Glu Thr Leu Leu Arg Ser Leu Leu Gly Gly His Tyr Asp  
 35 40 45  
 Pro Gly Phe Met Ala Thr Ser Pro Pro Glu Asp Arg Pro Gly Gly Gly  
 50 55 60  
 Gly Gly Ala Ala Gly Gly Ala Glu Asp Leu Ala Glu Leu Asp Gln Leu  
 65 70 75 80  
 Leu Arg Gln Arg Pro Ser Gly Ala Met Pro Ser Glu Ile Lys Gly Leu  
 85 90 95  
 Glu Phe Ser Glu Gly Leu Ala Gln Gly Lys Lys Gln Arg Leu Ser Lys  
 100 105 110  
 Lys Leu Arg Arg Lys Leu Gln Met Trp Leu Trp Ser Gln Thr Phe Cys  
 115 120 125  
 Pro Val Leu Tyr Ala Trp Asn Asp Leu Gly Ser Arg Phe Trp Pro Arg

58

130		135		140
Tyr Val Lys Val Gly Ser Cys Phe Ser Lys Arg Ser Cys Ser Val Pro				
145		150		155
Glu Gly Met Val Cys Lys Pro Ser Lys Ser Val His Leu Thr Val Leu				
	165		170	175
Arg Trp Arg Cys Gln Arg Arg Gly Gly Gln Arg Cys Gly Trp Ile Pro				
	180		185	190
Ile Gln Tyr Pro Ile Ile Ser Glu Cys Lys Cys Ser Cys				
	195	200		205

&lt;210&gt; 108

&lt;211&gt; 197

&lt;212&gt; PRT

&lt;213&gt; Gallus gallus

&lt;400&gt; 108

Gln His Tyr Leu His Ile Arg Pro Ala Pro Ser Asp Asn Leu Pro Leu				
1	5	10	15	
Val Asp Leu Ile Glu His Pro Asp Pro Ile Phe Asp Pro Lys Glu Lys				
	20	25	30	
Asp Leu Asn Glu Thr Leu Leu Arg Ser Leu Met Gly Gly His Phe Asp				
	35	40	45	
Pro Asn Phe Met Ala Met Ser Leu Pro Glu Asp Arg Leu Gly Val Asp				
	50	55	60	
Asp Leu Ala Glu Leu Asp Leu Leu Leu Arg Gln Arg Pro Ser Gly Ala				
	65	70	75	80
Met Pro Gly Glu Ile Lys Gly Leu Glu Phe Tyr Asp Gly Leu Gln Pro				
	85	90	95	
Gly Lys Lys His Arg Leu Ser Lys Lys Leu Arg Arg Lys Leu Gln Met				
	100	105	110	
Trp Leu Trp Ser Gln Thr Phe Cys Pro Val Leu Tyr Thr Trp Asn Asp				
	115	120	125	
Leu Gly Ser Arg Phe Trp Pro Arg Tyr Val Lys Val Gly Ser Cys Tyr				
	130	135	140	
Ser Lys Arg Ser Cys Ser Val Pro Glu Gly Met Val Cys Lys Pro Ala				
	145	150	155	160
Lys Ser Val His Leu Thr Ile Leu Arg Trp Arg Cys Gln Arg Arg Gly				
	165	170	175	
Gly Gln Arg Cys Thr Trp Ile Pro Ile Gln Tyr Pro Ile Ile Ala Glu				
	180	185	190	
Cys Lys Cys Ser Cys				
	195			

&lt;210&gt; 109

&lt;211&gt; 196

&lt;212&gt; PRT

&lt;213&gt; Xenopus laevis

&lt;400&gt; 109

Gln His Tyr Leu His Ile Arg Pro Ala Pro Ser Glu Asn Leu Pro Leu				
1	5	10	15	
Val Asp Leu Ile Glu His Pro Asp Pro Ile Tyr Asp Pro Lys Glu Lys				
	20	25	30	
Asp Leu Asn Glu Thr Leu Leu Arg Thr Leu Met Val Gly His Phe Asp				
	35	40	45	
Pro Asn Phe Met Ala Thr Ile Leu Pro Glu Glu Arg Leu Gly Val Glu				

59

50		55		60	
Asp	Leu	Gly	Glu	Leu	Asp
65					70
Met	Pro	Ala	Glu	Ile	Lys
					85
Lys	Lys	His	Arg	Leu	Ser
					100
Leu	Trp	Ser	Gln	Thr	Phe
					115
Gly	Thr	Arg	Phe	Trp	Pro
					130
Lys	Arg	Ser	Cys	Ser	Val
145					150
Ser	Met	His	Leu	Thr	Ile
					165
Gln	Lys	Cys	Ala	Trp	Ile
					180
Lys	Cys	Ser	Cys		
					195

<210> 110  
 <211> 195  
 <212> PRT  
 <213> Takifugu rubripes

<400> 110
Gln Pro Tyr Tyr Leu Leu Arg Pro Ile Pro Ser Asp Ser Leu Pro Ile
1 5 10 15
Val Glu Leu Lys Glu Asp Pro Gly Pro Val Phe Asp Pro Lys Glu Arg
20 25 30
Asp Leu Asn Glu Thr Glu Leu Lys Ser Val Leu Gly Asp Phe Asp Ser
35 40 45
Arg Phe Leu Ser Val Leu Pro Ala Glu Asp Gly His Ala Gly Asn
50 55 60
Asp Glu Leu Asp Asp Phe Asp Ala Gln Arg Trp Gly Gly Ala Leu Pro
65 70 75 80
Lys Glu Ile Arg Ala Val Asp Phe Asp Ala Pro Gln Leu Gly Lys Lys
85 90 95
His Lys Pro Ser Lys Lys Leu Lys Arg Arg Leu Gln Gln Trp Leu Trp
100 105 110
Ala Tyr Ser Phe Cys Pro Leu Ala His Ala Trp Thr Asp Leu Gly Ser
115 120 125
Arg Phe Trp Pro Arg Phe Val Arg Ala Gly Ser Cys Leu Ser Lys Arg
130 135 140
Ser Cys Ser Val Pro Glu Gly Met Thr Cys Lys Pro Ala Thr Ser Thr
145 150 155 160
His Leu Thr Ile Leu Arg Trp Arg Cys Val Gln Arg Lys Val Gly Leu
165 170 175
Lys Cys Ala Trp Ile Pro Met Gln Tyr Pro Val Ile Thr Asp Cys Lys
180 185 190
Cys Ser Cys
195

<210> 111  
 <211> 196  
 <212> PRT

&lt;213&gt; Danio rerio

&lt;400&gt; 111

Gln His Tyr Tyr Leu Leu Arg Pro Ile Pro Ser Asp Ser Leu Pro Ile  
 1 5 10 15  
 Val Glu Leu Lys Glu Asp Pro Asp Pro Val Leu Asp Pro Lys Glu Arg  
 20 25 30  
 Asp Leu Asn Glu Thr Glu Leu Arg Ala Ile Leu Gly Ser His Phe Glu  
 35 40 45  
 Gln Asn Phe Met Ser Ile Asn Pro Pro Glu Asp Lys His Ala Gly Gln  
 50 55 60  
 Asp Glu Leu Asn Glu Ser Glu Leu Met Lys Gln Arg Pro Asn Gly Ile  
 65 70 75 80  
 Met Pro Lys Glu Ile Lys Ala Met Glu Phe Asp Ile Gln His Gly Lys  
 85 90 95  
 Lys His Lys Pro Ser Lys Lys Leu Arg Arg Arg Leu Gln Leu Trp Leu  
 100 105 110  
 Trp Ser Tyr Thr Phe Cys Pro Val Val His Thr Trp Gln Asp Leu Gly  
 115 120 125  
 Asn Arg Phe Trp Pro Arg Tyr Leu Lys Val Gly Ser Cys Tyr Asn Lys  
 130 135 140  
 Arg Ser Cys Ser Val Pro Glu Gly Met Val Cys Lys Pro Pro Lys Ser  
 145 150 155 160  
 Ser His Leu Thr Val Leu Arg Trp Arg Cys Val Gln Arg Lys Gly Gly  
 165 170 175  
 Leu Lys Cys Ala Trp Ile Pro Val Gln Tyr Pro Val Ile Ser Glu Cys  
 180 185 190  
 Lys Cys Ser Cys  
 195

&lt;210&gt; 112

&lt;211&gt; 188

&lt;212&gt; PRT

&lt;213&gt; Mus musculus

&lt;400&gt; 112

Gln Gly Trp Gln Ala Phe Arg Asn Asp Ala Thr Glu Val Ile Pro Gly  
 1 5 10 15  
 Leu Gly Glu Tyr Pro Glu Pro Pro Pro Glu Asn Asn Gln Thr Met Asn  
 20 25 30  
 Arg Ala Glu Asn Gly Gly Arg Pro Pro His His Pro Tyr Asp Ala Lys  
 35 40 45  
 Gly Val Ser Glu Tyr Ser Cys Arg Glu Leu His Tyr Thr Arg Phe Leu  
 50 55 60  
 Thr Asp Gly Pro Cys Arg Ser Ala Lys Pro Val Thr Glu Leu Val Cys  
 65 70 75 80  
 Ser Gly Gln Cys Gly Pro Ala Arg Leu Leu Pro Asn Ala Ile Gly Arg  
 85 90 95  
 Val Lys Trp Trp Arg Pro Asn Gly Pro Asp Phe Arg Cys Ile Pro Asp  
 100 105 110  
 Arg Tyr Arg Ala Gln Arg Val Gln Leu Leu Cys Pro Gly Gly Ala Ala  
 115 120 125  
 Pro Arg Ser Arg Lys Val Arg Leu Val Ala Ser Cys Lys Cys Lys Arg  
 130 135 140  
 Leu Thr Arg Phe His Asn Gln Ser Glu Leu Lys Asp Phe Gly Pro Glu  
 145 150 155 160  
 Thr Ala Arg Pro Gln Lys Gly Arg Lys Pro Arg Pro Gly Ala Arg Gly

61

Ala Lys Ala Asn Gln Ala Glu Leu Glu Asn Ala Tyr  
165 170  
180 185

175